


PM 21

100-607

1 of 2



SUPPLEMENTAL LABELING

RIDOMIL® 2E FUNGICIDE  
EPA REG. NO. 100-607

STRAWBERRIES

Active Ingredient:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester .....	25.1%
Inert Ingredients:	74.9%
Total:	100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

PRECAUCIÓN AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

All applicable directions, restrictions, and precautions on the EPA-registered Ridomil 2E label are to be followed.

This label must be in the possession of the user at the time of pesticide application.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY AND/OR ILLEGAL RESIDUES.

Strawberries

Ridomil provides control of red stele (Phytophthora fragariae), vascular collapse (P. cactorum) and leather rot (P. cactorum) when used as directed. Applications may be made using ground application equipment or through irrigation systems.

Annual Strawberries (Vascular Collapse): Apply Ridomil 2E at 2 qts./treated acre in sufficient water to move the fungicide into the root zone of the plants. Make one application at the time of transplanting. Make an additional application 30 days before the start of harvest and another one in July before the second flush. For banded applications, a 12-inch band is recommended. Use the formula in the General Information section of this label to determine the amount of Ridomil needed per acre. When Ridomil is applied through drip irrigation systems, use the same amount as would be applied in a 12-inch band. Observe the precautions concerning application of Ridomil through irrigation systems in the General Information section of this label.

Perennial Strawberries (Red Stele and Leather Rot): Apply Ridomil 2E at 2 qts./treated acre in sufficient water to move the fungicide into the root zone of the plants. Make one application at the time of transplanting or in the spring before the plants start growth. Make another application in the fall after harvest. For control of leather rot, an application can be made during the growing season. Use sufficient water to move the Ridomil into the root zone. For banded applications, a 12-inch band is recommended. Use the formula in the General Information section of this label to determine the amount of Ridomil needed per acre.

Note: To avoid possible illegal residues, do not use more than a total of 6 qts. (3 lbs. ai) of Ridomil 2E per acre of strawberries per year.

<u>Rotational Crops</u>	<u>Planting Time From Last Ridomil 2E Application</u>
Strawberries	-0- days

Ridomil® trademark of CIBA-GEIGY Corporation for metalaxyl.

©1989 CIBA-GEIGY Corporation

Agricultural Division  
 CIBA-GEIGY Corporation  
 Greensboro, North Carolina 27419

CGA

(LABELS-R.1 - RDML2/SUP/F3)

November 4, 1988  
 Revised September 26, 1989