

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

100-617

3-30-2001

1/7



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number  
NOTIFICATION

**Application for Pesticide - Section I**

1. Company/Product Number 100-617	2. EPA Product Manager Mary Waller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Tilt@ Bravo@Twin-Pak™ Fungicide	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419  <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____ Product Name _____	

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b>  MAR 30 2001
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

**Explanation: Use additional page(s) if necessary. (For Section I and Section II.)**

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. The following changes are being made via this notification: 1) Company name and address have been updated to reflect Syngenta Crop Protection, Inc. 2) The Conditions of Sale and Warranty statement has been changed to reflect the name change. Because Syngenta has been formed by the merger of Novartis Crop Protection, Inc. and Zeneca Ag Products, we have chosen to use the former Zeneca warranty statement as the Syngenta warranty statement. No other changes occur in the statement other than the name change. 3) The copyright date reflects Syngenta. 4) Trademark statements have been updated to reflect Syngenta for those products for which Syngenta holds the trademark. 5) The Internet address has been changed to reflect Syngenta. 6) Other places in the label which referring to the company name have been updated.

**Section - III**

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal <input type="checkbox"/>		Plastic <input type="checkbox"/>
*Certification must be submitted		If "Yes" No. per Unit Packaging wgt. Container		Glass <input type="checkbox"/>	
If "Yes" No. per Unit Packaging wgt. Container		If "Yes" No. per Unit Packaging wgt. container		Paper <input type="checkbox"/>	Other (Specify) _____
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					


**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Nan S. Padgett		Title Label Group Leader		Telephone No. (Includc Area Code) 336-632-7567	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
4. Typed Name Nan S. Padgett		5. Date March 19, 2001			

MAR 30 2001

PULL HERE TO OPEN ►

# TILT®



## BRAVO®

### FUNGICIDE

For Control of Certain Diseases in Peanuts

See additional precautionary statements and directions for use inside booklet.

Using label instructions, apply the entire contents of this container to the target crop once the container is opened.

EPA Est. 34704-MS-2

SCP 617C-M2C 0201

Total **180 fl. oz.**  
U.S. Standard Measure

#### Tilt

Active Ingredient:	
Propiconazole (CAS No. 60207-90-1) . . .	41.8%
Other Ingredients:	58.2%
Total:	100.0%

EPA Reg. No. 100-617  
Product of Switzerland  
Formulated in the USA

**KEEP OUT OF REACH OF CHILDREN.**

**WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

20 fl. OZ. U.S. Standard Measure

#### Bravo Weather Stik®

Active Ingredient:	
Chlorothalonil (CAS No. 1897-45-6) . . .	54.0%
Other Ingredients:	46.0%
Total:	100.0%

Contains 6.0 lbs. chlorothalonil per gallon (720 grams per liter)

EPA Reg. No. 50534-188-10182

**KEEP OUT OF REACH OF CHILDREN.**

**WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

160 fl. OZ. U.S. Standard Measure

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

#### Tilt/Bravo

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

## GENERAL INFORMATION

Tilt/Bravo is a fungicide for the control of certain diseases in peanuts. Using label instructions, apply the entire contents of this container to the target crop once the container is opened.

**Important:** Do not use in greenhouses, nurseries, on turf, landscape plantings, or as a tree injection.

**FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.**

### Spray Equipment

Thorough coverage is necessary to provide good disease control.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate sprayer before use.

Use a pump with capacity to: (1) maintain 35-40 psi at nozzles, and (2) provide sufficient, **continuous** agitation in tank to keep mixture in suspension – this requires recirculation of 10% of tank volume per minute. Use a jet agitator or liquid sparge tube for agitation. Do not air sparge.

It is suggested that screens be used to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be *16-mesh or coarser*. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.

**Aerial Application:** Apply in a minimum of 5 gals. of water per acre. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals.

**Ground Application:** Apply Tilt/Bravo by ground equipment in a minimum of 15 gals. of water per acre.

**Chemigation:** Do not apply this product through any type of irrigation system.

### Mixing Instructions

**Note:** Slowly invert container several times to assure uniform mixture.

Prepare no more spray mixture than is required for the immediate operation. Use the entire contents of this container once the container is opened. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

**Tilt/Bravo Alone:** Add  $1/2$ - $2/3$  of the required amount of water to the spray or mixing tank. With the agitator running, add the Tilt/Bravo to the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the Tilt/Bravo has completely dispersed into the water. Maintain continuous agitation until all of the mixture has been sprayed.

**Tilt/Bravo + Tank Mixtures:** Consult your dealer or Syngenta sales representative for information concerning compatibility with other tank mix partners. **Note: When using Tilt/Bravo in mixture, any product in water-soluble packaging should be added to the tank prior to adding Tilt/Bravo. Allow water-soluble packaging to completely dissolve and the mixture product to completely disperse before adding Tilt/Bravo to the tank.**

Add  $1/2$ - $2/3$  of the required amount of water to the spray or mixing tank. Again, if tank mixing with any product packaged in water-soluble packets, add this product first with the agitator running. Allow the material to completely dissolve and disperse in the mix water. Continue agitation while adding all of the contents of both chambers of the Tilt/Bravo container at once. In general, additional tank mix partners should be added in the following order: wettable powders, dry flowable formulations, liquid flowable formulations, and emulsifiable concentrations. Always allow each tank mix partner to become fully dispersed before adding the next product. Maintain agitation while adding the rest of the water, while adding the other tank mix partners and throughout the mixing and spraying operation.

If using Tilt/Bravo in mixture with other products, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate should be exceeded and the most restrictive label precautions and limitations should be followed. Tilt/Bravo should not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products are permitted only in those states in which the products are registered.

**Important:** Do not combine Tilt/Bravo with Dipel® 4L, Triton AG-98, or Triton B-1956, as phytotoxicity may result.

### Rotational Crops

To avoid illegal residues, do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of Tilt/Bravo application to the preceding crop, unless the second crop is one of the following: Barley, Celery, Grasses Grown for Seed, Rice, Rye, or Wheat.

### Peanuts

For control of early leafspot (*Cercospora*), late leafspot (*Cercosporidium*), and web blotch (*Phoma*), use the entire contents of this container to treat 10 acres of peanuts. Begin application 30-40 days after planting or at the first appearance of disease; continue applications on a 10 to 14-day schedule. Under conditions highly favorable for disease development use the shorter spray interval.

**Notes:** (1) Do not apply within 14 days of harvest. (2) Do not apply more than 8 applications of Tilt/Bravo per season. (3) Do not exceed the equivalent of 16 fl. oz./A per season of Tilt, whether used alone or as a component of Tilt/Bravo. (4) Do not allow livestock to graze in treated areas. (5) Do not feed hay or threshings from treated fields to livestock. (6) Using label instructions, apply the entire contents of this container to the target crop once the container is opened.

## STORAGE AND DISPOSAL

### Storage Instructions

Store in the original container. Using label instructions, apply the entire contents of this container to the target crop once the container is opened. Store in a cool place. Protect from excessive heat. Keep from freezing.

### Pesticide Disposal

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

### Container Disposal

Do not reuse empty container. Triple rinse container (or equivalent) and then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. Stay out of smoke from burning containers.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

## PRECAUTIONARY STATEMENTS

### Tilt

#### Hazards to Humans and Domestic Animals

##### WARNING/AVISO

This product may cause skin sensitization reactions in certain individuals.

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing spray mist.

#### Statement of Practical Treatment

**If in eyes:** Flush with plenty of water. Get medical attention if irritation persists.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**If swallowed:** Do not induce vomiting. Promptly drink a large quantity of milk, egg whites, or gelatin solution. If these are not available, drink large quantities of water. Never give anything by mouth to an unconscious person. Call a physician or Poison Control Center immediately.

**If inhaled:** Move victim to fresh air.

**Note to Physician:** There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give a saline laxative and supportive therapy.

### Bravo Weather Stik

#### Hazards to Humans and Domestic Animals

##### WARNING/AVISO

Causes eye irritation. May cause skin irritation. May be a potential skin sensitizer. Do not get in eyes. Avoid prolonged contact with skin. Avoid breathing spray mist. Do not take internally.

**Note to User:** This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation, and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

#### Statement of Practical Treatment

**If in eyes:** Immediately flush eyes with plenty of water and continue for 15 minutes. Seek medical attention for eyes immediately.

**If on skin:** Wash exposed areas of skin with soap and warm water after handling or using.

**If swallowed:** Never give anything by mouth to an unconscious person.

**Note to Physician:** Persons having a temporary allergic reaction respond to treatment with antihistamines or steroid creams and/or system steroids.

## Tilt/Bravo

### Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category G on an EPA chemical resistance category selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear
- Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**Environmental Hazards**

**Tilt**

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

**Bravo Weather Stik**

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Do not apply when weather conditions favor drift from treated areas. Apply only to areas specified on label.

**Physical or Chemical Hazards**

Do not use, pour, spill, or store near heat, open flame, or hot surfaces.

---

Tilt® and Twin-Pak™ trademarks of Syngenta  
U.S. Patent No. 4,079,062

Bravo® Weather Stik® trademark of Zeneca Ag Products

Dipel® trademark of Abbott Laboratories

Triton AG-98 and Triton B-1956 trademarks of Rohm and Haas Company

©2001 Syngenta


Tilt:

Product of Switzerland  
Formulated in the USA

Syngenta Crop Protection, Inc.  
Greensboro, North Carolina 27409  
[www.syngenta-us.com](http://www.syngenta-us.com)

**SCP 617C-M2C 0201**

# TILT® BRAVO®



**Twin-Pak**

## FUNGICIDE

For Control of Certain  
Diseases in Peanuts

EPA Est. 34704-MS-2

©2001 Syngenta  
Syngenta Crop Protection, Inc.  
Greensboro, NC 27409  
www.syngenta-us.com  
**SCP 617C-M2C 0201**

Total **180 fl. oz.**  
U.S. Standard Measure

### Tilt

Active Ingredient:	
Propiconazole (CAS No. 60207-90-1)	41.8%
Other Ingredients:	58.2%
Total:	100.0%

EPA Reg. No. 100-617

Tilt® and Twin-Pak™ trademarks of Syngenta  
U.S. Patent No. 4,079,062

**20 fl. oz.**  
U.S. Standard Measure

### Bravo Weather Stik®

Active Ingredient:	
Chlorothalonil (CAS No. 1897-45-6)	54.0%
Other Ingredients:	46.0%
Total:	100.0%

Contains 6.0 lbs. chlorothalonil per gallon (720 grams per liter)

EPA Reg. No. 50534-188-10182

Bravo® Weather Stik® trademark of Zeneca Ag Products

**160 fl. oz.**  
U.S. Standard Measure

See directions for use in attached booklet.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.



**NEXT**

**LABEL**

100-617

3-30-2001

1/10

 <b>United States</b> <b>Environmental Protection Agency</b> Washington, DC 20460	<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width:30%;"><input type="checkbox"/> Registration</td> </tr> <tr> <td><input type="checkbox"/> Amendment</td> </tr> <tr> <td><input checked="" type="checkbox"/> Other</td> </tr> </table>	<input type="checkbox"/> Registration	<input type="checkbox"/> Amendment	<input checked="" type="checkbox"/> Other	OPP Identifier Number <b>NOTIFICATION</b>
<input type="checkbox"/> Registration					
<input type="checkbox"/> Amendment					
<input checked="" type="checkbox"/> Other					

**Application for Pesticide - Section I**

1. Company/Product Number 100-617	2. EPA Product Manager Mary Waller	3. Proposed Classification  <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Tilt® Fungicide	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419  <input type="checkbox"/> Check if this is a new address		6. <b>Expedited Review.</b> In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____  Product Name _____

**Section - II**

<input type="checkbox"/> Amendment - Explain below.  <input type="checkbox"/> Resubmission in response to Agency letter dated _____  <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application.  <input type="checkbox"/> Other - Explain below.
--	---

NOTIFICATION  
MAR 30 2001


**Explanation:** Use additional page(s) if necessary. (For Section I and Section II.)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. The following changes are being made via this notification: 1) Company name and address have been updated to reflect Syngenta Crop Protection, Inc. 2) The Conditions of Sale and Warranty statement has been changed to reflect the name change. Because Syngenta has been formed by the merger of Novartis Crop Protection, Inc. and Zeneca Ag Products, we have chosen to use the former Zeneca warranty statement as the Syngenta warranty statement. No other changes occur in the statement other than the name change. 3) The copyright date reflects Syngenta. 4) Trademark statements have been updated to reflect Syngenta for those products for which Syngenta holds the trademark. 5) The Internet address has been changed to reflect Syngenta. 6) Other places in the label which referring to the company name have been updated.

**Section - III**

<b>1. Material This Product Will Be Packaged In:</b>				<b>2. Type of Container</b>	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
<i>*Certification must be submitted</i> If "Yes" Unit Packaging wgt.    No. per Container		If "Yes" Unit Packaging wgt.    No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

**Section - IV**

<b>1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)</b>		
Name Nan S. Padgett	Title Label Group Leader	Telephone No. (Include Area Code) 336-632-7567
<b>Certification</b>		<b>6. Date Application Received (Stamped)</b>
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		_____ _____ _____
2. Signature 	3. Title Label Group Leader	
4. Typed Name Nan S. Padgett	5. Date March 19, 2001	

2/10

NOTIFICATION

MAR 30 2001

PULL HERE TO OPEN ►

# TILT®



## FUNGICIDE

For control of certain diseases in celery, cereals, corn, grasses grown for seed, peanuts, pineapple, and sugarcane

**KEEP OUT OF REACH OF CHILDREN. WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

Active Ingredient:	
Propiconazole (CAS No. 60207-90-1) . . . . .	41.8%
Other Ingredients:	58.2%
Total:	100.0%

SCP 617A-M2D 0201

EPA Reg. No. 100-617  
EPA Est. 34704-MS-2©  
EPA Est. 67545-AZ-1©  
(Superscript is first letter of lot number on jug)

**ONE GALLON**  
U.S. Standard Measure

Product of Switzerland  
Formulated in the USA



### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.** Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

## GENERAL INFORMATION

Tilt is a broad spectrum fungicide for the control of certain diseases in wheat, barley, rye, oats, rice, wild rice, celery, corn, peanuts, pineapple, sugarcane, and in grasses grown for seed.

**Important:** Do not use in nurseries, on turf, landscape plantings, or as a tree injection.

**FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.**

### Spray Equipment

Thorough coverage is necessary to provide good disease control.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate sprayer before use.

Use a pump with capacity to: (1) maintain 35-40 psi at nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension – this requires recirculation of 10% of tank volume per minute. Use a jet agitator or liquid sparge tube for agitation. Do not air sparge.

Although Tilt is an emulsifiable concentrate, it is suggested that screens be used to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be *16-mesh or coarser*. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.

### Application Instructions

**Aerial Application:** For those crops where aerial applications are indicated, apply in a minimum of 5 gals. of water per acre, unless specified otherwise. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals.

**Ground Application:** Apply Tilt by ground equipment in a minimum of 10 gals. of water per acre, unless specified otherwise.

### Application Through Irrigation Systems – Apply Tilt through irrigation equipment only to crops for which chemigation is specified on this label.

Tilt, alone or in combination with other pesticides which are registered for application through irrigation systems, may be applied through irrigation systems. Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**Note:** Do not inject Tilt at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part Tilt. Tilt is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used, but should be replaced once a year. Do not use Viton, Buna-N, Neoprene, or PVC seals.

### Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended.

### Center Pivot Irrigation Equipment

**Notes:** (1) Use only with drive systems which provide uniform water distribution. (2) Do not use end guns when chemigating Tilt through center pivot systems because of nonuniform application.

- Determine the size of the area to be treated.
- Determine the time required to apply  $1/8$ - $1/2$  inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. When applying Tilt through irrigation equipment, use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of Tilt required to treat the area covered by the irrigation system.
- Add the required amount of Tilt and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the Tilt solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the Tilt solution has cleared the sprinkler head.

### Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying Tilt through irrigation equipment, use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of Tilt required to treat the area covered by the irrigation system.

- Add the required amount of Tilt into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the Tilt solution has cleared the last sprinkler head.

### Mixing Instructions

Prepare no more spray mixture than is required for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

**Important: For wheat, barley, rye, and rice, custom applicators must use a closed system for mixing and loading this product.**

**Tilt Alone:** Add  $1/2$ - $2/3$  of the required amount of water to the spray or mixing tank. With the agitator running, add the Tilt to the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the Tilt has completely dispersed into the mix water. Maintain agitation until all of the mixture has been sprayed.

**Tilt + Tank Mixtures:** Tilt is usually compatible with all tank mix partners listed on this label. To determine the physical compatibility of Tilt with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Add  $1/2$ - $2/3$  of the required amount of water to the spray or mixing tank. With the agitator running, add the tank mix partner into the tank. Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and the Tilt to the spray tank. Allow the Tilt to completely disperse. Spray the mixture with the agitator running.

If using Tilt in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate should be exceeded, and the most restrictive label precautions and limitations should be followed. This product should not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

### Rotational Crops

Soybeans may be planted as a double crop following a cereal crop which has been treated with Tilt. Do not use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding.

To avoid possible illegal residues, do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of Tilt application to the preceding crop, unless the second crop appears on this label.

## CELERY

For control of early blight (*Cercospora*) and late blight (*Septoria*), apply 4 fl. oz. of Tilt per acre on a 7-day schedule, either by ground or aerial application. If desired, Tilt may be tank mixed with an appropriate spreader-sticker.

**Notes:** To avoid possible illegal residues, (1) Do not apply more than 16 fl. oz. of Tilt per crop (4 applications), and (2) Do not apply during the last 14 days before harvest.

## CEREALS

### Wheat, Barley, Rye, and Oats

In wheat, barley, rye, and oats, Tilt controls rusts (*Puccinia* spp.), powdery mildew (*Erysiphe* spp.), leaf blight and glume blotch (*Septoria* spp.), tan spot (*Pyrenophora tritici-repentis*), Helminthosporium leaf blight, spot blotch, barley scald (*Rhynchosporium secalis*), and net blotch (*Pyrenophora teres*). Apply 4 fl. oz. of Tilt per acre by ground or aerial application or through irrigation equipment. Highest yields are normally obtained when Tilt is applied to the emerging flag leaf. Tilt can be applied until the ligule of the flag leaf emerges. (Feekes growth stage 8). Do not apply after this growth stage to avoid possible illegal residues. Tilt may be applied earlier if disease symptoms appear (especially applicable to barley).

### Wheat Only

For control of foot rot (*Pseudocercospora* spp.), apply 4 fl. oz. of Tilt per acre plus half rates of other EPA-registered fungicides such as Benlate®, Mertect®, and Topsin® M. Apply at tillering, but before elongation has occurred. Do not apply more than 4 fl. oz. of Tilt per season.

**Notes:** To avoid possible illegal residues: (1) Do not apply more than 4 fl. oz. of Tilt per acre per season. (2) Do not apply to oats within 40 days of harvest. (3) Do not graze or feed livestock treated wheat, barley, or rye forage or cut the green crop for hay or silage. After harvest, the straw from these crops may be used for bedding or feed. (4) Tilt-treated oat forage may be grazed, and oat forage and hay may be fed to livestock.

**Rice**

In rice, Tilt controls sheath blight (*Rhizoctonia solani*), brown leaf spot (*Helminthosporium oryzae*), narrow brown leaf spot and brown blotch (*Cercospora oryzae*), leaf smut (*Entyloma oryzae*), sheath spot (*Rhizoctonia oryzae*), kernel smut (*Tilletia barclayana*), and black sheath rot (*Gaeumannomyces graminis*). Tilt also suppresses stem rot (*Sclerotium oryzae*) and false smut (*Ustilagoideae virens*). Apply Tilt on either of the following schedules as an aerial spray in 5-10 gals. of water per acre:

- A. 6 fl. oz./A at first internode elongation (up to 2-inch panicle) and repeat at swollen boot. Make the second application 10-14 days after the first application, but before the boot splits and head emerges. Tilt provides best control of sheath blight when the first application is applied at disease appearance in the field. The first application is recommended when 5% or fewer of the tillers are infected.
- B. 10 fl. oz./A at first internode elongation (up to 2-inch panicle). The 10 oz. rate is recommended if greater than 10% of the tillers are infected with sheath blight. If disease reappears, use another registered fungicide for the second application.

**Important:** To avoid possible illegal residues: (1) Do not apply to stubble or ratoon crop rice. (2) Do not use in rice fields where commercial farming of crayfish will be practiced. (3) Do not drain water from treated rice fields into ponds used for commercial catfish farming. (4) Do not use water drained from treated fields to irrigate other crops. (5) Do not use in CA.

**Endangered Species Restrictions:** The use of Tilt (propiconazole) on rice is restricted to protect the endangered fat pocketbook pearly mussel (*Potamilus capax*) and its habitat. Use is prohibited in the following areas of AR:

**Mississippi County** – Within the basin that drains directly into the Right Hand Chute of Little River, south of Big Lake National Wildlife Refuge.

**Poinsett County** – Between Crowley's Ridge and the levee east of the Right Hand Chute of Little River and the St. Francis Floodway. Use is also prohibited west of Rt. 140 and north of Rt. 63 at the siphon near Marked Tree. Except that the prohibited area does not include the area bounded by Arkansas highway 373 on the west, highway 63 on the east, and highway 14 on the south.

**Cross, St. Francis, and Lee Counties** – Between Crowley's Ridge and the levee east of the Right Hand Chute of Little River, and the St. Francis Floodway, as far south as the confluence of L'Anguille River (Lee County).

**Wild Rice (MN only)**

For control of *Helminthosporium* leaf blight and brown spot (*Bipolaris* spp.), apply 6 fl. oz. of Tilt per acre at both booting and heading, or make a single application of 8 fl. oz./A booting. For aerial applications, apply in a spray volume of 5-10 gals./A.

**Important:** Do not use water drained from treated fields to irrigate other crops.

**CORN (FIELD CORN, FIELD CORN GROWN FOR SEED, SWEET CORN, AND POPCORN)**

For control of *Helminthosporium* leaf blights (*Helminthosporium maydis*, *H. turcicum*, and *H. carbonum*), rusts (*Puccinia* spp.), gray leaf spot (*Cercospora zeae-maydis*), and eye spot (*Kabatiella zeae*), apply Tilt by ground or aerial application or through irrigation equipment according to the following schedule:

**Helminthosporium Leaf Blights:** Apply 2-4 fl. oz. of Tilt per acre when disease first appears and continue on a 7 to 14-day schedule.

**Rusts:** Apply 4 fl. oz. of Tilt per acre when rust pustules first appear and continue on a 7 to 14-day schedule.

**Gray Leaf Spot and Eye Spot:** Apply 4 fl. oz. of Tilt per acre when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule.

**Important:** To avoid possible illegal residues: (1) Do not apply Tilt to field corn and field corn grown for seed after silking. (2) Do not apply more than 16 fl. oz. of Tilt per acre per season. (3) Do not apply to sweet corn within 14 days of harvest. (4) Do not harvest field corn, field corn grown for seed, or popcorn for forage within 30 days of application. (5) Do not harvest sweet corn for forage within 14 days of application.

**GRASSES GROWN FOR SEED (NEBRASKA, OREGON, WASHINGTON, IDAHO, AND MINNESOTA ONLY)**

Tilt controls rusts (*Puccinia* spp.), powdery mildew (*Erysiphe* spp.), and *Selenophoma* stem eyespot (*Selenophoma*) in perennial ryegrass, fescues, bluegrass, orchardgrass, and/or wheatgrasses.

Mix 4-8 fl. oz. of Tilt (maximum 4 fl. oz. on bluegrass) in a minimum of 20 gals. of water per acre for ground application, or in a minimum of 10 gals. of water per acre for aerial application. Tilt may also be applied through irrigation equipment. Apply when powdery mildew and *Selenophoma* infections or rust pustules are noticeable and increasing in number, in late spring or early summer. Repeat at 14 to 21-day intervals. To maximize control under severe rust pressure, use the higher rate of 8 fl. oz./A (except on bluegrass), and make applications at 14-day intervals until the seed is mature. Do not apply more than 32 fl. oz. of Tilt per acre per growing cycle. Make the last application at least 20 days before seed matures. For bluegrass, it is important to begin application early in the growing season.

**Important:** To avoid possible illegal residues, do not: (1) feed hay cut within 20 days of the last application; or (2) graze treated areas within 140 days of the last application.

## PEANUTS

### Peanuts – Tilt Alone For Follar Disease Control

**Late Leaf Spot (*Cercosporidium*)** – Apply 4 fl. oz. of Tilt per acre beginning applications 35-40 days after planting or at the first appearance of disease. Continue applications on a 10 to 14-day schedule. Tilt also may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

**Early Leaf Spot (*Cercospora*)** – Apply 2.5-4 fl. oz. of Tilt per acre beginning applications 35-40 days after planting or at the first appearance of disease. Continue applications on a 14-day schedule. Under heavy disease pressure use higher recommended application rates. Tilt also may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

**Important:** To avoid possible illegal residues: (1) Do not apply more than 16 fl. oz. of Tilt per acre per season. (2) Do not feed green vines to livestock or graze livestock in treated area. (3) Do not apply within 14 days of harvest.

### Peanuts – Tilt Plus Chlorothalonil For Follar Disease Control

**Early and Late Leaf Spot** – Combinations of Tilt with products containing chlorothalonil may be used for early and late leaf spot control. Apply 2 fl. oz. of Tilt as a tank mixture with 0.75 lb. active ingredient chlorothalonil per acre. Use the following table as a guide to determine the appropriate amount of chlorothalonil to use in the tank mixture. Begin applications 35-40 days after planting or at the first appearance of disease and continue applications on a 10 to 14-day schedule. Tilt plus chlorothalonil also may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

**Important:** Use the following table as a guide to determine the amount of product needed for the recommended tank mix ratios of Tilt with various chlorothalonil formulations. To avoid possible illegal residues: (1) Do not apply more than 16 fl. oz. of Tilt per acre per season. (2) Do not apply within 14 days of harvest. (3) Do not graze livestock in treated area. (4) Do not feed hay or threshings from treated fields to livestock.

Number of acres to be treated	Amount of Tilt required for treated area (total acres to be treated)	Amount of chlorothalonil product required for treated area by formulation type (total acres to be treated)	
		6 lbs. a.i. per Gal.	4.17 lbs. a.i. per Gal.
1	2 fl. oz. (1/8 pt.)	1 pt.	1.5 pts.
5	10 fl. oz. (5/8 pt.)	5 pts.	7.5 pts.
10	1.25 pts.	10 pts.	15 pts.
50	6.25 pts.	50 pts.	75 pts.
100	12.5 pts.	100 pts.	150 pts.

### Peanuts – Tilt Alone for Southern Stem Rot Control

For the control of Southern Stem Rot (*Sclerotium rolfsii*), apply Tilt using one of the following schedules:

A. Apply 4 fl. oz. of Tilt per acre to the crown and pegging zones of the plant using chemigation or directed ground application. Begin applications 45 days after planting, or at the first appearance of disease, and repeat on a 14-day schedule.

**Important:** To avoid possible illegal residues: (1) Do not apply more than 16 fl. oz. of Tilt per acre per season. (2) Do not feed green vines to livestock or graze livestock in treated area. (3) Do not apply within 14 days of harvest.

B. Apply 8 fl. oz. of Tilt per acre to the crown and pegging zones of the plant using chemigation or directed ground application. Make 2 applications: the first at pegging (approximately 60 days after planting) or at the first appearance of disease, and the second application 3-4 weeks later.

**Irrigation:** When applying Tilt in irrigation water for Southern Stem Rot Control of Peanuts, use a minimum of 0.25-0.5 inch of irrigation water per acre. Use enough water so that the fungicide penetrates the peanut canopy and reaches the crown of the plant where *Sclerotium rolfsii* is most active.

**Ground Application:** When applying Tilt by ground equipment, use a minimum of 20-60 GPA directed to the crown of the plant where *Sclerotium rolfsii* is most active. For best disease control, the higher carrier volume is recommended. Canopy openers also may be used to improve fungicide placement.

**When applying Tilt via irrigation or as a directed ground application, additional methods should be employed for leaf spot control.**

**Banded Application:** For banded applications, the treated area is the area covered by the band, not total cropland planted. The following formula can be used to calculate the amount of Tilt needed per acre of crop when banded applications are made.

$$\frac{\text{band width in inches}}{\text{row spacing in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

**Important:** To avoid possible illegal residues: (1) Do not apply more than 16 fl. oz. of Tilt per acre per season. (2) Do not graze livestock in treated area. (3) Do not feed hay or threshings from treated fields to livestock. (4) Do not apply within 21 days of harvest.

**PINEAPPLE (HAWAII ONLY)**

For control of butt rot disease of pineapple (*Ceratocystis paradoxa*), apply 0.75 fl. oz. (22 ml) of Tilt per 100 gals. of water (1:17,000). Treatments can be made in either a cold or hot water dip.

**Cold Water Dip** – Immerse crowns to give thorough wetting, remove, and allow to drain.

**Hot Water Dip** – Maintain water temperature at 125°F (52°C). Soak crowns for 20-30 minutes, remove, and allow to drain.

**Important:** To avoid possible illegal residues: (1) Do not use treated crowns for food or feed. (2) Do not graze while plant is growing. (3) Do not graze tops until fruit is harvested. (4) Dispose of used dip solution according to local, state, and federal regulations.

**SUGARCANE (HAWAII ONLY)**

For control of pineapple disease of sugarcane (*Ceratocystis paradoxa*), apply Tilt to cut seed pieces using 0.75 fl. oz. (22 ml) per 100 gals. of water (1:17,000). Treatments can be applied in either a cold or hot water dip. Do not use treated seed pieces for food or feed purposes. Dispose of spent dip solution according to state and federal regulations.

**Cold Water Dip** – Immerse seed pieces to give thorough wetting, remove, and allow to drain.

**Hot Water Dip** – Maintain water temperature at 125°F (52°C). Soak the seed pieces for 20-30 minutes, remove, and allow to drain.

**STORAGE AND DISPOSAL****Pesticide Disposal**

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

**Container Disposal**

Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. Stay out of smoke from burning containers.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

**PRECAUTIONARY STATEMENTS****Hazards to Humans and Domestic Animals****WARNING/AVISO**

This product may cause skin sensitization reactions in certain individuals.

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing spray mist.

**Statement of Practical Treatment**

**If in eyes:** Flush with plenty of water. Get medical attention if irritation persists.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**If swallowed:** Do not induce vomiting. Promptly drink a large quantity of milk, egg whites, or gelatin solution. If these are not available, drink large quantities of water. Never give anything by mouth to an unconscious person. Call a physician or Poison Control Center immediately.

**If inhaled:** Move victim to fresh air.

**Note to Physician:** There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give a saline laxative and supportive therapy.

**Personal Protective Equipment**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category G on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

##### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### Environmental Hazards

This pesticide is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Refer to product labeling for use restrictions to protect endangered species.

#### Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

Tilt® and Mertect® trademarks of Syngenta  
U.S. Patent No. 4,079,062

Benlate® trademark of E. I. duPont de Nemours and Company, Inc.


Topsin® trademark of Pennwalt Corporation

©2001 Syngenta

Product of Switzerland  
Formulated in the USA

Syngenta Crop Protection, Inc.  
Greensboro, North Carolina 27409  
[www.syngenta-us.com](http://www.syngenta-us.com)

**SCP 617A-M2D 0201**



# TILT®

## FUNGICIDE

For control of certain diseases in celery, cereals, corn, grasses grown for seed, peanuts, pineapple, and sugarcane

Active Ingredient:	
Propiconazole	
(CAS No. 60207-90-1) . . . . .	41.8%
Other Ingredients:	58.2%
Total:	100.0%

EPA Reg. No. 100-617  
EPA Est. 34704-MS-2®  
EPA Est. 67545-AZ-1®  
(Superscript is first letter of lot number on jug)

See directions for use in attached booklet.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Tilt® trademark of Syngenta  
U.S. Patent No. 4,079,062

©2001 Syngenta

Syngenta Crop Protection, Inc.  
Greensboro, North Carolina 27409  
www.syngenta-us.com

SCP 617A-M2D 0201

ONE GALLON  
U.S. Standard Measure

## KEEP OUT OF REACH OF CHILDREN. WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

This product may cause skin sensitization reactions in certain individuals.

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing spray mist.

#### Statement of Practical Treatment

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If swallowed: Do not induce vomiting. Promptly drink a large quantity of milk, egg whites, or gelatin solution. If these are not available, drink large quantities of water. Never give anything by mouth to an unconscious person. Call a physician or Poison Control Center immediately.

If inhaled: Move victim to fresh air.

Note to Physician: There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give a saline laxative and supportive therapy.

#### Environmental Hazards

This pesticide is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Refer to product labeling for use restrictions to protect endangered species.

#### Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

#### Chemigation

Refer to supplemental labeling in attached booklet for use directions on chemigation. Do not apply this product through any type of irrigation system unless the supplemental labeling on chemigation is followed.

syngenta

**NEXT**

**LABEL**

100-617

3-30-2001

1/6

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number NOTIFICATION
	<b>Application for Pesticide - Section I</b>		

1. Company/Product Number 100-617	2. EPA Product Manager Mary Waller	3. Proposed Classification  <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Tilt@Plus Fungicide	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419  <input type="checkbox"/> Check if this is a new address	6. <b>Expedited Review.</b> In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____  Product Name _____	

**Section - II**


<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b>  MAR 30 2001
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

**Explanation:** Use additional page(s) if necessary. (For Section I and Section II.)  
 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. The following changes are being made via this notification: 1) Company name and address have been updated to reflect Syngenta Crop Protection, Inc. 2) The Conditions of Sale and Warranty statement has been changed to reflect the name change. Because Syngenta has been formed by the merger of Novartis Crop Protection, Inc. and Zeneca Ag Products, we have chosen to use the former Zeneca warranty statement as the Syngenta warranty statement. No other changes occur in the statement other than the name change. 3) The copyright date reflects Syngenta. 4) Trademark statements have been updated to reflect Syngenta for those products for which Syngenta holds the trademark. 5) The internet address has been changed to reflect Syngenta. 6) Other places in the label which referring to the company name have been updated.

**Section - III**

<b>1. Material This Product Will Be Packaged In:</b>				2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No			
*Certification must be submitted		If "Yes" Unit Packaging wgt.    No. per Container	If "Yes" Unit Packaging wgt.    No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Nan S. Padgett	Title Label Group Leader	Telephone No. (Include Area Code) 336-632-7567
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Label Group Leader	
4. Typed Name Nan S. Padgett	5. Date March 19, 2001	

MAR 30 2001

**TILT<sup>®</sup> PLUS****FUNGICIDE**

For control of foot rot and early season control of foliar diseases in fall-seeded wheat in Idaho, Oregon, and Washington only

SHAKE WELL BEFORE USING.

See additional precautionary statements and directions for use inside booklet.

Using label instructions, apply the entire contents of this container to the target crop once the container is opened.

EPA Est. 34704-MS-2Ⓢ, EPA Est. 5905-GA-1Ⓢ  
(Superscript is first letter of lot number on container)

SCP 617D-M18D 0201

Total 140 fl. oz.  
U.S. Standard Measure

**Tilt**

PULL HERE TO OPEN ►

Active Ingredient:	
Propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl] methyl]-1H-1,2,4-triazole . . . . .	41.8%
Inert Ingredients:	58.2%
Total:	100.0%

EPA Reg. No. 100-617

**KEEP OUT OF REACH OF CHILDREN.**

**WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

40 fl. oz. U.S. Standard Measure

**Topsin<sup>®</sup> M 4.5F**

Active Ingredient:	
Thiophanate-methyl(dimethyl[[(1,2-phenylene)-bis(iminocarbonothioyl)] bis(carbamate)]*) . . . . .	46.2%
Inert Ingredients:	53.8%
Total:	100.0%

\*Also known as dimethyl 4,4'-O-phenylene bis [3-thiallophanate]

EPA Reg. No. 4581-352

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

100 fl. oz. U.S. Standard Measure

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

#### Tilt Plus

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

### GENERAL INFORMATION

Tilt Plus is a combination of fungicides for the control of foot rot (*Pseudocercospora* sp.) and early season foliar diseases on fall-seeded wheat in ID, OR, and WA only. Using label instructions, apply the entire contents of this container to the target crop once the container is opened.

**Important:** Do not apply to other cereals, such as barley, oats, and rye. Do not use in nurseries, on turf, landscape plantings, or as a tree injection.

**FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.**

#### Spray Equipment

Thorough coverage is necessary to provide good disease control.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate sprayer before use.

Use a pump with capacity to: (1) maintain 35-40 psi at nozzles, and (2) provide sufficient, *continuous* agitation in tank to keep mixture in suspension – this requires recirculation of 10% of tank volume per minute. Use a jet agitator or liquid sparge tube for agitation. Do not air sparge.

It is suggested that screens be used to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.

**Aerial Application:** Apply in a minimum of 5 gals. of water per acre. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals.

**Ground Application:** Apply Tilt Plus by ground equipment in a minimum of 15 gals. of water per acre.

**Chemigation:** Do not apply this product through any type of irrigation system.

#### Mixing Instructions

Prepare no more spray mixture than is required for the immediate operation. When the container is opened, immediately pour the entire contents of this container into the spray tank. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

**Important:** Custom applicators must use a closed system for mixing and loading this product.

**Tilt Plus Alone:** Add  $\frac{1}{2}$ - $\frac{2}{3}$  of the required amount of water to the spray or mixing tank. Set the agitator to producing a rippling or rolling movement on the surface of the mix water. With the agitator running, add all of the contents of both chambers of the Tilt Plus container to the tank immediately. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the Tilt Plus has completely dispersed into the mix water. Maintain agitation until all of the mixture has been sprayed.

**Tilt Plus + Tank Mixtures:** Consult your dealer or Syngenta sales representative for information concerning the compatibility with other tank mix partners. **Note: When using Tilt Plus in mixture, any product in water-soluble packaging should be added to the tank prior to adding Tilt Plus. Allow water-soluble packaging to completely dissolve and the mixture product to completely disperse before adding Tilt Plus to the tank.**

Add  $\frac{1}{2}$ - $\frac{2}{3}$  of the required amount of water to the spray tank. With the agitator running, add the tank mix partners to the spray tank. In general, tank mix partners should be added in this order: wettable powders, dry flowable formulations, liquid flowable formulations, and emulsifiable concentrations. Always allow each tank mix partner to become fully dispersed before adding the next product. Continue agitation while adding the remainder of the water and the Tilt Plus to the spray tank. Add all of the contents of both chambers of the Tilt Plus container to the tank at once. Allow the Tilt Plus to become completely dispersed into the mix water. Maintain agitation throughout the spraying operation.

If using Tilt Plus in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix partner label. No label dosage rate should be exceeded and the most restrictive label precautions and limitations should be followed. This product should not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the products are registered.

**Important:** Do not tank mix with highly alkaline pesticides.

### Rotational Crops

Soybeans may be planted as a double crop following a wheat crop which has been treated with Tilt Plus. Do not use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding.

To avoid possible illegal residues, do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of Tilt Plus application to the preceding crop, unless the second crop appears on the following list: Barley, Celery, Grasses Grown for Seed, Rice, Rye, Wheat.

## FALL-SEEDED WHEAT IN IDAHO, OREGON, AND WASHINGTON ONLY

For the control of foot rot (*Pseudocercospora* sp.) and the early season control of rusts (*Puccinia* sp.), powdery mildew (*Erysiphe graminis*), leaf and glume blotch (*Septoria* sp.), tan spot (*Pyrenophora tritici-repentis*), and Helminthosporium leaf blight, use the entire contents of this container to treat 10 acres of fall-seeded wheat. Apply Tilt Plus at tillering, but before stem elongation has occurred.

**Notes:** (1) Do not make more than one application of Tilt Plus per growing season. After applying Tilt Plus to fall-seeded wheat, do not apply Tilt, Tilt Gel, Topsin 4.5M, Topsin M 70W, or any other product containing thiophanate-methyl during the remainder of the growing season. (2) Do not graze or feed livestock-treated forage or cut the green crop for hay or silage. After harvest, the straw may be used for bedding or feed. (3) Using label instructions, apply the entire contents of this container to the target crop once the container is opened.

## STORAGE AND DISPOSAL

**Storage Instructions:** Store in the original container at temperatures above 32°F. Using label instructions, apply the entire contents of this container to the target crop once the container is opened.

### Pesticide Disposal

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

### Container Disposal

Do not reuse empty container. Triple rinse container (or equivalent) and then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. Stay out of smoke from burning containers.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

## PRECAUTIONARY STATEMENTS

### Tilt

#### Hazards to Humans and Domestic Animals

#### WARNING/AVISO

This product may cause skin sensitization reactions in certain individuals.

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing spray mist.

#### Statement of Practical Treatment

**If in eyes:** Flush with plenty of water. Get medical attention if irritation persists.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**If swallowed:** Do not induce vomiting. Promptly drink a large quantity of milk, egg whites, or gelatin solution. If these are not available, drink large quantities of water. Never give anything by mouth to an unconscious person. Call a physician or Poison Control Center immediately.

**If inhaled:** Move victim to fresh air.

**Note to Physician:** There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give a saline laxative and supportive therapy.

**Topsin M 4.5F****Hazards to Humans and Domestic Animals****CAUTION**

**Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation.** Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

**Statement of Practical Treatment**

**If in eyes:** Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

**If on skin:** Immediately flush skin with plenty of water. Get medical attention if irritation persists.

**If swallowed:** Induce vomiting immediately by giving 2 glasses of water and sticking finger down throat. Call a physician. Never give anything by mouth to an unconscious person.

**Tilt Plus****Personal Protective Equipment**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category G on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Control Statements**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations****Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

**Physical or Chemical Hazards**

Do not use, pour, spill, or store near heat or open flame.

Tilt® and Twin-Pak™ trademarks of Syngenta

U.S. Patent No. 4,079,062

Topsin® trademark of Nippon Soda Company, Ltd. and is licensed to Elf Atochem North America

©2001 Syngenta

Syngenta Crop Protection, Inc.

Greensboro, North Carolina 27409

www.syngenta-us.com

SCP 617D-M18D 0201

**TILT<sup>®</sup> PLUS****Twin-Pak™****FUNGICIDE**

For control of foot rot and early season control of foliar diseases in fall-seeded wheat in Idaho, Oregon, and Washington only

EPA Est. 34704-MS-2Ⓢ

EPA Est. 5905-GA-1Ⓢ

(Superscript is first letter of lot number on container)

©2001 Syngenta

Syngenta Crop Protection, Inc.  
Greensboro, NC 27409  
www.syngenta-us.com

**SCP 617D-M18D 0201**

Total **140 fl. oz.**  
U.S. Standard Measure

See directions for use in attached booklet.

SHAKE WELL BEFORE USING.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

**Tilt**

Active Ingredient:

Propiconazole: 1-[[2-(2,4-dichloro-phenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1*H*-1,2,4-triazole. . . . . 41.8%

Inert Ingredients: 58.2%

Total: 100.0%

EPA Reg. No. 100-617

Tilt® and Twin-Pak™ trademarks of Syngenta

U.S. Patent No. 4,079,062

**40 fl. oz. U.S. Standard Measure****Topsin® M4.5F**

Active Ingredient:

Thiophanate-methyl (dimethyl [(1,2-phenylene)-bis(iminocarbonothioyl)] bis [carbamate])\* . . . . . 46.2%

Inert Ingredients: 53.8%

Total: 100.0%

\*Also known as dimethyl 4,4'-*O*-phenylene bis [3-thioallophanate]

EPA Reg. No. 4581-352

Topsin® trademark of  
Nippon Soda Company, Ltd. and is licensed to  
Elf Atochem North America

**100 fl. oz. U.S. Standard Measure**