

Caparol

4L

Herbicide

For selective weed control in cotton and celery

Active Ingredient
Prometryn 2,4-bis-
isopropylamino-
6-methylthio-
s-triazine 44.4

Inert
Ingredients 55.6%
Total 100.0%

2 1/2
Gallons
U.S. Standard
Measure

Caparol 4L contains 4.4 lbs.
active ingredient per gallon.
Mix well before using.

Keep Out of Reach
of Children

Warning

See additional precautionary
statements at end of
label booklet

See directions for use
inside booklet

Caparol trademark of
CIBA-GEIGY for prometryn

EPA Reg. No. 100-620
EPA Est. 100-AL-1

1981 CIBA-GEIGY
Corporation

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, NC 27419
CGA 17L6

CIBA-GEIGY

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the Directions for Use and Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product are hereby incorporated by reference into this sale. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product. CIBA-GEIGY makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller disavow any liability for the use or handling of this product in violation of the foregoing Conditions of Sale and Warranty. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

CIBA-GEIGY warrants that this product conforms to the description on the label and to the specifications of the manufacturer.

CIBA-GEIGY makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller disavow any liability for the use or handling of this product in violation of the foregoing Conditions of Sale and Warranty. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

General Information

This product is for use in the control of weeds in cotton and celery.

Caparol 4L is a selective herbicide for use in cotton and celery. It controls a wide range of weeds, including annual and perennial species. The active ingredient, prometryn, is a triazine herbicide that acts by inhibiting photosynthesis in the weeds. Caparol 4L is formulated as a liquid concentrate, which allows for easy mixing and application. It is important to read the label and follow the directions for use carefully to ensure effective control of weeds while minimizing damage to the crop. The product is registered with the EPA and is labeled for use in cotton and celery. It is not intended for use in other crops or for other purposes. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

FAILURE TO FOLLOW PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PERFORMANCE AND OR ILLEGAL RESIDUES

Equipment and Mixing

Caparol 4L should be mixed with water in a clean container. The mixture should be applied immediately after mixing. Do not store the mixture for more than 24 hours.

Caparol 4L should be mixed with water in a clean container. The mixture should be applied immediately after mixing. Do not store the mixture for more than 24 hours. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

Nozzles

Use a nozzle that provides a fine spray. Do not use a nozzle that produces a coarse spray or a stream. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

Mixing

Caparol 4L should be mixed with water in a clean container. The mixture should be applied immediately after mixing. Do not store the mixture for more than 24 hours. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

Cleaning

Caparol 4L should be cleaned with water. Do not use solvents. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

Seedbed Preparation

Caparol 4L should be applied to the seedbed. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

Band Treatment

Caparol 4L should be applied in a band. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

Caparol 4L should be applied in a band. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

Cotton

Caparol 4L should be applied to cotton. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

Caparol 4L should be applied to cotton. The user must read and follow the Directions for Use and Conditions of Sale and Warranty before using this product.

Note: Do not use Caparol 4L on livestock or in areas where livestock are present.



Recap: $\mathcal{L} = \{L_1, \dots, L_n\}$

2. (1) 1988年11月1日

P. 121

8. 10. 1990. 41

235

Rotational Crops

510 4E

Tretlan EC Tank Mixture

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

Table 6 Tank Mixture with Trellan EC

Soil texture	Broadcast rate per acre	
	Trellan EC	Caparol 4L
2. Heavy sand	1.0	0.5
3. Medium sand	1.0	0.4 (0.2) *
4. Medium silt	1.0	0.35
5. Fine silt	0.5	0.25
6. Heavy silt	0.5	0.2

less than 30% reduction, in Arizona and California.

Precautions: Do not use old seeds of new strains. Use seeds of fresh process. Seed in barrels where flooding is controlled. Key: 1. Put seed in a barrel that is 1/2 full of water when process is 1/2 full of water. 2. When water is added, add water to the barrel after each seed is added. 3. When seed is added, add water to the barrel after each seed is added.

Note: Do not feed treated hay to livestock or graze treated areas.

Rotational Crops: Cattle and sheep graze on the fields and are moved in the fall after a spring application of fertilizer. Cattle and sheep bare the field when they are moved in the fall. They are then moved and are used in feed. Refer to the farm for the current status of the crops.

Trelian EC Split Application - Arizona and California

Any direct and incorporated application of Treated E.C. is considered a direct application, and the appropriate label must be used. If the Treated E.C. is used in any other way, the label must be changed to reflect the use. For example, if Treated E.C. is incorporated into a product, the label must reflect the use of the product.

DSMA, SSMA

the 1990s, the number of people in the world who are undernourished has declined from 1.1 billion to 800 million. The number of people who are malnourished has declined from 1.5 billion to 1 billion. The number of people who are obese has increased from 100 million to 300 million. The number of people who are overweight has increased from 100 million to 300 million. The number of people who are obese and overweight has increased from 100 million to 300 million. The number of people who are obese and overweight has increased from 100 million to 300 million.

— 252 —

• **References**

the 1990s, the number of people in the world who are under 15 years of age is expected to increase from 1.1 billion to 1.5 billion. The number of people aged 65 and over is expected to increase from 200 million to 400 million. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion. The number of people aged 15 and over is expected to increase from 3.5 billion to 4.5 billion.

Direct seeded Celery

[illegible]

the 1990s, the number of people in the United States who are 65 years of age or older is projected to increase from 20 million to 30 million, and the number of people 75 years of age or older is projected to increase from 10 million to 15 million (U.S. Census Bureau, 1996). The number of people 85 years of age or older is projected to increase from 2 million to 4 million (U.S. Census Bureau, 1996). The number of people 90 years of age or older is projected to increase from 500,000 to 1 million (U.S. Census Bureau, 1996). The number of people 95 years of age or older is projected to increase from 100,000 to 200,000 (U.S. Census Bureau, 1996). The number of people 100 years of age or older is projected to increase from 10,000 to 20,000 (U.S. Census Bureau, 1996).

As the number of nodes in the network increases, the number of nodes that are not connected to any other nodes increases. This is because the number of nodes that are not connected to any other nodes is proportional to the number of nodes in the network. As the number of nodes in the network increases, the number of nodes that are not connected to any other nodes increases.

[illegible]

Transplants

[illegible]

Table 7 Transplanted Celery

State	Soil	Broadcast rate per acre
Alabama	1 lb. per acre	1.0-2.0 pts.
California	1 lb. per test plot	1.0-2.0 pts.
	1 lb. per test plot	1.0-4.0 pts.
	1 lb. per test plot	3.2-4.0 pts.
Florida	1 lb. per test plot	1.0-6.0 pts.
Georgia	1 lb. per test plot	1.0-6.0 pts.
Illinois	1 lb. per test plot, 1 lb. per acre	2.4 pts.
Mississippi	1 lb. per test plot	1.0-4.0 pt.

Rotational Crops

The following information was obtained from the above mentioned
the file of the "El Centro de la Raza" (E.C.R.), located at 1000
N. Main Street, El Centro, California.

Storage and Disposal

[illegible]

Precautionary Statements

WARNING

First Aid

Environmental Hazards

Caparol

4L

Herbicide

For selective weed control in cotton and celery

Active Ingredient
Prometryn
2,4-bis(isopropylamino)-
6-(methylthio)-
triazine 44.4%

Inert ingredients 55.6%
Total 100.0%

Caparol 4L contains 4 lbs active ingredient per gal.

Shake well before using.

See directions for use in attached booklet.

EPA Reg. No. 100-620

EPA Est. 100-AL-1

2½
Gallons
U.S. Standard Measure

Keep Out of Reach
of Children.

Warning

Precautionary Statements

Hazard to Humans and Domestic Animals

First Aid

Environmental Hazards

Caparol, trademark of CIBA-GEIGY for prometryn.

© 1981 CIBA-GEIGY Corporation

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419
CGA 17L6

CIBA-GEIGY