

APR 15 1992

Ms. Karen S. Stumpf  
Agricultural Division  
Ciba-Geigy Corporation  
P.O. Box 18300  
Greensboro, NC 27419-8300

545110 | 300  
|  
18

Dear Ms. Stumpf:

Subject: Amended Labeling  
Ridomil MZ58 Fungicide  
EPA Registration No. 100-629  
Your Submissions Dated February 26, 1992 and  
March 3, 1992

The amendments referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, are acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. On page 5, in the last paragraph, revise the third sentence to "For aerial application, apply in 3-10 gallons of water/acre". The phrase "are generally needed" can imply that the spray volume may be more or less than that indicated. A minimum of 2 gallons per acre is permitted by the PD4.

b. On page 6, under Ridomil MZ58 + Tank Mixtures, include only those EBDC fungicides which contain mancozeb since applications may not be made to areas where other EBDC fungicides are used. Delete "Manzate" which contains maneb. Dithane M45 and Manzate 200 both contain mancozeb and may be retained. If recommendations are retained, specific directions for use on each crop must include the maximum rate of mancozeb per application.

c. Under potatoes and tomatoes, delete tank mix recommendations for Manzate 80W and Dithane M-22 since both contain maneb.

d. For tomatoes, the tank mix directions are confusing. The statement "Do not apply more than 2 lbs./acre of (products) in combination with Ridomil MZ58" can be taken to mean 2 lbs. of (product) + 1.5 lbs. Ridomil MZ58 or to mean 2 lbs. of the

H7505C:J.Fairfax:100-629:4, 14/92

combination. This sentence must be rewritten to more clearly define the rate. If 2 lbs. of the 80% product is used, the maximum rate of mancozeb per application would be exceeded for areas west of the Mississippi River. The directions should be revised to state "When making applications of Manzate 200 or Dithane M45, use 1.0 lb. of product and 1.5 lbs. Ridomil MZ58 per acre every 14 days for control of late blight in areas west of the Mississippi River or 2.0 lbs. of product and 1.5 lbs. Ridomil MZ58 on the same schedule for areas east of the Mississippi River." Also, in footnote (5), delete nabam and zineb since these products are no longer marketed for crop uses.


e. For specific crops add the following statement following the sentence stating maximum mancozeb limits per growing season: "If tank mixed with other products containing mancozeb, do not exceed \_\_\_\_\_ lbs. mancozeb active ingredient per acre per application." The maximum mancozeb rate/application for cucumbers, melons, squash, potatoes and sugar beets is 1.6 lbs.; for onions, 2.4 lbs. and for tomatoes, 1.6 lbs. west of the Mississippi, 2.4 lbs. east of the Mississippi.

f. On page 9, qualify the use on melons by listing the types of melons approved for mancozeb. Following melons, add "(cantaloupe, casaba, crenshaw, honeydew and watermelons)."

Submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

  
Susan T. Lewis  
Product Manager (21)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosure

4121 100-629

30418

Water Soluble Bag

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

APR 15 1992

Ridomil® MZ58

Fungicide

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No.

For control of certain diseases of cucumbers, melons,  
summer squash, onions, potatoes, sugar beets, and tomatoes

100-629

Five Lbs.                      Ten Lbs.  
Net Weight                      Net Weight

Active Ingredients:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester .....	10.0%
A coordination product of zinc ion and manganese ethylene bisdithiocarbamate* .....	48.0%
in which the ingredients are:	
Manganese <sup>++</sup> .....	9.6%
Zinc <sup>++</sup> .....	1.2%
Ethylene bisdithiocarbamate ion (C <sub>4</sub> H <sub>6</sub> N <sub>2</sub> S <sub>4</sub> ) .....	37.2%
<u>Inert Ingredients:</u>	<u>42.0%</u>
Total:	100.0%

\*Same as the active ingredient found in Dithane® M-45 and Manzate® 200.

Ridomil MZ58 is a wettable powder

KEEP OUT OF REACH OF CHILDREN.

WARNING

AVISO

PRECAUCIÓN AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

See additional precautionary statements on back of container.

See directions for use on panel at right.

EPA Reg. No. 100-629

EPA Est. No. 32761-MO-3

CIBA-GEIGY

**BEST AVAILABLE COPY**

This outer protective container contains Ridomil MZ58 in an inner water-soluble bag. Entire inner bag and contents dissolve in water. After opening outer container, immediately dump entire unopened inner bag into the partially filled sprayer or mix tank. Do not handle the soluble bag or expose it to moisture, since this may cause rupturing.

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

BEST AVAILABLE COPY

50418

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons, except those knowingly involved in the application. The area being treated must be vacated by unprotected persons.

Re-entry and Worker Protection Statements

Worker Safety Rules

Keep all unprotected persons, children, livestock and pets away from treated area or where there is a danger of drift. Do not rub eyes or mouth with hands. See Statements of Practical Treatment.

Personal Protective Equipment - For Mixers, Loaders, Applicators and Early Re-entry Workers.

HANDLE THIS PRODUCT ONLY WHEN WEARING THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT: All agricultural workers/handlers (i.e., mixers, loaders, and applicators) applying EBDC's must wear coveralls over long-sleeved shirt and long pants; shoes, socks, and chemical-resistant gloves. During mixing and loading, a chemical-resistant apron and goggles or a face shield must also be worn. For agricultural workers, where completely enclosed cabs with positive pressure filtration or an enclosed cockpit for aerial application are used, a long-sleeved shirt and long pants may be worn in place of the protective clothing described above. Chemical-resistant gloves must be available in the cab or cockpit and worn upon exiting. The gloves must be kept in an enclosed container in the cab or cockpit to prevent contamination of the inside of the cab or cockpit. During aerial application, human flaggers are prohibited unless in totally enclosed vehicles.

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before eating, smoking or drinking. Always wash hands and arms with soap and water before using the toilet.

After work take off all clothes and shoes. Shower using soap and water. Wear only clean clothes. Do not use contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Personal clothing worn

during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with Ridomil MZ58 must be destroyed according to state and local regulations.

HEAVILY CONTAMINATED OR DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED. During aerial application, human flaggers are prohibited unless in totally enclosed vehicles.

REENTRY STATEMENTS

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Warnings must include the following information: "WARNING - Area treated with Ridomil MZ58 on (date of application). Do not enter without appropriate protective clothing within 24 hours of application. In case of contact, flush skin or eyes with plenty of water; for eyes, consult a physician if irritation persists."

After sprays have dried, do not enter or allow entry into treated areas until the 24-hour re-entry interval has expired unless wearing the personal protective equipment listed on this label.

General Information

Ridomil MZ58 contains 10% of the active ingredient metalaxyl, and 48% of the active ingredient mancozeb. Metalaxyl is a systemic fungicide for use on selected crops to control diseases caused by members of the Phycomycete family of fungi. Mancozeb is effective against a wide range of fungal pathogens.

**Note:** Metalaxyl is a systemic fungicide having a specific mode of action and could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, CIBA-GEIGY cannot assume liability for crop damage resulting from resistant strains of fungi. If treatment is not effective following the use of Ridomil MZ58 as recommended, a resistant strain of fungi may be present. If the treatment is ineffective due to the presence of a metalaxyl resistant strain of fungi, neither Ridomil MZ58 nor any other fungicide with similar action will effectively control that disease. Consideration should then be given to the prompt use of other types of suitable fungicides. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance in your particular crop and disease control situation. DO NOT USE RIDOMIL MZ58 FOR DISEASE CONTROL IN CROPS GROWN IN GREEN-HOUSES, LATH HOUSES, OR IN BEDDING PLANT STRUCTURES.

Do not make applications when weather conditions favor drift from target area. Avoid spray overlap as crop injury may result.

Precaution: If this product is used on a crop, no other product containing a different EBDC active ingredient may be used on the same crop during the same growing season.

Mixing Instructions

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary to maintain tank mix uniformity. Maintain a maximum of agitation throughout the spraying operation. Flush spray tank thoroughly with water daily after use and dispose of pesticide rinsate by application to an already treated area.

Amount of water needed per acre will vary according to the amount of plant growth. For dilute ground applications, the water volume will vary from 20-150 gals./A. For aerial applications, 3-10 gals. of water/A are generally needed. To minimize the potential of drift from target area, applications at wind speeds greater than 10 mph are not recommended. DO NOT USE LESS THAN 5 GALLONS PER ACRE IN CALIFORNIA.

**BEST AVAILABLE COPY**

Precaution: Water-soluble prepacks of Ridomil MZ58 must be completely dissolved and dispersed in water before any other tank mix partner, including micronutrients or other liquid or dry fertilizers, are added to the spray solution.

Ridomil MZ58 Alone: Add 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble bags of Ridomil MZ58 into the tank all at once. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the pouches have dissolved, and the material has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Ridomil MZ58 + Tank Mixtures: Ridomil MZ58 is usually compatible with Bravo®, Dithane M-45, D.z.n®, Difolatan®, Manzate, and Manzate 200. Add 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble bags of Ridomil MZ58 into the tank all at once. Continue agitation while adding the remainder of the water. Allow the bags of Ridomil MZ58 to dissolve and the product to completely disperse into the mix water. Then add the desired amount of other products recommended for tank mixture and allow them to become completely dispersed. Continue agitation to maintain a uniform suspension until all of the spray solution has been applied.

Note: Do not let spray mixture stand overnight in the spray tank. Rinse spray tank at the end of the day.

#### Application Through Irrigation Systems

Ridomil MZ58 alone or in combination with other fungicides which are registered for sprinkler irrigation may be applied through irrigation systems.

Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**BEST AVAILABLE COPY**



Operating Instructions:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Application Instructions

Ridomil MZ58 must be applied on the schedule specified in the specific crop use recommendations, not according to the irrigation schedule. If irrigation schedules are used, ground or aerial applications must supplement chemigation applications to achieve adequate disease control.

With the exception of potatoes, Ridomil MZ58 has not been sufficiently tested to assure consistent product performance for all labeled uses when applied through chemigation systems. The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler irrigation equipment. Users should check with state and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

**BEST AVAILABLE COPY**

Center-Pivot Irrigation Equipment (Use only with drive systems which provide uniform water distribution)

1. Determine the size of the area to be treated.
2. Determine the time required to apply 1/4 inch water over the area to be treated when the system and injection equipment is operated at normal pressures recommended by the equipment manufacturer. Run the system at 80-95% of the manufacturer's rated capacity.
3. Using water, determine the injection pump output when operated at normal line pressure.
4. Determine the amount of Ridomil MZ58 required to treat the area covered by the irrigation system.
5. Add the required number of Ridomil MZ58 soluble bags and sufficient water to meet the injection time requirements to the solution tank. (See Mixing Instructions section of this label.)
6. Maintain constant solution tank agitation during the injection period.
7. Stop injection equipment after treatment is completed. Continue to operate the system until the Ridomil MZ58 solution has cleared the sprinkler head.

Solid Set, Hand Move and Moving Wheel Irrigation Equipment:

1. Determine the acreage covered by the sprinklers.
2. Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30 minute interval.
3. Determine the amount of Ridomil MZ58 required to treat the area covered by the irrigation system.
4. Add the required number of Ridomil MZ58 soluble bags into the same quantity of water used to calibrate the injection period. (See Mixing Instructions section of this label.)
5. Operate system at the same pressure and time interval established during the calibration.
6. Inject Ridomil MZ58 at the end of the irrigation cycle or as a separate application to maximize retention of the fungicide by the foliage.
7. Stop injection equipment after treatment is completed. Continue to operate the system under the Ridomil MZ58 solution has cleared the last sprinkler head.

1104

IMPORTANT

Do not allow Ridomil MZ58 to become wet or overheated during storage as this may lead to the soluble bag rupturing or to chemical changes which will impair the fungicidal effectiveness of Ridomil MZ58 and which may also generate flammable vapors.

Cucumbers, Melons and Summer Squash

Ridomil MZ58 provides effective control of downy mildew caused by Pseudoperonospora cubensis when used as a foliar spray in a preventative disease control program.

Apply 1 1/2-2 lbs. Ridomil MZ58 per acre in sufficient water to obtain thorough coverage. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until the threat of disease is over. Under heavy disease pressure, use the higher rate of Ridomil MZ58.

Notes: (1) To minimize the potential for resistance, do not make more than 4 applications of Ridomil MZ58 per crop. (2) If downy mildew continues to be a problem after making 4 applications of Ridomil MZ58, use other fungicides registered for downy mildew control. To avoid possible illegal residues, (3) Do not make the last application within 5 days of harvest and (4) If other registered fungicides containing mancozeb are used during the growing season, do not exceed a total of 19.2 lbs. of mancozeb active ingredient per acre.

Onions (Dry Bulb)

Ridomil MZ58 provides effective control of downy mildew caused by Peronospora destructor when used as a foliar spray in a preventative disease control program.

Apply 1 1/2-2 lbs. Ridomil MZ58 per acre in sufficient water to obtain thorough coverage. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until the threat of disease is over. Use a suitable spreader-sticker at rates recommended on the product label. Under heavy disease pressure, use the higher rate of Ridomil MZ58.

Notes: (1) To minimize the potential for resistance, do not make more than 4 applications of Ridomil MZ58 per crop. (2) If downy mildew continues to be a problem after making 4 applications of Ridomil MZ58, use other fungicides registered for downy mildew control. (3) Do not apply to exposed bulbs. To avoid possible illegal residues, (4) Do not make the last application within 7 days of harvest and (5) If other registered fungicides containing mancozeb are used during the growing season, do not

exceed a total of 24 lbs. of mancozeb active ingredient per acre.

Potatoes

Ridomil MZ58, when used as a foliar application in a preventative disease control program, will provide effective control of late blight and tuber rot caused by Phytophthora infestans, Pythium leak caused by Pythium spp, and pink rot caused by Phytophthora ethroseptica. Make applications of Ridomil MZ58 in sufficient water to obtain thorough coverage for ground applications and in a minimum of 3 gals. of water per acre for aerial applications.

It is recommended that this product be used within an Integrated Pest Management Program. Also, vine-kill should occur 14-days before harvest.

Late Blight: Apply 1 1/2-2 lbs. Ridomil MZ58 per acre. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until the threat of disease is over. Under heavy disease pressure, use the higher rate of Ridomil MZ58.

Tuber Rot, Pythium Leak and Pink Rot: Apply 2 lbs. Ridomil MZ58 per acre. Make the first application at flowering and another one 14 days later. If the field has a history of tuber disease problems, make a third application 14 days after the second application.

Note: If applications of Ridomil MZ58 are being made for late blight control that correspond to the timing of applications for tuber disease control, additional applications for tuber disease control are not needed.

For supplemental control of early blight, add 0.8 lb. of Manzate 200 80W, Dithane M-45 80W, Manzate 80W or Dithane M-22 80W to the Ridomil MZ58 spray or make an additional application of any registered early blight fungicide between each Ridomil MZ58 spray. Where early blight disease pressure is heavy, use early blight fungicides according to product use directions, and use Ridomil MZ58 only when late blight threatens. Do not apply more than 0.8 lb. per acre per application of Manzate 200 80W, Dithane M-45 80W, Manzate 80W, or Dithane M-22 80W in combination with Ridomil MZ58.

Notes: To avoid possible illegal residues, (1) Do not apply Ridomil MZ58 within 3 days of harvest in CT, FL, ME, MA, NH, NY, PA, VT and WI or at least 14 days elsewhere. (2) If other registered fungicides containing mancozeb are used during the growing season, do not exceed a total of 11.2 lbs. of mancozeb active ingredient per acre. (3) To minimize the potential for

BEST AVAILABLE COPY

resistance, do not make more than 4 applications of Ridomil MZ58 per crop. (4) If late blight is still a problem after making 4 applications of Ridomil MZ58, use other fungicides registered for late blight control. (5) Do not tank mix with amobam or Polyram®.

### Sugar Beets:

Ridomil MZ58, when used as a foliar application in a preventative disease control program, will provide effective control of downy mildew caused by Peronospora schachtii.

Apply 1-1/2 - 2 lbs. Ridomil MZ58 per acre. Begin application when conditions are favorable for disease, but before infection, and continue at 14-day intervals until the threat of disease is over. Under heavy disease pressure, use the higher rate of Ridomil MZ58.

Notes: (1) To minimize the potential for resistance, do not make more than four applications of Ridomil MZ58 per crop. To avoid possible illegal residues, (2) Do not make the last application within 14 days of harvest, (3) If other registered fungicides containing mancozeb are used during the growing season, do not exceed a total of 11.2 lbs. of mancozeb active ingredient per acre, and (4) Do not feed treated tops to livestock.

### Tomatoes

Ridomil MZ58, when used as a foliar application in a preventative disease control program, will provide effective control of late blight caused by Phytophthora infestans.

Apply 1 1/2-2 lbs. Ridomil MZ58 per acre as a foliar spray in sufficient water to provide thorough coverage. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until the threat of disease is over. Under heavy disease pressure, use the higher rate of Ridomil MZ58.

Note: When conditions are favorable for development of diseases other than late blight, any registered fungicides which control those diseases may be applied according to their label directions. When making applications of Manzate 200, Dithane M-45, Manzate, or Dithane M-22, 1 1/2 lbs./A of Ridomil MZ58 may be added to the spray mix every 14 days for control of late blight. Do not apply more than 2 lbs. per acre per application of Dithane M-45 80W, Manzate 200 80W, Dithane M-22 80W, or Manzate 80W in combination with Ridomil MZ58. Observe all precautions and restrictions that appear on the labels of other products applied with Ridomil MZ58.

Notes: To avoid possible illegal residues, (1) Do not apply Ridomil MZ58 within 14 days of harvest and (2) If other regis-

5151

tered fungicides containing mancozeb are used during the growing season, do not exceed a total of 16.8 lbs. of mancozeb active ingredient per acre on tomatoes grown east of the Mississippi River and 6.4 lbs. of mancozeb active ingredient on tomatoes grown west of the Mississippi River. (3) To minimize the potential for resistance, do not make more than 4 applications of Ridomil MZ58 per crop. (4) If late blight continues to be a problem after making 4 applications of Ridomil MZ58, use other fungicides registered for late blight control. (5) Do not tank mix with amobam, Polyram, nabam, thiram, zineb, or ziram.

Plant Back Restrictions

Do not plant any crop which is not registered for use with metalaxyl in metalaxyl-treated soil for a period of 12 months.

Planting Time From  
Last Application of  
Crop of Ridomil MZ58

Almonds	-0- Days
Alfalfa	-0- Days
Apples	-0- Days
Asparagus	-0- Days
Avocados	-0- Days
Blueberries	-0- Days
Broccoli	-0- Days
Cabbage	-0- Days
Cauliflower	-0- Days
Chinese Broccoli (gai lon, white flowering broccoli)	-0- Days
Chinese Cabbage (tight-heading varieties only)	-0- Days
Citrus	-0- Days
Cotton	-0- Days
Cucurbit Vegetables	-0- Days
Deciduous Fruits and Nuts	-0- Days
Hops	-0- Days
Head Lettuce	-0- Days
Legume Vegetables	-0- Days
Onions (dry bulb and seed only)	-0- Days
Papaya	-0- Days
Peanuts	-0- Days
Pineapples	-0- Days
Potatoes	-0- Days
Raspberries	-0- Days
Root and Tuber Vegetables	-0- Days
Soybeans	-0- Days
Spinach	-0- Days
Stone Fruits	-0- Days
Strawberries	-0- Days
Sugar Beets	-0- Days
Tobacco	-0- Days
Tomatoes	-0- Days
Walnuts	-0- Days
Wheat, Barley, Oats	12 Months
Corn	12 Months
Crops <u>Not Intended</u> for Food or Feed	-0- Days
Other Crops <u>Intended</u> for Food or Feed	12 Months

**BEST AVAILABLE COPY**

177218

Storage and Disposal

Storage - Store in a cool, dry place. Do not store this product under wet conditions.

Pesticide Disposal

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Disposal

Do not reuse outer container. Dispose of empty outer container in a sanitary landfill or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

Harmful if swallowed, inhaled or absorbed through the skin. Causes eye irritation. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. This product may cause allergic skin reaction in some individuals. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

Clothing and Equipment: All agricultural workers/handlers (i.e., mixers, loaders, and applicators) applying EBDC's must wear coveralls over long-sleeved shirt and long pants; shoes, socks, and chemical-resistant gloves. During mixing and loading, a chemical-resistant apron and goggles or a face shield must also be worn. Refer to Re-entry and Worker Protection Statements for additional Worker Safety rules.

BEST AVAILABLE COPY



Statement of Practical Treatment

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention.

If inhaled: Remove victim to fresh air. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Note to Physician: There is no specific antidote for Ridomil MZ58. Lavage stomach. Give a saline laxative and supportive therapy.

Environmental Hazards

This product is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Apply only as specified on this label. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Ridomil® trademark of CIBA-GEIGY for metalaxyl  
U.S. Patent No. 4,151,299

Bravo® trademark of ISK Biotech for chlorothalonil

Difolatan® trademark of Chevron Chemical Company for captafol

Dithane® trademark of Rohm and Haas Company for maneb/mancozeb

D·z·n® trademark of CIBA-GEIGY for diazinon

Manzate® trademark of E. I. duPont de Nemours & Company for maneb/mancozeb

Polyram® trademark of BASF Corporation for metiram

**BEST AVAILABLE COPY**

©1992 CIBA-GEIGY Corporation

Agricultural Division  
CIBA-GEIGY Corporation  
Greensboro, North Carolina 27419

	February 9, 1983
Revised	June 28, 1983
Revised	February 10, 1984
Revised	April 23, 1985
Revised	October 30, 1985
Revised	May 15, 1986
Revised	February 25, 1987
Revised	July 20, 1987
Revised	March 28, 1988
	(corrected Chinese cabbage)
Revised	April 19, 1989
	(water soluble bag)
Revised	September 18, 1989
	Metalaxyl FRSTR labeling
Revised	January 22, 1990
	Precautionary Statement additions
Revised	October 15, 1991
	New graphics
Revised	February 24, 1992 -
	cucurbits, sugar beets, hand
	move irrigation

CGA 65L22  
CGA 65L25

[GANNONC.LABELR]RIDOMIL-M258-F40 - 3/2/92