

(DRAFT)

Supplemental Labeling  
Dual® 25G Granular Herbicide

EPA Reg. No. 100-638

ACCEPTED  
Sept. 12, 1988  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, the pesticide  
registered under  
EPA Reg. No. 100-638

Active Ingredient:

Metolachlor: 2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl) acetamide .....	25.0%
Inert Ingredients:	75.0%
Total:	100.0%

Keep Out of Reach of Children


CAUTION

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

General Information

Dual 25G will control many annual grasses and certain broadleaf weeds in corn, peanuts, potatoes, and soybeans. It should be applied prior to the emergence of weeds. It may be preplant surface-applied (corn and soybeans only), or incorporated into the surface 1-2 inches of soil prior to planting or left on the soil surface during or after planting. Dry weather following application of Dual 25G may reduce its effectiveness. Shallow cultivate if weeds develop or if tillage is necessary to improve the physical condition of the soil.



Rotational Crops

(1) If crop treated with Dual 25G is lost, any crop on this label may be replanted immediately. Do not make a second broadcast application of Dual 25G. If the original application was banded and the second crop is planted in the untreated row middles, a second banded treatment may be applied. (2) Barley, oats, rye, or wheat may be planted 4 1/2 months following application; alfalfa may be planted 4 months following application. (3) Any crop on this label in addition to cotton, pod crops, safflower, grain sorghum\*, stone fruits, tree nuts, root crops, barley, buckwheat, milo, oats, rice, rye and wheat may be planted in the spring following treatment. Clovers may be seeded 9 months following application. Do not graze or feed forage or fodder from cotton to livestock. All other rotational crops may be planted 18 months after application. (4) Any crop on this label, in addition to cotton, pod crops, safflower, sorghum, stone fruits and tree nuts may be planted in the spring following a lay-by treatment or multiple treatments applied the previous season. All other rotational crops may be planted 18 months after a lay-by application.

\*provided the seed has been properly treated with Concep® seed protectant.

Important Notes: To avoid injury to rotational alfalfa or clover, (1) Do not apply more than 3 lbs. ai of metolachlor (12 lbs. Dual 25G) to corn, peanuts, potatoes, or soybeans either preplant surface, preplant incorporated, preemergence or postplant incorporated and (2) Do not make a lay-by or after-hilling application of Dual 25G.

Follow all applicable directions, restrictions, and precautions on the EPA-registered label.

This label must be in the user's possession at the time of application.

Dual® trademark of CIBA-GEIGY Corporation for metolachlor  
U.S. Patent No. 3,937,730

Agricultural Division  
CIBA-GEIGY Corporation  
P. O. Box 18300  
Greensboro, NC 27419

(LABELS-D.1 - DUAL25G/F5 - DRAFT)

February 8, 1988

