

Supplemental Labeling
Dual® 25G Granular Herbicide

EPA Reg. No. 100-638

Rotational Tomatoes

Active Ingredient:

Metolachlor: 2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl) acetamide	25.0%
Inert Ingredients:	75.0%
Total:	100.0%

Dual 25G is a granular herbicide.

KEEP OUT OF REACH OF CHILDREN

CAUTION


All applicable directions, restrictions and precautions on the EPA-registered Dual 25G label are to be followed.

This label must be in the possession of the user at the time of application.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.



Rotational Crops: (1) If crop treated with Dual 25G is lost, any crop on this label may be replanted immediately. Do not make a second broadcast application of Dual 25G. If the original application was banded and the second crop is planted in the untreated row middles, a second banded treatment may be applied. (2) Barley, oats, rye, or wheat may be planted 4 1/2 months following application; alfalfa may be planted 4 months following application. Tomatoes may be planted 6 months following application. (3) Any crop on this label in addition to cotton, cabbage, peppers, pod crops, safflower, grain sorghum*, stone fruits, tree nuts, root crops, tobacco, barley, buckwheat, milo, oats, rice, rye and wheat may be planted in the spring following treatment. Clover may be seeded 9 months following application. Do not graze or feed forage or fodder from cotton to livestock. All other rotational crops may be planted 12 months after application. (4) Any crop on this label, in addition to cotton, cabbage, peppers, tobacco, pod crops, safflower, grain sorghum*, stone fruits, and tree nuts, may be planted in the spring following a lay-by treatment or multiple treatments applied the previous season. All other rotational crops may be planted 12 months after a lay-by application.

*Provided the seed has been properly treated with Concep® seed protectant.

Important Notes: To avoid injury to rotational alfalfa or clover, (1) Do not apply more than 3 lbs. ai of metolachlor/A (12 lbs. Dual 25G) either preplant surface, preplant incorporated, preemergence, or postplant incorporated, and (2) Do not make a lay-by or after-hilling application of Dual 25G.

Dual® trademark of CIBA-GEIGY Corporation for metolachlor

©1992 CIBA-GEIGY Corporation

Agricultural Division
CIBA-GEIGY Corporation
P. O. Box 18300
Greensboro, NC 27419

Revised

July 12, 1989
April 30, 1992

[GANNONC.LABELD]DUAL25G-SUPP - 4/28/92