

PM25 100-640

File
10/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 27 1994

Theresa Kamicker
CIBA-GEIGY CORP.
P. O. Box 18300
Greensboro, NC 27419

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 09/13/93 in Response to PR Notice 93-7
EPA Reg. No. 100-640
PRIME +

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber

2 of 15


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

3015

WPS
(Front Cover)

Prime +Ⓢ

A Plant Growth Regulator for
Control of Tobacco Suckers

One Gallon
U. S. Standard Measure

or

One Quart
U.S. Standard Measure

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 27 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

100-640

Active Ingredient:	
2-chloro-N-[2,6-dinitro-4-	
(trifluoromethyl)phenyl]-N-ethyl-	
6-fluorobenzenemethanamine	15%
Inert Ingredients:	85%
Total:	100%

Prime + contains 1.2 lbs. active ingredient
per gallon.

EPA Reg. No. 100-640
EPA Est. 100-LA-1

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se
la explique a usted en detalle. (If you do not understand the
label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use
inside booklet.

CGA 73L9H 093 (1 Gal.)
CGA 73L11G 093 (w/o est. no.) (1 Gal.)
CGA 73L10B 093 (1 Qt.)

Ciba-Geigy

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. Ciba-Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.

5045

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR SUCKER CONTROL AND/OR CROP INJURY.

Do not apply this product through any type of irrigation system.

General Information

Prime + is a plant growth regulator for the control of sucker growth on flue-cured, burley, dark, Maryland, and cigar tobacco.

One application of Prime + will provide full season control of excessive sucker development in tobacco. Read this label carefully and observe all directions and precautions.

Mixing Instructions

Mix 1 gal. of Prime + in 49 gals. of water or 2.5 fl. oz. Prime + per one gal. of water. Provide sufficient agitation during mixing and application to maintain a uniform mixture.

Note: Do not mix Prime + with insecticides or fungicides.

Application Instructions

Time of Application

Apply Prime + when most tobacco plants are in the elongated button to early flower stage, either before or after topping. If Prime + is applied after topping, apply within 24 hours of topping. During topping, remove all suckers one inch or longer from all plants. (See notes listed below.)

Precision Hand Application

Mix 1 gal. of Prime + in 49 gals. of water and apply as a coarse spray or drench to the top of the tobacco stalk using a hand-held drop line, or mix 2.5 fl. oz. of Prime + per gal. of water when using the jug method or backpack sprayer. Apply only once per plant per season. Apply the spray mixture at a rate sufficient to run down the stalk contacting each leaf axil and just reach the ground line. Excessive spray volume may result in carryover injury to rotational crops. Spray volume per plant will generally be 1/3-2/3 fl. oz. depending on plant height. Do not exceed 30 gals. spray volume per acre when using hand application methods. (See notes listed below.)

Application with Powered Spray Equipment (3 nozzles per row)

Mix 1 gal. of Prime + in 49 gals. of water and apply at a rate of 30 to 50 gals. spray volume per acre. Use the high spray volume in tall types of tobacco, such as flue-cured, burley, Maryland, and cigar. Use the low spray volume only in short types of tobacco, such as dark. Apply only once per season. Avoid application to plants treated by precision hand application method. Spray volume should be sufficient to run down the

70/15

stalk contacting each leaf axil and just reach the ground line. Excessive spray volume beyond the point of runoff to the soil may result in carryover injury to rotational crops. (See notes listed below.)

Apply Prime + as a coarse spray directed downward to the top of the tobacco stalk. This is best accomplished with three full cone nozzles per row; one nozzle directed downward over the center of the row and one 7-9 in. to each side of the center, directed at the top of the stalk. Maintain nozzles at a height of 12-16 in. above the top of the stalk during application. Maintain boom pressure at 20-25 psi, not exceeding 30 psi.

Sequential Applications with Maleic Hydrazide (MH)

Prior to Use of Maleic Hydrazide

Mix 2 qts. of Prime + in 49.5 gals. of water and apply either by Precision Hand Application or Application with Powered Spray Equipment as described in those sections of this label. Timing of Prime + should be during the elongated button to early flower stage of tobacco.

Approximately one week after the Prime + Application, apply maleic hydrazide according to the directions, precautions, and restrictions on that label.

After Use of Maleic Hydrazide

For extended control of tobacco suckers, particularly during late tobacco harvests in FL, GA, NC, SC and VA.

First, apply the regular schedule of contacts. Follow with the labeled rate of maleic hydrazide in accordance with its label. Follow with 2 qts. of Prime + in sufficient water to total 50 gals. to treat one acre 3-4 weeks after applying maleic hydrazide, depending on the development of suckers. To increase the effectiveness of Prime +, add 0.25% of X-77@ (1 pt./50 gals.) to the spray solution.

Apply Prime + as a coarse spray using powered spray equipment (3 nozzles per row) at least 7 days before the next leaf harvest. The spray volume should be sufficient to run down the stalk contacting each leaf axil where suckers are present. Refer to the Application with Powered Spray Equipment section for specific instructions.

Notes: 1) Apply Prime + only once per tobacco season.
2) Prime + will not control large suckers. All suckers one inch or longer on all tobacco plants must be removed before application of Prime +. 3) Follow all precautions, notes and other instructions on this label.

8045

Tank Mix with Maleic Hydrazide (MH)

Mix 2 qts. of Prime + with labeled rate of maleic hydrazide in sufficient water to total 50 gals. and apply at the recommended time for application of maleic hydrazide. Apply the mixture as a coarse spray at a volume of 50 gals. per acre with a boom pressure maintained at 20-25 psi by Powered Spray Equipment as described in that section of this label. For best results, follow precautions and restrictions on the Prime + and maleic hydrazide labels.

Application sequence: 1) Apply contact material according to its label. 2) Apply the tank mixture of Prime + and maleic hydrazide at or near the time suggested for application of maleic hydrazide on its label. 3) Make only one application of the tank mixture of Prime + plus maleic hydrazide per tobacco crop.

Notes

1. Timing is important; applications should be made at timing described above so that number and size of suckers will be small. Applications made before the elongated button stage of growth may result in chemical topping or distortion of leaves that were too immature at time of application. Spraying too late may result in poor sucker control.
2. Prime + will not control large suckers. All suckers one inch or longer on all plants must be removed before application of Prime +. Suckers that escape treatment should be removed by hand about 10-14 days after application.
3. Where a contact (fatty alcohol) is commonly used to control early suckers and provide a more uniform crop, the normal sequence would be: (a) apply contact material according to its label, and (b) about 5 days after application of contact material, apply Prime +. Timing of Prime + application should be during the elongated button to early flower stage.
4. Where a contact (fatty alcohol) is not used and a wide range of flowering exists, early flowering plants should be topped and suckers removed by hand. Then, when most plants are in the elongated button to early flower stage, remove all suckers one inch or longer, and apply Prime +. If Prime + is applied using precision hand application, early flowering plants can be topped and treated selectively; later flowering plants can be topped and treated as they reach the proper treatment stage; and a contact may not be needed.

9/15

5. For proper sucker control, Prime + must contact the small suckers on the axil where the leaf and stalk connect. Prime + will not control suckers on plants that are leaning at time of application. Applications with powered spray equipment during windy periods are not recommended. High wind will disrupt the spray pattern of nozzles on boom sprayers and turn leaves causing improper contact and flow of the spray solution, resulting in poor sucker control. Application to wet plants, or drought stressed plants may result in poor sucker control.
6. Carefully follow the use rates and spray volume recommended, and spray field uniformly. Volume should be sufficient to run down the stalk, contact each leaf axil, and just reach the ground line.
7. Rain occurring within 2 hours after spraying may reduce effectiveness.

Precautions: 1) To avoid illegal residues and possible injury to crops planted in the fall or following spring, do not apply excessive spray volume to the point of runoff onto the soil. Prime + residues in the soil may interact with dinitroaniline herbicides such as Balan®, Basalin®, Paarlan®, Prowl®, or Treflan® to increase early season crop stunting. Thorough disking and deep tillage in the fall are recommended where excessive amounts of Prime + may have reached the soil. 2) When tobacco is in a state of very rapid, succulent growth due to high nitrogen and rainfall, Prime + may cause injury. Under these conditions, use only the precision hand application method and avoid contacting leaves with the spray solution.

Storage and Disposal

Pesticide Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product are acutely hazardous. Improper disposal of unused pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Disposal

Do not reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

100415

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372 day or night.

Store at temperatures above 32°F.

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER/PELIGRO

Corrosive - causes eye damage. Causes skin irritation or skin sensitization on repeated contact. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist.

First Aid: In case of contact with the eyes, immediately flush with plenty of water and get medical attention. In case of contact with skin, wash immediately with soap and water. If swallowed, drink plenty of water. Do not induce vomiting. Avoid alcohol. Call a physician immediately. In case of inhalation exposure, move from the contaminated area.

Note to Physician: There is no specific antidote for Prime +. Lavage stomach. Give a saline laxative or slurry of activated carbon in water and supportive therapy.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to lakes, streams, or ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Prime +® trademark of Ciba-Geigy
U.S. Patent No. 4,169,721

Balan®, Paarlán® and Treflan® trademarks of DowElanco

Basalin® trademark of BASF/Wyandotte Corp.

Prowl® trademark of American Cyanamid

X-77® trademark of Chevron Chemical Company

©1993 Ciba-Geigy Corporation

Ciba Plant Protection
Ciba-Geigy Corporation
Greensboro, North Carolina 27419

CGA 73L9H 093 (1 Gal.)
CGA 73L11G 093 (w/o est. no.) (1 Gal.)
CGA 73L10B 093 (1 Qt.)

Ciba-Geigy		August 17, 1983
	Revised	February 3, 1984
	Revised	December 8, 1984
	Revised	September 4, 1985
	Revised	May 12, 1987
	Revised	November 20, 1989
	Revised	April 25, 1991
	Revised	January 17, 1992
	Revised	September 2, 1993
	Revised	September 8, 1993

[GANNONC.LABELP] - PRIMEWPS - 9/8/93

(Back Cover)

Prime +⊙

A Plant Growth Regulator for
Control of Tobacco Suckers

Active Ingredient:

2-chloro-N-[2,6-dinitro-4-(trifluoromethyl)phenyl]-N-ethyl-6-fluorobenzenemethanamine	15%
---	-----

Inert Ingredients:	85%
--------------------	-----

Total:	100%
--------	------

Prime + contains 1.2 lbs. active ingredient per gallon.

See directions for use in attached booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-640
EPA Est. 100-LA-1

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

©1993 Ciba-Geigy Corporation Corporation

One Gallon
U.S. Standard Measure

or

One Quart
U.S. Standard Measure

Precautionary Statements

Hazards to Humans and Domestic Animals

Corrosive - causes eye damage. Causes skin irritation or skin sensitization on repeated contact. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist.

First Aid: In case of contact with the eyes, immediately flush with plenty of water and get medical attention. In case of contact with skin, wash immediately with soap and water. If swallowed, drink plenty of water. Do not induce vomiting. Avoid alcohol. Call a physician immediately. In case of inhalation exposure, move from the contaminated area.

Note to Physician: There is no specific antidote for Prime +. Lavage stomach. Give a saline laxative or slurry of activated carbon in water and supportive therapy.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to lakes, streams, or ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply this product through any type of irrigation system.

STORE AT TEMPERATURES ABOVE 32°F

Prime +® trademark of Ciba-Geigy Corporation
U.S. Patent No. 4,169,721

©1993 Ciba-Geigy Corporation

Ciba Plant Protection
Ciba-Geigy Corporation
Greensboro, North Carolina 27419

CGA 73L9H 093 (1 Gal.)
CGA 73L11G 093 (w/o est. no.) (1 Gal.)
CGA 73L10B 093 (1 Qt.)

Ciba-Geigy

	August 17, 1983
Revised	February 3, 1984
Revised	December 7, 1984
Revised	September 4, 1985
Revised	May 12, 1987
Revised	November 20, 1989
Revised	April 25, 1991
Revised	January 17, 1992
Revised	September 2, 1993
Revised	September 8, 1993

[GANNONC.LABELP]PRIMEWPS - 9/8/93