

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Ms. Eileen D. King-Watson
Agricultural Division
Ciba-Geigy Corporation
P.O. Box 18300
Greensboro, NC 27419-8300

FEB 5 1992

Dear Ms. King-Watson:

Subject: Amended Labeling
Banner
EPA Registration No. 100-641
Your Submission Dated September 20, 1991

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable provided that you:

- 1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. For tulip bulbs, indicate the length of the dip treatment.
 - b. Under Environmental Hazards, delete the third sentence. The following sentence is intended to replace this sentence.
- 2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Additionally, we are awaiting a review from the Product Chemistry Review Section concerning your revised Confidential Statement of Formula submitted with the submission referred to above.

Sincerely yours,

Susan T. Lewis

Susan T. Lewis
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (H7505C)

BEST AVAILABLE COPY

Enclosure

CONCURRENCES

H7505C:J. Fairfax:100-641:1/16/92

SYMBOL	SURNAME	DATE					

PM 21 100-641

Highlighted copy 2/18

MASTER LABEL BOOKLET
(For Update Purposes Only)

ACCEPTED
with COMMENTS
by EPA Office of Pesticides

FEB 06 1992

Under the provisions of the
Fungicide, and other Pesticide Act,
and related laws, this pesticide
registered under the Act.

100-641

BANNER®

FUNGICIDE

Broad spectrum and systemic
disease control for turf and
ornamentals

Active Ingredient:

Propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-
1,3-dioxolan-2yl]methyl]-1H-1,2,4-

triazole 14.3%

Inert Ingredients: 85.7%

Total: 100.0%

EPA Reg. No. 100-641

EPA Est. 100-AL-1

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

PRECAUCIÓN AL USUARIO: Si usted no lee inglés,
no use este producto hasta que la etiqueta haya
sido explicado ampliamente.

See additional precautionary statements
and directions for use inside booklet.

One Quart (CGA)

One Gallon (CGA)

U.S. Standard Measure

CIBA-GEIGY

BEST AVAILABLE COPY

[GANNONC.LABELB] - [BANNER-DRAFT] - 10/01/91

Added
ISO-approx
Common name

ac 2d
Spanish

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

Reentry Statement

Do not apply this product in such a manner as to directly or through drift expose workers or other persons, except those knowingly involved in the application. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must inform workers of areas or fields that may not be entered without specific protective clothing until sprays have dried, and appropriate actions to take in case of accidental exposure, as described under Precautionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "WARNING. Area treated with Banner on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental eye contact, flush eyes with plenty of water. Get medical attention. In case of contact with skin, wash thoroughly with soap and water. Call physician if irritation occurs. Remove and wash contaminated clothing before reuse."

Added

General Information

Banner is a systemic fungicide for use on turfgrasses for the control of dollar spot (*Sclerotinia homoeocarpa*), brown patch (*Rhizoctonia solani*), anthracnose (*Colletotrichum graminicola*), red thread (*Corticium fuciforme*), rust (*Puccinia graminis*), powdery mildew (*Erysiphe graminis*), stripe smut (*Ustilago striiformis* and *Urocystis agropyri*), summer patch (*Magnaporthe poae*), spring dead spot (*Leptosphaeria korrea*), gray leafspot (*Pyricularia grisea*), pink snowmold (*Fusarium nivale*) and gray snowmold (*Typhula* spp.).

Banner also controls numerous diseases on ornamentals and other landscape and nursery plantings. It controls powdery mildews, rusts, leafspots, and blights. Refer to the appropriate section for specified diseases and plants.

Do not use on home lawns. Do not apply this product through any type of irrigation system. Do not use Banner as a tree injection.

} added

Mixing Instructions

Fill the spray tank 1/2-3/4 full with water, add the proper amount of Banner, then add the rest of the water. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

If Banner is tank mixed with other products, use the following sequence:

1. Always check the compatibility of the tank mix using a jar test with proportionate amounts of Banner, other chemicals to be used, and water, before mixing in the spray tank.
2. Provide sufficient jet or mechanical agitation during filling and application to keep the tank mix uniformly suspended.
3. Fill tank at least 1/2 full of clean water.
4. Add wettable powders to the tank first, allowing them to completely suspend in the tank before proceeding. This process can be hastened by premixing the product in water before adding to the tank.
5. Add flowables or suspensions next.
6. Add Banner and other emulsifiable concentrates last.
7. Do not leave tank mix combinations in the spray tank for prolonged periods without agitation. Mix and apply them the same day.

Turfgrass and Dichondra Disease Control

1. USE BANNER IN A PREVENTATIVE DISEASE CONTROL PROGRAM.
2. Apply in sufficient water to ensure thorough coverage.
3. Apply after mowing OR allow sprayed area to completely dry before mowing.
4. For control of foliar diseases, allow sprayed area to completely dry before irrigation.

- 5. For control of soil-borne diseases, Banner can be watered in after application.
- 6. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
- 7. For optimum turf quality and disease control, use Banner in conjunction with turf management practices that promote good plant health and optimum disease control.
- 8. Addition of surfactants may result in phytotoxicity.
- 9. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

CAUTION: Bermudagrass can be sensitive to Banner. Do not exceed 4 fl. ozs. per 1,000 sq. ft. every 30 days on any variety of bermudagrass. In Florida, do not apply Banner to Bermudagrass golf course greens when temperatures exceed 90°F.

NOTE: Do not graze animals on treated areas. Do not feed clippings from treated areas to livestock or poultry.

Tank Mixes

For broader spectrum control, Banner can be tank mixed with other fungicides. For example, Subdue® or Pace® may be tank mixed with Banner or used alone when conditions are favorable for Pythium blight. Banner is also compatible with numerous herbicides and insecticides. Check compatibility before tank mixing. Add Unite® (3 pts./100 gals.) to tank mixes which are incompatible. Do not mix Banner and Tersan® 1991 in a fertilizer solution. Follow the directions under "Mixing Instructions" for tank mixes. Observe all directions, precautions, and limitations on labeling of all products used in tank mixes.

} added

Turfgrass - Specific Diseases, Rates and Application Timing

<u>Disease</u>	<u>Fl. Oz. Per 1,000 sq. ft.</u>	<u>Application Interval/ Timing</u>	<u>Instructions</u>
Dollar Spot	0.5	7-14 days	Apply when conditions are favorable for disease development.
	1-2	14-28 days	
Brown Patch	2-4	10-21 days	Begin applications in May or June before disease is present. If disease is present, mix 2 fl. oz. Banner per 1,000 sq. ft. with a registered contact fungicide at the labeled rate. Under conditions of high temperature and high humidity, use the higher rates and shorter intervals.
Powdery Mildew, Rust, Anthracnose	1-2	14-28 days	Apply when conditions are favorable for disease development.
Red Thread	2	14-21 days	Apply when conditions are favorable for disease development.

Disease	Fl. Oz. Per 1,000 sq. ft.	Application Interval/ Timing	Instructions
Stripe Smut	1-2	Fall or Spring	Apply once in the fall after grass becomes dormant or in the early spring before grass starts to grow.
Gray Leafspot (St. Au- gustinegrass sod produc- tion)	2	14 days	Apply when conditions are favorable for disease development.
Melting Out, Leaf Spot	2-4	14 days	Under light to moderate pressure, apply Banner to reduce the severity of leaf spot and melting out caused by Helminthosporium type pathogens.
Summer Patch, Poa Patch	4	30 days	Apply Banner at monthly intervals beginning in April. Make 2-3 applications depending on local recommendations.
Spring Dead Spot (California Only)	4	30 days	Apply Banner at monthly intervals beginning in August. Make three applications.

<u>Disease</u>	<u>Fl. Oz. Per 1,000 sq. ft.</u>	<u>Application Interval/ Timing</u>	<u>Instructions</u>
Snowmold			
Gray - <u>Typhula</u> spp.	4	Late Fall	Apply one appli- cation in the late fall before snow cover. Do not apply on top of snow.
Pink - <u>Fusarium</u> <u>nivale</u>			

Dichondra - Specific Diseases, Rate and Application Timing

<u>Disease</u>	<u>Fl. Oz. Per 1,000 sq. ft.</u>	<u>Application Interval/ Timing</u>	<u>Instructions</u>
Dichondra Rust	2	14-21 days	Apply when condi- tions are favorable for disease development.

Establishment of Cool Season Turfgrass

Banner provides control of many diseases of turf, and its primary use is as a fungicide for use against the diseases listed on this label. As an additional benefit, Banner will improve the rate of establishment when it is applied to cool season grass seedlings or sod.

New Seedlings: Apply 1 oz./1,000 sq. ft. at the two to three leaf stage of growth for faster root development and top growth.

Sod: Apply 1 oz./1,000 sq. ft. 2-6 weeks before cutting for increased sod knitting and rate of establishment after setting.

} added

Disease Control in Nurseries (Field) and Landscape Plantings

1. USE BANNER IN A PREVENTATIVE DISEASE CONTROL PROGRAM. INITIATE APPLICATIONS ACCORDING TO LOCAL RECOMMENDATIONS OR AT FIRST SIGNS OF DISEASE.
2. Apply in enough water to thoroughly cover foliage to the point of runoff.

- 3. Allow spray to dry before overhead irrigation is applied.
- 4. Optimum benefit of Banner is obtained when used in conjunction with sound disease management practices.

Note: Banner has been shown to be safe on the species of plants and effective in controlling the diseases listed in the Directions For Use. Other plant species may be sensitive to Banner and other diseases may not be controlled. Before using Banner on plants or for diseases that are not listed in the Directions For Use, test Banner on a small scale basis first. Do not apply Banner to African violets, begonias, Boston fern, or geraniums. Apply the recommended rates for a particular type of disease, i.e., rust, powdery mildew, etc., and evaluate for phytotoxicity and disease control prior to widespread use.

Relocated
from General
Information

Bedding and Flowering Plants

<u>Species</u>	<u>Disease</u>	<u>Recommended Rates and Timing</u>
Calendula	Powdery Mildew (<u>E. cichoracearum</u>)	Apply 5 fl. oz. per 100 gals. every 30 days.
Chrysanthemums	Ray Blight (<u>Ascochyta chrysanthemi</u>)	Apply 12 fl. oz. per 100 gals. at 7-day intervals or 20 fl. oz. per 100 gals. at 14-day intervals.
Delphinium	Powdery mildew (<u>Erysiphe sp.</u>)	Apply 5 fl. oz. per 100 gals. every 10-14 days.
Roses (Field-grown only)	Powdery Mildew (<u>S. pannosa</u>)	Apply 5 fl. oz. per 100 gals. on a 10-14 day interval. Thorough coverage is needed for best results. Begin applications when conditions are favorable for disease development.
	Rust (<u>Phragmidium spp.</u>)	Apply 5-10 fl. oz. per 100 gals. on a 10-14 day interval. Use the highest rate and shortest interval when conditions for disease are severe.
	Blackspot (<u>D. rosea</u>)	Tank mix 5 fl. oz. per 100 gals. with a registered contact fungicide labeled for blackspot control.

<u>Species</u>	<u>Disease</u>	<u>Recommended Rates and Timing</u>
Snapdragon	Rust (<u>Puccinia sp.</u>)	Apply 10 fl. oz. per 100 gals. every 7-10 days.
Sweet William (<u>Dianthus barbatus</u>)	Leafspot (<u>Heterosporium echinulatum</u>)	Apply 6 fl. oz. per 100 gals. every 14 days.
Tulip Bulbs	Blue Mold (<u>Penicillium sp.</u>)	Dip bulbs in a suspension of 24-48 fl. oz. per 50 gals. ¹ Allow bulbs to dry before placing in storage.

} added

¹Dispose of waste by using for another purpose as specified on this labeling or according to the instructions given in the Storage and Disposal section.

Woody Plants

<u>Species</u>	<u>Disease</u>	<u>Recommended Rates and Timing</u>
Crabapple	Scab (<u>Venturia inaequalis</u>)	Apply 2 fl. oz. per 100 gals. every 14 days. Begin at green tip.
	Rust (<u>G. juniperi</u>)	Apply 2 fl. oz. per 100 gals. every 7 days. Make 3 applications beginning at green tip.
	Powdery Mildew	Apply 6-8 fl. oz. per 100 gals. every 14 days.
Crape Myrtle	Powdery Mildew (<u>E. lagerstroemia</u>)	Apply 16 fl. oz. per 100 gals. every 21 days.
Dogwood	Anthraxnose (<u>Discula spp.</u>)	Apply 2-4 fl. oz. per 100 gals. every 14 days or apply 8 oz./100 gal. every 28 days.
Douglas Fir	Needle Rust (<u>Melampsora occidentalis</u>)	Apply 16 fl. oz. per 100 gals. Apply once in May.
Jeffrey Pine	Tip blight (<u>Sirococcus strobolinus</u>)	Apply 16 fl. oz. per 100 gals. Initiate applications in mid-late winter (Nov.-Jan.) Apply 3 times at 2-month intervals.
Oaks	Leafspot (<u>Tubakia dryina</u>)	Apply 16 fl. oz. per 100 gals. every 14 days.

} added

} added

<u>Species</u>	<u>Disease</u>	<u>Recommended Rates and Timing</u>
Red Tip Photinia	Leafspot (<u>Fabraea maculata</u> syn. <u>Entomosporium maculatum</u>)	Apply 12-20 fl. oz. per 100 gals. every 7-14 days. Reductions of plant height may occur with Banner applications, especially on plants 6 inches or smaller.
Shasta Fir	Needle Rust (<u>Pucciniastrum goeppertianum</u>)	Apply 16 fl. oz. per 100 gals.

} Added

Non-Bearing Fruits and Nuts
(Nurseries and Landscape Plantings)

<u>Species</u>	<u>Disease</u>	<u>Recommended Rates and Timing</u>
Apple	Scab (<u>Venturia inaequalis</u>)	Apply 2 fl. oz. per 100 gals. beginning at green tip. Apply every 14 days.
	Rust (<u>G. juniperi</u>)	Apply 2 fl. oz. per 100 gals. every 7 days. Make 3 applications.
	Powdery Mildew (<u>P. leucotricha</u>)	Apply 6-8 fl. oz. per 100 gals. every 14 days.
Cherry	Powdery Mildew	Apply 2-4 fl. oz. per 100 gals. on a 14-21 day interval. Apply 4 fl. oz. per 100 gals. on a 14-21 day interval.
	Leafspot (<u>Coccomyces hiemalis</u>)	
	Blossom Blight (<u>Monilinia fructicola</u>)	
Citrus	Greasy Spot (<u>Mycosphaerella citri</u>)	Apply 20-24 fl. oz. per 100 gals.
Peach Plum Nectarine	Blossom Blight (<u>M. fructicola</u> or <u>M. laxa</u>)	Apply 3-4 fl. oz. per 100 gals. Apply at 5-10% bloom and again at 70-100% bloom.

} Rates/100 gals rather than 1A

} Rates/100 gal rather than 1A

} Added

} Rates/100 gal rather than per A.

<u>Species</u>	<u>Disease</u>	<u>Recommended Rates and Timing</u>
Pecan	Scab	Apply 12 fl. oz. per 100 gals. beginning at bud break. Apply 3 times on a 14-day interval.
	(Cladosporium)	
	Powdery Mildew	
	Downy Spot	
	Zonate Leafspot	
Walnut	Fungal Leaf Scorch	Apply 8.5 fl. oz. per 100 gals. at 14-21 day intervals.
	Brown Spot	
	Anthracnose	

Rates / 100 gals rather than 1A

Note: To avoid possible illegal residues, do not apply to apple, cherry, citrus, peach, plum, nectarine, pecan or walnut trees that will bear harvestable fruit within 12 months.

Storage and Disposal

Removed 0°F minimum storage temp. restriction.

Pesticide Disposal

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Disposal

Do not reuse empty container. Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. Stay out of smoke from burning containers.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372 day or night.

Revised

14919

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING/AVISO

Causes substantial but temporary eye injury. Wear goggles or face shield. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapor or spray mist. Wear rubber gloves and a long-sleeved shirt when mixing, handling, and spraying the product. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

Statement of Practical Treatment

Revised

If in eyes: Immediately flush eye with a steady, gentle stream of water. Get medical attention.

If on skin: Wash thoroughly with soap and water. Get medical attention if irritation occurs.

If swallowed: Do not induce vomiting. Drink plenty of water and contact a physician, hospital or local Poison Control Center.

If inhaled: Move victim to fresh air.

Notes to Physician: If ingested, lavage stomach to avoid aspiration. A slurry of activated charcoal in water can be left in the stomach. Give a saline laxative and supportive therapy.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment wash waters.

} added

Physical or Chemical Hazards

Do not use or store near heat or open flame.

Subdue® and Pace® trademarks of CIBA-GEIGY for metalaxyl

Tersan® 1991 trademark of E. I. duPont de Nemours and Co., Inc.
for benomyl

Unite® trademark of Hopkins Chemical Co.

©1991 CIBA-GEIGY Corporation

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419

CGA (QUART)
CGA (GALLON)

Product ID: 58010
Product ID: 52840

This booklet manufactured using post-consumer, recycled paper.

[GANNONC.LABELB]BANNER-DRAFT - 10/01/91

Booklet Back Panel

BANNER®

FUNGICIDE

Broad spectrum and systemic disease control for turf and ornamentals.

Added ISO approved Common name

Active Ingredient:

(Propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2yl]methyl]-1H-1,2,4-triazole 14.3%

Inert Ingredients: 85.7%

Total: . 100.0%

See directions for use in attached booklet.

EPA Reg. No. 100-641
EPA Est. 100-AL-1

Banner® trademark of CIBA-GEIGY
U.S. Patent No. 4,079,062

©1991 CIBA-GEIGY Corporation

One Quart
One Gallon

U.S. Standard Measure

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

ac 2d Spanish

PRECAUCIÓN AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes substantial but temporary eye injury. Wear goggles or face shield. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapor or spray mist. Wear rubber gloves and a long-sleeved shirt when mixing, handling, and spraying the product. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

BEST AVAILABLE COPY

Statement of Practical Treatment

Revised

- If in eyes: Immediately flush eyes with a steady, gentle stream of water. Get medical attention.
- If on skin: Wash thoroughly with soap and water. Get medical attention if irritation occurs.
- If swallowed: Do not induce vomiting. Drink plenty of water and contact a physician, hospital or local Poison Control Center.
- If inhaled: Move victim to fresh air.

Notes to Physician: If ingested, lavage stomach to avoid aspiration. A slurry of activated charcoal in water can be left in the stomach. Give a saline laxative and supportive therapy.

Environmental Hazards

added

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment wash waters.

Physical or Chemical Hazards

Do not use or store near heat or open flame. In the event of spill, fire, or other emergency, call 1-800-888-8372 day or night.

← Removed 0°F minimum storage temperature restriction.

added

Chemigation

Do not apply this product through any type of irrigation system.

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419

CGA (QUART)
CGA (GALLON)

Product ID: 58010
Product ID: 52840

CIBA-GEIGY

- June 16, 1987
- Revised March 3, 1988
- May 8, 1989
- August 15, 1989
- September 27, 1989
- May 16, 1990
- August 8, 1991

BEST AVAILABLE COPY

11/2/19.

Revisions to typed label:

3/3/88

Added claims for summer patch, spring dead spot, and dichondra rust. Deleted claim for aid in control of fusarium blight; revision of use directions to specify rates on each disease, double maximum rate to 4 fl. oz./1000 sq. ft., allow use on bermudagrass and St. Augustine grass, add Tank Mixes section; added "Do not contaminate water when disposing of equipment wash waters;" added trademark statements for six other products.

8/15/89

Added claims for gray leafspot, pink snowmold and gray snowmold in turf section, added ornamentals section and made changes in general section parallel to those approved for Tilt.

9/27/89

Added one-quart size to master label copy.

5/16/90

Revised storage statement (0° F lower limit).

8/8/91

Added claims for turfgrass establishment, disease control on tulip bulbs, dogwood, oaks, red tip Phctinia and citrus.