


WATER SOLUBLE BAG

Ridomil®/Bravo® 81W

Fungicide

For the control of certain diseases in broccoli, cabbage, cauliflower, cucurbit vegetables, onions, potatoes and tomatoes.



Active Ingredients:

Metalaxyl: N-(2,6-dimethylphenyl)-	
N-(methoxyacetyl)alanine methyl ester	9.0%
Chlorothalonil: Tetrachloroisophthalonitrile.....	72.0%
Inert Ingredients:	19.0%
Total:	100.0%

Ridomil/Bravo 81W is a wettable powder.

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

PRECAUCIÓN AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

Statement of Practical Treatment

If in eyes: Flush with plenty of water and get medical attention.

If swallowed: Induce vomiting by touching back of throat with finger. Drink promptly a large quantity of milk, egg whites, or gelatin solution. If none of these are available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention immediately.

Continued ... 2

1715

If on skin: Wash promptly with plenty of soap and water. Rinse thoroughly. Get medical attention.

Note to Physician

If swallowed, probable mucosal damage may contraindicate the use of gastric lavage.

If in eyes, specialized ophthalmologic attention may be required.

See additional precautionary statements on left side of container.

See directions for use on right side of container.

EPA Reg. No. 100-658

EPA Est. 10163-AZ-1

This outer protective container contains Ridomil/Bravo 81W in an inner water-soluble bag. Entire inner bag and contents dissolve in water. After opening outer container, immediately dump entire unopened inner bag into the partially filled sprayer or mix tank. Do not handle the soluble bag or expose it to moisture, since this may cause rupturing.

Five Pounds
Net Weight

Ten Pounds
Net Weight

CIBA-GEIGY

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons, except those knowingly involved in the application. The area being treated must be vacated by unprotected persons.

Reentry Statements

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in the treated area or in an area about to be treated with this product. Oral warnings must be given which inform workers of areas or fields that may not be entered without specific protective clothing until sprays have dried and appropriate actions to take in case of accidental exposure, as described under Precautionary Statements on the back of this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "DANGER. Area treated with Ridomil/Bravo 81W on (date of application). Do not enter without appropriate protective clothing until sprays have dried. Flush eyes or skin with plenty of water. Call a physician if irritation persists. Remove contaminated clothing and wash before reuse."

General Information

Ridomil/Bravo 81W contains 9% of the active ingredient metalaxyl and 72% of the active ingredient chlorothalonil. Metalaxyl is a systemic fungicide that provides control of downy mildew and late blight diseases of selected crops. Chlorothalonil is a broad-spectrum protectant fungicide that controls many diseases of vegetables.

Note: Metalaxyl is a systemic fungicide having a specific mode of action and could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, CIBA-GEIGY cannot assume liability for crop damage resulting from resistant strains of fungi. If treatment is not effective following the use of Ridomil/Bravo 81W as recommended, a resistant strain of fungi may be present. If the treatment is ineffective due to the presence of a metalaxyl resistant strain of fungi, neither Ridomil/Bravo 81W nor any other fungicide with similar action will effectively control that disease. Consideration should then be given to the prompt use of other types of suitable fungicides. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance in your particular crop and disease control situation. DO NOT USE RIDOMIL/BRAVO 81W FOR DISEASE CONTROL IN CROPS GROWN IN GREENHOUSES, LATH HOUSES, OR IN BEDDING PLANT STRUCTURES.

Mixing Instructions

Ridomil/Bravo 81W is an effective fungicide when used according to label directions for control of a broad spectrum of plant diseases. Thorough, uniform coverage is essential for disease control. Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate thoroughly before and during application. Flush spray tank thoroughly with water daily after use and dispose of pesticide rinsate by application to an already treated area.

Amount of water needed per acre will vary with crop and amount of plant growth. Apply in a minimum of 20 gals./A for dilute sprays and 5 gals./A for concentrate ground sprays and aerial applications.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

Precaution: Water-soluble prepacks of Ridomil/Bravo 81W must be completely dissolved and dispersed in water before any other tank mix partner, including micronutrients or other liquid or dry fertilizers, are added to the spray solution.

Ridomil/Bravo 81W Alone: Add 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble bags of Ridomil/Bravo 81W into the tank all at once. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the pouches have dissolved, and the material has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Ridomil/Bravo 81W + Tank Mixtures: Add 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble bags of Ridomil/Bravo 81W into the tank all at once. Continue agitation while adding the remainder of the water. Allow the bags of Ridomil/Bravo 81W to dissolve and the product to completely disperse into the mix water. Then add the desired amount of products recommended for tank mixture and allow them to become completely dispersed. Continue agitation to maintain a uniform suspension until all of the spray solution has been applied.

Note: Do not let spray mixture stand overnight in the spray tank. Rinse spray tank at the end of the day.

Preharvest Intervals

Unless otherwise stated in the directions for use for a specific crop, there is no preharvest interval.

Application Through Irrigation Systems

Ridomil/Bravo 81W alone or in combination with other fungicides which are registered for sprinkler irrigation may be applied through irrigation systems.

Apply this product only through center pivot, solid set, hand move or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Posting Requirements

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public such as golf courses or retail greenhouses.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

Operating Instructions:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional

pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended.

Application Instructions

Ridomil/Bravo 81W must be applied on the schedule specified in the specific crop use recommendations, not according to the irrigation schedule. If irrigation schedules are used, ground or aerial applications must supplement chemigation applications to achieve adequate disease control.

With the exception of potatoes, Ridomil/Bravo 81W has not been sufficiently tested when applied through irrigation systems to assure consistent product performance for all labeled uses. The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler irrigation equipment. Users should check with state and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

Center-Pivot Irrigation Equipment (Use only with drive systems which provide uniform water distribution):

1. Determine the size of the area to be treated.
2. Determine the time required to apply 1/4 inch water over the area to be treated when the system and injection equipment is operated at normal pressures recommended by the equipment manufacturer. Run the system at 80-95% of the manufacturer's rated capacity.
3. Using water, determine the injection pump output when operated at normal line pressure.
4. Determine the amount of Ridomil/Bravo 81W required to treat the area covered by the irrigation system.
5. Add the required number of Ridomil/Bravo 81W soluble bags and sufficient water to meet the injection time requirements to the solution tank. (See Mixing Instructions section of this label.)

6. Maintain constant solution tank agitation during the injection period.
7. Stop injection equipment after treatment is completed. Continue to operate system until the Ridomil/Bravo 81W solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

1. Determine the acreage covered by the sprinklers.
2. Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30 minute interval.
3. Determine the amount of Ridomil/Bravo 81W required to treat the area covered by the irrigation system.
4. Add the required number of Ridomil/Bravo 81W soluble bags into the same quantity of water used to calibrate the injection period. (See Mixing Instructions section of this label.)
5. Operate system at the same pressure and time interval established during the calibration.
6. Inject Ridomil/Bravo 81W at the end of the irrigation cycle or as a separate application to maximize retention of the fungicide by the foliage.
7. Stop injection equipment after treatment is completed. Continue to operate the system until the Ridomil/Bravo 81W solution has cleared the last sprinkler head.

Cole Crops (Broccoli, Cabbage, and Cauliflower)*

*Broccoli, Chinese broccoli (gai lan, white flowering broccoli), cabbage, Chinese cabbage (tight-heading varieties only) and cauliflower only.

Ridomil/Bravo 81W, when used as a foliar application in a preventive disease control program, provides effective control of downy mildew and Alternaria leaf spot.

Downy Mildew: Apply 1 1/2 - 2 lbs./A. Begin applications when conditions are favorable for disease, but before infection and continue at 14-day intervals until the threat of disease is over. Use the higher rate under severe disease pressure.

Alternaria Leaf Spot: Apply 1 1/2-2 lbs./A. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until threat of disease is over. Under severe disease pressure, use other effective EPA-registered fungicides according to label directions between the Ridomil/Bravo 81W sprays.

Notes: 1) To avoid possible illegal residues, do not apply within 7 days of harvest. 2) To minimize the potential for resistance, do not make more than 4 applications of Ridomil/Bravo 81W per crop.

Cucurbit Vegetables*

*Cucumber, cantaloupe, casaba, crenshaw, honeydew melon, honey balls, muskmelon, Persian melon, watermelon, pumpkin, summer squash, winter squash and cucurbit hybrids only.

Ridomil/Bravo 81W, when used as a foliar spray in a preventive disease control program, provides effective control of many diseases of cucurbit vegetables.

Downy Mildew: Apply 1 1/2 - 2 lbs./A. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until the threat of disease is over. Use the higher rate under severe disease pressure.

Anthracnose, Cercospora Leaf Spot, Gummy Stem Blight, Leaf Blight and Scab: Apply 2-3 lbs./A. Begin applications when conditions are favorable for disease, but before infection, and continue at 10-14-day intervals. Under severe disease pressure, alternate Ridomil/Bravo 81W sprays with other effective EPA-registered fungicides every 7 days.

Note: To minimize the potential for resistance, do not make more than 4 applications of Ridomil/Bravo 81W per crop.

Onions (Dry Bulb, Seed and Green Onions)

Ridomil/Bravo 81W, when used as a foliar spray in a preventive disease control program, provides effective control of downy mildew, Botrytis leaf blight and purple blotch.

Downy Mildew: Apply 1 1/2 - 2 lbs./A. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until the threat of disease is over. Use a compatible spreader-sticker at rates recommended on the product label. Use the higher rate under severe disease pressure.

Botrytis Leaf Blight (Blast) and Purple Blotch: Apply 2 lbs./A on a 14-day schedule and apply other effective EPA-registered fungicides between the Ridomil/Bravo 81W sprays; or apply 2 lbs./A of Ridomil/Bravo 81W on a 7-day schedule.

Notes: 1) To avoid possible illegal residues, do not apply to dry bulb onions within 7 days of harvest or to green onions within 21 days of harvest. 2) Do not make more than 3 applications to green onions per season. If more than 14 days remain after 3 applications of Ridomil/Bravo 81W, it is recommended

that another EPA- registered fungicide which does not contain chlorothalonil be used. 3) To minimize the potential for resistance, do not make more than 4 applications of Ridomil/Bravo 81W to dry bulb or seed onions.

Potatoes

Ridomil/Bravo 81W, when used as a foliar spray in a preventive disease control program, provides effective control of late blight, early blight and Botrytis vine rot. Two to three applications will also provide control of tuber rot in the field caused by Phytophthora spp. and storage rot caused by Pythium and Phytophthora spp.

Late Blight: Apply 1 1/2 - 2 lbs./A. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until threat of disease is over. Use the higher rate under severe disease pressure.

Tuber Rot and Storage Rot (Phytophthora and Pythium spp.): Apply 2 lbs./A. Make one application at flowering and another 14 days later. If the field has a history of tuber rot or if foliar late blight is present, make a third application 14 days after the second.

Early Blight and Botrytis Vine Rot: Apply 1 1/2 - 2 lbs./A. Begin applications when plants are six inches high or when conditions are favorable for disease, but before infection, and continue at 14-day intervals until the threat of disease is over. Under severe disease pressure, use 1 1/2 lbs./A on a 7-day schedule or use other effective EPA-registered fungicides between the Ridomil/Bravo 81W sprays.

Notes: 1) To avoid possible illegal residues, do not apply within 7 days of harvest. 2) To minimize the potential for resistance, do not make more than 4 applications of Ridomil/Bravo 81W per crop.

Tomatoes

Ridomil/Bravo 81W, when used as a foliar spray in a preventive disease control program, provides effective control of several tomato diseases.

Late Blight and Fruit Rot Caused by Phytophthora spp: Apply 1 1/2 - 2 lbs./A. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals until threat of disease is over. Use the higher rate under severe disease pressure.

Early Blight, Gray Leaf Spot, Gray Leaf Mold, and Septoria Leaf Spot: Apply 2 lbs./A. Begin applications when conditions are favorable for disease, but before infection, and continue at 14-day intervals. Under severe disease pressure, use other effective EPA-registered fungicides between the Ridomil/Bravo 81W sprays.

Anthracnose, Alternaria Fruit Rot (Black Mold), Rhizoctonia Fruit Rot and Botrytis Gray Mold: Apply 2-3 lbs./A. Begin applications when conditions are favorable for disease, but before infection, and continue at 10-14 day intervals. Under severe disease pressure, use other effective EPA-registered fungicides between the Ridomil/Bravo 81W sprays.

Note: To minimize the potential for resistance, 1) Do not make more than 4 applications of Ridomil/Bravo 81W per crop. 2) To avoid possible illegal residues, do not apply within 7 days of harvest.

Rotation (Plant Back) Restrictions

Do not plant any crop which is not registered for use with metalaxyl in soil treated with Ridomil/Bravo 81W for a period of 12 months.

Rotation (Plant Back) Restrictions

<u>Rotation Crop</u>	<u>Planting Time From Last Application of Ridomil/Bravo 81W</u>
Alfalfa (inc. birdsfoot trefoil)	-0- days
Almonds	-0- days
Apples*	-0- days
Asparagus	-0- days
Avocadoes	-0- days
Blueberries	-0- days
Broccoli	-0- days
Citrus	-0- days
Corn	12 months
Cabbage	-0- days
Celery	12 months
Cauliflower	-0- days
Carrots	-0- days
Cranberry*	-0- days
Brussels Sprouts	12 months
Eggplant	-0- days
Mint	12 months
Grass Grown for Seed	-0- days
Shallots	12 months
Garlic	12 months
Papaya	-0- days
Peppers	-0- days
Potatoes	-0- days
Cotton	-0- days
Cucurbit Vegetables	-0- days
Deciduous Fruits and Nuts*	-0- days
Hops	-0- days
Head Lettuce	-0- days
Legume Vegetables	-0- days
Onions (Bulb, Green and Seed)	-0- days
Peanuts	-0- days
Pineapples	-0- days
Raspberries	-0- days
Soybeans	-0- days
Spinach	-0- days
Stone Fruits	-0- days
Strawberries	-0- days
Sugar Beets	-0- days
Tobacco	-0- days
Tomatoes	-0- days
Walnuts	-0- days
Wheat/Barley/Oats	14 days
Root and Tuber Vegetables	-0- days
Crops <u>Not For</u> Food or Feed	-0- days
All Other Crops <u>For</u> Food or Feed	12 months

*These crops may be planted immediately following last application of Ridomil/Bravo 81W, provided they will not bear harvestable fruit within 12 months.

Storage and Disposal

Storage - Store in a cool, dry place. Do not store this product under wet conditions. Handle outer container carefully when stored at temperatures less than 50°F to avoid breakage of inner soluble bag.

Pesticide Disposal

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this pesticide are acutely hazardous. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures.

For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container

Do not reuse outer container. Dispose of empty outer container in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. Stay out of smoke from burning bags.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372 day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER/PELIGRO

Corrosive. Causes irreversible eye damage. May be fatal if swallowed or inhaled. Harmful if absorbed through skin. Do

not get in eyes, on skin, or on clothing. Wear goggles or face shield when handling this product. Do not breathe spray mist. Wear a mask or a jointly approved pesticide respirator (by the Mining Enforcement and Mining Administration and the National Institute for Occupational Safety and Health) when handling this product. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco. Remove contaminated clothing and wash before reuse.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor drift from treated area. Do not contaminate water when disposing of equipment wash waters.

Ridomil® trademark of CIBA-GEIGY Corporation for metalaxyl
U.S. Patent 4,151,299

Bravo® trademark of ISK Biotech for chlorothalonil

©1992 CIBA-GEIGY Corporation

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419

Revised June 15, 1980
Revised October 1987 -
Chemigation
Revised February 8, 1988 -
Administrative Amendments
Revised October 12, 1988
water soluble bag
Revised September 18, 1989
Metalaxyl FRSTR labeling.
Revised December 13, 1990
New graphics, new est. #
Revised June, 1992
Bravo reduced plantbacks,
minor adm. changes

CGA (DRAFT)

[GANNONC.LABELR.]RIDOMIL-BRAVO-DRFT - 6/22/92