

Page 1

10-6-88

1319

APR 8 1988

206551  
|  
17

Ms. Karen Stumpf  
Ciba-Geigy Corporation  
Agricultural Division  
P.O. Box 18300  
Greensboro, N.C. 27419

Dear MS. Stumpf:

Subject: Ridomil PC 11G Granular Fungicide  
EPA Registration Number 100-664  
Your Submission Dated April 7, 1988  
Addition of beans to the label

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
  - a. revise the section under "Environmental Hazards" to read:  
"This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water or wetlands. Cover or incorporate all granules. Do not contaminate water when disposing of equipment washwaters."
3. Submit one (1) copy of your final printed labeling before you release the product for shipment.

PM 21

-2-

100-264

3419

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

SL

Susan Lewis  
Acting Product Manager (21)  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Ridomil® PC 11G

Granular  
Fungicide

For the control of diseases of  
beans, cotton, and peanuts  
caused by Pythium and Rhizoctonia

50 Pounds  
Net Weight

Active Ingredients:

|  |               |
|--|---------------|
| PCNB: Pentachloronitrobenzene .....  | 10.0%         |
| Metalaxyl: <u>N</u> -(2,6-dimethylphenyl)- <u>N</u> -<br>(methoxyacetyl)alanine methyl ester ..... | 1.0%          |
| <u>Inert Ingredients:</u>  | <u>89.0%</u>  |
| <u>Total:</u>  | <u>100.0%</u> |

Ridomil PC 11G is a granular fungicide

Keep Out of Reach of Children

Caution

See additional precautionary  
statements on back of bag.

See directions for use on back of bag.

ACCEPTED  
with comments  
to EPA

APR - 6 1989

100-664

EPA Reg. No. 100-664

Control No. 54810

©1987 CIBA-GEIGY Corporation

CIBA-GEIGY

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR CROP INJURY.

General Information

Ridomil PC 11G is a granular fungicide containing 1% metalaxyl, a systemic fungicide, and 10% PCNB (pentachloronitrobenzene). It provides control of damping-off and seed and seedling rot diseases caused by Pythium and Rhizoctonia in beans, cotton, and peanuts. Ridomil PC 11G will also provide suppression of pod rot in peanuts caused by Sclerotium rolfsii.

Note: Metalaxyl is a systemic fungicide having a specific mode of action. Metalaxyl could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, CIBA-GEIGY cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance or ways to control any possible metalaxyl resistant strains of fungi which may occur.

Applicator Settings

It is important that the user calibrate each individual granular applicator before its use. The settings in the charts under the Cotton and Peanuts sections below SHOULD ONLY BE USED AS A STARTING POINT. Even after calibration, it is best to check the actual application rate under your operating conditions. Application rates can vary due to changing moisture conditions, equipment wear, normal variation in product bulk density and particle size distribution.

-4-

Calibration: To calibrate a granular applicator, follow the procedure as outlined below:

1. Fill the hopper with Ridomil PC 11G.
2. Attach a plastic bag to the spout opening.
3. Refer to the appropriate chart in either the Cotton or Peanuts section of this label and adjust the setting on the applicator to match the desired rate per acre.
4. Measure off 1,000 linear feet and drive at the speed selected.
5. Weigh or measure the volume of each bag, change settings as needed to obtain the desired rate and repeat steps 2-5 until the setting delivers the correct number of ounces per 1,000 ft.
6. Repeat steps 1-5 for each applicator box.

Conversions:

1. Time to cover 1,000 feet at:  

|                          |
|--------------------------|
| 2 mph = 5 mins. 40 secs. |
| 3 mph = 3 mins. 54 secs. |
| 4 mph = 2 mins. 50 secs. |
| 5 mph = 2 mins. 16 secs. |
| 6 mph = 1 min. 57 secs.  |
2. Ounces (wt.) of granules to approximate ounces (volume):  
6 oz. wt. = 8 fl. oz. (1 level cup)

When calibrating granular applicators, follow the recommendations below:

1. To reduce granule flow variability, adjust orifice openings by moving the controls in one direction only, such as from the closed position to open.
2. Granule flow through an orifice is not always proportional to ground speed, so make calibration checks at the same speed at which you plant and maintain that ground speed as constantly as possible when planting.

3. Granular pesticide products vary considerably in size, density, and particle shape, so recalibrate applicators whenever you change brands and/or formulations. The same brand can even change from year to year.
4. Changing humidity and temperature during the day can change the flow rate of granules through the applicator's orifice. Recalibrate when conditions change and periodically readjust orifice settings to compensate.
5. Know the recommended rate. For band applications, the area actually is the area covered by the band, not total cropland area planted. Some row-crop recommendations are based on treating in-the-row and these rates generally are specified as amounts (ounces) of granules per certain row length (often 1,000 feet).
6. Calibrate frequently. There are several acceptable calibration methods. Learn one. Do the arithmetic first, then complete the material collection phase under field conditions.

Calculate the amount of Ridomil PC 11G needed for band treatment as follows:

$$\frac{\text{Band Width in Inches}}{\text{Row Spacing in Inches}} \times \text{Broadcast Rate per Acre} = \text{Band Rate per Acre}$$

Beans

Ridomil PC 11G will control damping-off and seed and seedling rots of dry and green beans (bush and pole types) caused by *Pythium* and *Rhizoctonia* when used at planting.

Apply 3/4 lb. (12 oz.) of Ridomil PC 11G per 1,000 linear feet of row at the time of planting. Adjust the application equipment so the granules are mixed with the soil surrounding the seed.

Note: Do not allow the feeding of vines or grazing of foliage by livestock, or illegal residues may result.

Cotton

Ridomil PC 11G will control damping-off, seed and seedling rot diseases caused by *Pythium* spp. and *Rhizoctonia solani*.

Chart 1: Cotton - Suggested Applicator Settings

|   | ←----- 4 MPH -----→ |      |             |      | ←----- 5 MPH -----→ |      |      |             | ←----- 6 MPH -----→ |      |      |             |
|---|---------------------|------|-------------|------|---------------------|------|------|-------------|---------------------|------|------|-------------|
| 40" Rows<br>In-Furrow<br>Lbs./A                                 | 10                  | 9    | 8           | 7    | 10                  | 9    | 8    | 7           | 10                  | 9    | 8    | 7           |
| Eqv. Ozs./<br>1,000 ft.   | 12.3                | 11.1 | 9.8         | 8.6  | 12.3                | 11.1 | 9.8  | 8.6         | 12.3                | 11.1 | 9.8  | 8.6         |
| Gandy   | 28                  | 26.8 | 25          | 23.5 | 31                  | 30   | 28   | 26.3        | 34.5                | 32.5 | 30.5 | 29          |
| Allis<br>Chalmers   | 2.4                 | 2.1  | 1.9         | 1.6  | 3                   | 2.8  | 2.4  | 2.05        | 3.6                 | 3.2  | 2.9  | 2.5         |
| Noble<br>1972 and Before  | 19                  | 17   | 15          | 13   | 23                  | 21   | 19   | 17          | 27                  | 25   | 22   | 20          |
| Noble<br>1973 and Later   | 14                  | 13   | 12 or<br>11 | 10   | 17 or<br>16         | 15   | 14   | 13 or<br>12 | 19                  | 18   | 16   | 15 or<br>14 |
| John Deere<br>Metal Box   | 2/12                | 2/10 | 2/8         | 2/5  | 2/18                | 2/15 | 2/12 | 2/9         | 2/22                | 2/19 | 2/16 | 2/13        |
| John Deere<br>1978 and Before<br>Meter Scale<br>0--10--20--etc. | 28                  | 27   | 24          | 22   | 31                  | 30   | 28   | 26          | 33                  | 32   | 31   | 29          |
| John Deere<br>Post 1978<br>Meter Scale<br>0--5--15--etc.        | 18                  | 17   | 16          | 15   | 21                  | 20   | 18   | 17          | 23                  | 22   | 21   | 19          |
| I.H.<br>1981 and Later  | 3.1                 | 2.9  | 2.6         | 2.35 | 3.6                 | 3.4  | 3.1  | 2.8         | 4.5                 | 4.0  | 3.55 | 3.2         |

**BEST AVAILABLE COPY**

100-659

100-659

100-659



-7-

For best results, use high quality seed and plant into a properly prepared seedbed.

Apply 7-10 lbs. of Ridomil PC 11G per 13,000 linear feet of row (8.6-12.3 oz./1,000 linear feet) at the time of planting. Adjust the application equipment so the granules are mixed with the soil surrounding the seed. See Chart 1 below for suggested applicator settings and the General Information section of this label for calibration instructions.

Note: Do not allow the feeding or grazing of cotton foliage by livestock, or illegal residues may result.

### Peanuts

Ridomil PC 11G applied at planting will provide control of seed and seedling rots of peanuts caused by Pythium and Rhizoctonia. Applications of Ridomil PC 11G during pegging will provide control of pod rot caused by Pythium and Rhizoctonia and will provide suppression of Sclerotium rolfsii.

Seed and Seedling Rots: Apply 12.5-25 lbs. of Ridomil PC 11G/A in a 9-inch band at the time of planting. Incorporate with mechanical equipment to a depth of 2-4 inches. For row spacings other than 36 inches and band widths other than 9 inches, calculate the amount of Ridomil PC 11G needed per acre based on the formula in the General Information section of this label, using a broadcast rate of 50-100 lbs./A. Use the higher rate in fields with a history of seedling disease problems or when weather conditions are expected to be unfavorable for rapid germination and growth of the seedlings.

Pod Rot: Apply 50-100 lbs./A (21.5-43 oz./1,000 linear feet) (broadcast basis) at early pegging. For banded applications, adjust the equipment so the granules are distributed over the row. The higher rate should be used in areas where pod rot is severe. See Chart 2 below for suggested applicator settings and the General Information section of this label for calibration instructions.

Sclerotium rolfsii: When Ridomil PC 11G is used for pod rot control, some suppression of Sclerotium rolfsii will be obtained. In areas where Sclerotium rolfsii is a problem use other fungicides that are labeled for this disease.

Notes: (1) Do not allow the feeding or grazing of peanut foliage or harvest residues by livestock, or illegal residues may result. (2) Do not apply more than 100 lbs. of Ridomil PC 11G/A/season.

**BEST AVAILABLE COPY**

CHART 2: Peanuts - Suggested Applicator Settings for Pod Rot Control

| Rate for 14" Band Width<br>Oz./1,000 linear ft.*<br>(one opening per row) |           | 43   | 38.6 | 34.3 | 30   | 25.7 | 21.5 |
|---|-----------|------|------|------|------|------|------|
| Lbs./A for 40" rows   |           | 35   | 31.5 | 28   | 24.5 | 21.0 | 17.5 |
| Applicator Brand  | Speed mph |      |      |      |      |      |      |
| Gandy   | 2         | 36   | 34   | 32   | 30   | 28   | 26   |
|   | 3         | 42.5 | 40.5 | 39   | 36.5 | 33.5 | 31   |
|   | 4         | 47.5 | 45.5 | 43   | 41.5 | 39   | 36   |
|   | 5         | 53.5 | 50   | 47.5 | 45   | 42.5 | 40   |
|   | 6         | 61   | 56   | 52   | 49   | 45   | 42.5 |
| John Deere<br>1978 & Before<br>Meter Scale<br>0..10..20..<br>Etc.         | 2         | 33   | 32   | 30   | 28   | 26   | 24   |
|   | 3         | 39   | 37   | 36   | 34   | 32   | 30   |
|   | 4         | Max  | 43   | 40   | 38   | 36   | 33   |
|   | 5         |      |      | Max  | 43   | 39   | 36   |
|   | 6         |      |      |      | Max  | 43   | 39   |
| John Deere<br>1979 & Later<br>Meter Scale<br>0..5..15..<br>Etc.           | 2         | 24   | 22   | 21   | 20   | 18   | 16   |
|   | 3         | 31   | 28   | 26   | 25   | 22   | 20   |
|   | 4         | 37   | 34   | 32   | 30   | 27   | 24   |
|   | 5         | 43   | 40   | 37   | 34   | 31   | 28   |
|   | 6         |      |      | 41   | 38   | 34   | 31   |
| Noble<br>1973 & Later   | 2         | 20   | 18   | 17   | 15   | 12   | 10   |
|   | 3         | 28   | 26   | 24   | 22   | 18   | 16   |
|   | 4         | 34   | 32   | 30   | 27   | 24   | 20   |
|   | 5         | 38   | 36   | 34   | 31   | 28   | 24   |
|   | 6         | 40   | 39   | 37   | 35   | 32   | 28   |
| Noble<br>1972 & Before  | 2         | 28   | 26   | 23   | 20   | 17   | 13   |
|   | 3         | 37   | 34   | 32   | 29   | 26   | 22   |
|   | 4         | 41   | 40   | 38   | 35   | 32   | 28   |
|   | 5         | 45   | 43   | 41   | 39   | 37   | 32   |
|   | 6         | 50   | 46   | 44   | 42   | 40   | 37   |

Continued ... 9

**BEST AVAILABLE COPY**

|  |   |      |      |      |      |      |      |
|--|---|------|------|------|------|------|------|
| John Deere<br>Metal Box                    | 2 | 2/25 | 2/21 | 2/19 | 2/16 | 2/11 | 2/06 |
|  | 3 | 3/03 | 2/32 | 2/30 | 2/26 | 2/21 | 2/16 |
|  | 4 | 3/12 | 3/08 | 3/04 | 3/00 | 2/29 | 2/24 |
|  | 5 | 3/21 | 3/15 | 3/12 | 3/08 | 3/02 | 2/30 |
|  | 6 | 3/33 | 3/23 | 3/20 | 3/14 | 3/08 | 3/02 |
| International<br>Harvester<br>1981 & Later | 2 | 5/3  | 4/7  | 4/2  | 3/6  | 3/1  | 2/6  |
|  | 3 | 7/8  | 7/0  | 6/2  | 5/4  | 4/7  | 3/9  |
|  | 4 | 10/3 | 9/3  | 8/3  | 7/3  | 6/2  | 5/2  |
|  | 5 |      | 11/6 | 10/3 | 9/0  | 7/7  | 6/4  |
|  | 6 |      |      |      | 10/8 | 9/3  | 7/8  |
| Allis<br>Chalmers                          | 2 | 5/1  | 4/6  | 4/2  | 3/8  | 3/3  | 2/9  |
|  | 3 | 7/3  | 6/6  | 6/0  | 5/3  | 4/6  | 4/0  |
|  | 4 | 9/6  | 8/6  | 7/7  | 6/8  | 5/9  | 5/0  |
|  | 5 | 11/6 | 10/5 | 9/4  | 8/3  | 7/3  | 6/1  |
|  | 6 |      |      | 11/2 | 9/9  | 8/5  | 7/2  |

\*If an applicator has a dual metering unit (2 openings/row) set guage at half indicated rate.

Note: To calculate the amount of Ridomil PC 11G needed for row spacings other than 40 inches and band widths other than 14 inches, refer to the formula in the General Information section.

Rotation Crops\*

\*These crops represent those which may be planted (rotated to) according to the interval listed following an application of Ridomil PC 11G to beans, cotton, or peanuts.

Rotational Interval

|   |           |
|---|-----------|
| Asparagus                                 | 0 days    |
| Beans, Dry and Snap                       | 0 days    |
| Broccoli                                  | 0 days    |
| Brussels Sprouts                          | 18 months |
| Cabbage                                   | 0 days    |
| Cauliflower                               | 0 days    |
| Corn                                      | 9 months  |
| Cotton                                    | 0 days    |
| Cucurbit Vegetables                       | 0 days    |
| Garlic                                    | 12 months |
| Head Lettuce                              | 0 days    |
| Hops                                      | 0 days    |
| Legume Vegetables Other Than Beans        | 0 days    |
| Onions                                    | 0 days    |
| Other Root Crops                          | 12 months |
| All Other Crops Intended for Food or Feed | 18 months |

|          |         |
|----------|---------|
| Peanuts  | 0 days  |
| Peppers  | 0 days  |
| Potatoes | 0 days  |
| Soybeans | 0 days  |
| Spinach  | 0 days  |
| Tomatoes | 0 days  |
| Wheat    | 14 days |

Storage and Disposal

Pesticide

Store in a dry place. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide is a violation of federal law. Pesticide that cannot be used according to label instructions must be disposed of according to federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Disposal

Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call (919) 292-7100 day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through the skin or inhaled.  
Causes moderate eye irritation. Avoid contact with

Continued ... 11

100 664 / High Quality Copy

# Ridomil PC<sup>®</sup>

## 11G

**NOT REVIEWED**  
In Accordance with PR Notice 82-2.  
Based on Draft Labeling Dated

APRIL 6, 1989

## Granular Fungicide

For the control of diseases of beans, cotton, and peanuts caused by *Pythium* and *Rhizoctonia*

**50 Pounds**  
Net Weight

Active Ingredients:  
PCNB: Pentachloro-  
nitrobenzene . . . . . 10.0%  
Metalaxyl: *N*-(2,6-  
dimethylphenyl)-*N*-  
methoxyacetyl) alanine  
methyl ester . . . . . 1.0%

Inert Ingredients: 89.0%

Total: 100.0%

Control No. **54810**

Ridomil PC 11G is a granular fungicide.

Keep Out of Reach of Children.

### Caution

See additional precautionary statements on back of bag.

EPA Reg. No. 100-664

EPA Est. 8590-PA-1

See directions for use on back of bag.

# CIBA-GEIGY

## Ridomil PC 11G

### DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

**IMPORTANT:** Read the entire **Directions for Use** and the **Conditions of Sale and Warranty** before using this product

#### Conditions of Sale and Warranty

The **Directions for Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use** subject to the inherent risks referred to above. **CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.** CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing **Conditions of Sale and Warranty**, which may be varied only by agreement in writing and signed by a duly authorized representative of CIBA-GEIGY.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR CROP INJURY.**

#### General Information

Ridomil PC 11G is a granular fungicide containing 1% metalaxyl, a systemic fungicide, and 10% PCNB (pentachloronitrobenzene). It provides control of damping-off and seed and seedling rot diseases caused by *Pythium* and *Rhizoctonia* in beans, cotton, and peanuts. Ridomil PC 11G will also provide suppression of pod rot in peanuts caused by *Sclerotium rolfsii*.

**Note:** Metalaxyl is a systemic fungicide having a specific mode of action. Metalaxyl could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, CIBA-GEIGY cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance or ways to control any possible metalaxyl resistant strains of fungi which may occur.

#### Applicator Settings

It is important that the user calibrate each individual granular applicator before its use. The settings in the charts under the **Cotton** and **Peanuts** sections below SHOULD ONLY BE USED AS A STARTING POINT. Even after calibration, it is best to check the actual application rate under your operating conditions. Application rates can vary due to changing moisture conditions, equipment wear, normal variation in product bulk density and particle size distribution.

**Calibration:** To calibrate a granular applicator, follow the procedure as outlined below.

- 1 Fill the hopper with Ridomil PC 11G
- 2 Attach a plastic bag to the spout opening
- 3 Refer to the appropriate chart in either the **Cotton** or **Peanuts** section of this label and adjust the setting on the applicator to match the desired rate per acre
- 4 Measure off 1 000 linear feet and drive at the speed selected
- 5 Weigh or measure the volume of each bag, change settings as needed to obtain the desired rate and repeat steps 2-5 until the setting delivers the correct number of ounces per 1 000 ft
- 6 Repeat steps 1-5 for each applicator box

## Ridomil PC 11G

### Conversions

1. Time to cover 1,000 feet at:
  - 2 mph = 5 mins. 40 secs.
  - 3 mph = 3 mins. 54 secs.
  - 4 mph = 2 mins. 50 secs.
  - 5 mph = 2 mins. 16 secs.
  - 6 mph = 1 min. 57 secs.
2. Ounces (wt.) of granules to approximate ounces (volume):
  - 6 oz. wt. = 8 fl. oz. (1 level cup)

When calibrating granular applicators, follow the recommendations below:

1. To reduce granule flow variability, adjust orifice openings by moving the controls in one direction only, such as from the closed position to open.
2. Granule flow through an orifice is not always proportional to ground speed, so make calibration checks at the same speed at which you plant and maintain that ground speed as constantly as possible when planting.
3. Granular pesticide products vary considerably in size, density, and particle shape, so recalibrate applicators whenever you change brands and/or formulations. The same brand can even change from year to year.
4. Changing humidity and temperature during the day can change the flow rate of granules through the applicator's orifice. Recalibrate when conditions change and periodically readjust orifice settings to compensate.
5. Know the recommended rate. For band applications, the area actually is the area covered by the band, not total cropland area planted. Some row-crop recommendations are based on treating in-the-row and these rates generally are specified as amounts (ounces) or granules per certain row length (often 1,000 feet).
6. Calibrate frequently. There are several acceptable calibration methods. Learn one. Do the arithmetic first, then complete the material collection phase under field conditions.

Calculate the amount of Ridomil PC 11G needed for band treatment as follows:

$$\frac{\text{Band Width in Inches}}{\text{Row Spacing in Inches}} \times \frac{\text{Broadcast Rate}}{\text{Per Acre}} = \frac{\text{Band Rate}}{\text{Per Acre}}$$

### Beans

Ridomil PC 11G will control damping-off and seed and seedling rots of dry and green beans (bush and pole types) caused by *Pythium* and *Rhizoctonia* when used at planting.

Apply 3/4 lb. (12 oz.) of Ridomil PC 11G per 1,000 linear feet of row at the time of planting. Adjust the application equipment so the granules are mixed with the soil surrounding the seed.

**Note:** Do not allow the feeding of vines or grazing of foliage by livestock, or illegal residues may result.

### Cotton

Ridomil PC 11G will control damping-off, seed and seedling rot diseases caused by *Pythium* spp. and *Rhizoctonia solani*. For best results, use high quality seed and plant into a properly prepared seedbed.

Apply 7-10 lbs. of Ridomil PC 11G per 13,000 linear feet of row (8.6-12.3 oz /1,000 linear feet) at the time of planting. Adjust the application equipment so the granules are mixed with the soil surrounding the seed. See Chart 1 below for suggested applicator settings and the **General Information** section of this label for calibration instructions.

**Note:** Do not allow the feeding or grazing of cotton foliage by livestock, or illegal residues may result.

part

100-664

16319

Ridomil PC 11G

Chart 1: Cotton — Suggested Applicator Settings

| Applicator Brand                                       | Speed MPH | 40" Rows In-Furrow Lbs./A |      |          |          |
|--|-----------|---------------------------|------|----------|----------|
|  |           | 10                        | 9    | 8        | 7        |
|  |           | Equivalent Oz./1,000 ft.  |      |          |          |
|  |           | 12.3                      | 11.1 | 9.8      | 8.6      |
| Gandy  | 4         | 28                        | 26.8 | 25       | 23.5     |
|  | 5         | 31                        | 30   | 28       | 26.3     |
|  | 6         | 34.5                      | 32.5 | 30.5     | 29       |
| Allis Chalmers   | 4         | 2.4                       | 2.1  | 1.9      | 1.6      |
|  | 5         | 3                         | 2.8  | 2.4      | 2.05     |
|  | 6         | 3.6                       | 3.2  | 2.9      | 2.5      |
| Noble 1972 and Before                                  | 4         | 19                        | 17   | 15       | 13       |
|  | 5         | 23                        | 21   | 19       | 17       |
|  | 6         | 27                        | 25   | 22       | 20       |
| Noble 1973 and Later                                   | 4         | 14                        | 13   | 12 or 11 | 10       |
|  | 5         | 17 or 15                  | 15   | 14       | 13 or 12 |
|  | 6         | 19                        | 18   | 16       | 15 or 14 |
| John Deere Metal Box                                   | 4         | 2/12                      | 2/10 | 2/8      | 2/5      |
|  | 5         | 2/18                      | 2/15 | 2/12     | 2/9      |
|  | 6         | 2/22                      | 2/19 | 2/16     | 2/13     |
| John Deere 1978 and Before Meter Scale 0..10..20..etc. | 4         | 28                        | 27   | 24       | 22       |
|  | 5         | 31                        | 30   | 28       | 26       |
|  | 6         | 33                        | 32   | 31       | 29       |
| John Deere 1979 and Later Meter Scale 0.5..15..etc.    | 4         | 18                        | 17   | 16       | 15       |
|  | 5         | 21                        | 20   | 18       | 17       |
|  | 6         | 23                        | 22   | 21       | 19       |
| International Harvester 1981 and Later                 | 4         | 3.1                       | 2.9  | 2.6      | 2.35     |
|  | 5         | 3.6                       | 3.4  | 3.1      | 2.8      |
|  | 6         | 4.5                       | 4    | 3.55     | 3.2      |

**Peanuts**

Ridomil PC 11G applied at planting will provide control of seed and seedling rots of peanuts caused by *Pythium* and *Rhizoctonia*. Applications of Ridomil PC 11G during pegging will provide control of pod rot caused by *Pythium* and *Rhizoctonia* and will provide suppression of *Sclerotium rolfsii*.

**Seed and Seedling Rots:** Apply 12.5-25 lbs. of Ridomil PC 11G/A in a 9-inch band at the time of planting. Incorporate with mechanical equipment to a depth of 2-4 inches. For row spacings other than 36 inches and band widths other than 9 inches, calculate the amount of Ridomil PC 11G needed per acre based on the formula in the **General Information** section of this label, using a broadcast rate of 50-100 lbs./A. Use the higher rate in fields with a history of seedling disease problems or when weather conditions are expected to be unfavorable for rapid germination and growth of seedlings.

**Pod Rot:** Apply 50-100 lbs /A (21.5-43 oz /1,000 linear feet) (broadcast basis) at early pegging. For banded applications, adjust the equipment so the granules are distributed over the row. The higher rate should be used in areas where pod rot is severe. See Chart 2 below for suggested applicator settings and the **General Information** section of this label for calibration instructions.

**Sclerotium rolfsii:** When Ridomil PC 11G is used for pod rot control, some suppression of *Sclerotium rolfsii* will be obtained. In areas where *Sclerotium rolfsii* is a problem use other fungicides that are labeled for this disease.



100-664  
17-19

**Ridomil PC™ 11G**

**Notes:** (1) Do not allow the feeding or grazing of peanut foliage or harvest residues by livestock, or illegal residues may result (2) Do not apply more than 100 lbs. of Ridomil PC 11G/A/season

**Chart 2: Peanuts — Suggested Applicator Settings for Pod Rot Control**

| Applicator Brand   | Speed MPH | Rate for 14" Band Width<br>Oz./1,000 linear ft. *<br>(one opening per row) |      |      |      |      |      |
|--|-----------|--|------|------|------|------|------|
|  |           | 43   | 38.6 | 34.3 | 30   | 25.7 | 21.5 |
|  |           | Lbs./A for 40" rows  |      |      |      |      |      |
|  |           | 35   | 31.5 | 28   | 24.5 | 21   | 17.5 |
| Gandy  | 2         | 36   | 34   | 32   | 30   | 28   | 26   |
|  | 3         | 42.5   | 40.5 | 39   | 36.5 | 33.5 | 31   |
|  | 4         | 47.5   | 45.5 | 43   | 41.5 | 39   | 36   |
|  | 5         | 53.5   | 50   | 47.5 | 45   | 42.5 | 40   |
|  | 6         | 61   | 56   | 52   | 49   | 45   | 42.5 |
| John Deere<br>1978 and Before<br>Meter Scale<br>0.10..20..etc. | 2         | 33   | 32   | 30   | 28   | 26   | 24   |
|  | 3         | 39   | 37   | 36   | 34   | 32   | 30   |
|  | 4         | Max  | 43   | 40   | 38   | 36   | 33   |
|  | 5         | -  | -    | Max  | 43   | 39   | 36   |
|  | 6         | -  | -    | -    | Max  | 43   | 39   |
| John Deere<br>1979 and Later<br>Meter Scale<br>0.5..15..etc.   | 2         | 24   | 22   | 21   | 20   | 18   | 16   |
|  | 3         | 31   | 28   | 26   | 25   | 22   | 20   |
|  | 4         | 37   | 34   | 32   | 30   | 27   | 24   |
|  | 5         | 43   | 40   | 37   | 34   | 31   | 28   |
|  | 6         | -  | -    | 41   | 38   | 34   | 31   |
| Noble<br>1973 and Later  | 2         | 20   | 18   | 17   | 15   | 12   | 10   |
|  | 3         | 28   | 26   | 24   | 22   | 18   | 16   |
|  | 4         | 34   | 32   | 30   | 27   | 24   | 20   |
|  | 5         | 38   | 36   | 34   | 31   | 28   | 24   |
|  | 6         | 40   | 39   | 37   | 35   | 32   | 28   |
| Noble<br>1972 and Before                                       | 2         | 28   | 26   | 23   | 20   | 17   | 13   |
|  | 3         | 37   | 34   | 32   | 29   | 26   | 22   |
|  | 4         | 41   | 40   | 38   | 35   | 32   | 28   |
|  | 5         | 45   | 43   | 41   | 39   | 37   | 32   |
|  | 6         | 50   | 46   | 44   | 42   | 40   | 37   |
| John Deere<br>Metal Box  | 2         | 2/25   | 2/21 | 2/19 | 2/16 | 2/11 | 2/06 |
|  | 3         | 3/03   | 2/32 | 2/30 | 2/26 | 2/21 | 2/16 |
|  | 4         | 3/12   | 3/08 | 3/04 | 3/00 | 2/29 | 2/24 |
|  | 5         | 3/21   | 3/15 | 3/12 | 3/08 | 3/02 | 2/30 |
|  | 6         | 3/33   | 3/23 | 3/20 | 3/14 | 3/08 | 3/02 |

Part 1

100 66-1

100 10

### Ridomil PC™ 11G

|   |   |      |      |      |      |     |     |
|---|---|------|------|------|------|-----|-----|
| International Harvester<br>1981 and Later | 2 | 5/3  | 4/7  | 4/2  | 3/6  | 3/1 | 2/6 |
|   | 3 | 7/8  | 7/0  | 6/2  | 5/4  | 4/7 | 3/9 |
|   | 4 | 10/3 | 9/3  | 8/3  | 7/3  | 6/2 | 5/2 |
|   | 5 | -    | 11/6 | 10/3 | 9/0  | 7/7 | 6/4 |
|   | 6 | -    | -    | -    | 10/8 | 9/3 | 7/8 |
| Allis Chalmers                            | 2 | 5/1  | 4/6  | 4/2  | 3/8  | 3/3 | 2/9 |
|   | 3 | 7/3  | 6/6  | 6/0  | 5/3  | 4/6 | 4/0 |
|   | 4 | 9/6  | 8/6  | 7/7  | 6/8  | 5/9 | 5/0 |
|   | 5 | 11/6 | 10/5 | 9/4  | 8/3  | 7/3 | 6/1 |
|   | 6 | -    | -    | 11/2 | 9/9  | 8/5 | 7/2 |

\* If an applicator has a dual metering unit (2 openings/row) set gauge at half indicated rate.

**Note:** To calculate the amount of Ridomil PC 11G needed for row spacings other than 40 inches and band widths other than 14 inches, refer to the formula in the **General Information** section.

#### Rotation Crops\*

| Crop  | Rotational Interval |
|---|---------------------|
| Asparagus   | 0 days              |
| Beans (Dry and Snap), Broccoli                    |                     |
| Cabbage, Cauliflower, Cotton, Cucurbit Vegetables |                     |
| Head Lettuce, Hops                                |                     |
| Legume Vegetables Other than Beans                |                     |
| Onions  |                     |
| Peanuts, Peppers, Potatoes                        |                     |
| Soybeans, Spinach                                 |                     |
| Tomatoes  |                     |
| Brussels Sprouts                                  |                     |
| Corn  |                     |
| Garlic  |                     |
| Other Root Crops                                  |                     |
| Wheat   |                     |
| All Other Crops Intended for Food or Feed         |                     |

\* These crops represent those which may be planted (rotated to) according to the interval listed following an application of Ridomil PC 11G to beans, cotton, or peanuts. Do not plant any crop not registered for use with metalaxyl and PCNB in soil treated with this product for a period of 12 months.

## Storage and Disposal

### Pesticide

Store in a dry place. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide is a violation of federal law. Pesticide that cannot be used according to label instructions must be disposed of according to federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

100-29

## Ridomil PC\* 11G

### Container Disposal

Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night.

## Precautionary Statements

### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Protective clothing including gloves, long-sleeved shirt, long pants, socks and shoes should be worn during mixing/loading procedures.

### Statement of Practical Treatment

**If in eyes:** Flush eyes with plenty of water. Call a physician if irritation persists.

**If on skin:** Wash with plenty of soap and water. Get medical attention.

**If inhaled:** Move to fresh air.

### Environmental Hazards

This product is toxic to fish and aquatic organisms. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Cover or incorporate all granules. Do not contaminate water when disposing of equipment wash waters.

Ridomil\* trademark of CIBA-GEIGY for metalaxyl

Ridomil PC\* trademark of CIBA-GEIGY

©1989 CIBA-GEIGY Corporation

Agricultural Division

CIBA-GEIGY Corporation

Greensboro, North Carolina 27419

CGA 81L3D

049