

BEST AVAILABLE COPY

GAVICIDE® SUPER 90

**"440 SPRAY OIL"
— NARROW RANGE —
FOR FRUIT & NUT CROPS**

Active Ingredient:	Unulfonated Residue:
Mineral Oil 99.0%	(ASTM D-483) Min. 93%
Inert Ingredients 1.0%	Petroleum Oil Classification:
Total 100.0%	Unclassified

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (& DOMESTIC ANIMALS)

Caution: Do not get on skin, eyes, or clothing. Harmful if swallowed or inhaled. Avoid breathing spray mist. NOTE TO PHYSICIAN - Emergency information: Call CHEMTREC (800) 424-9300.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on the label.

This product is slightly toxic to bees exposed to direct treatment. Can be applied at any time but bees are temporarily incapacitated by direct contact and some loss may occur. Consult your Agricultural Extension Service for further protective information.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store only in original container in locked storage areas. If a spill occurs, visible puddles or drops should be covered with a dry absorbent such as clay, sawdust, lime or soda ash. Sweep-up and dispose of in a permitted disposal facility. Avoid cross contamination with other pesticides and fertilizers during handling and storage. Do not store near heat or open flame. Protect from temperatures below 0°F. Not for use or storage in or around the home.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not reuse container.

In case of spill or accident, call CHEMTREC (800) 424-9300 day or night for assistance.

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON THE LEFT PANEL

STATEMENT OF PRACTICAL TREATMENT

If swallowed do not induce vomiting, call a physician immediately.
If on skin, immediately remove contaminated clothing and flush skin with plenty of water.
If inhaled, remove victim to fresh air and get medical attention if symptoms persist.
If in eyes, rinse with plenty of water. Get medical attention if irritation persists.

WARRANTY LIMITATIONS AND DISCLAIMER

1. Unocal warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. Unocal neither makes, nor authorizes any agent or representative to make, any other warranty of fitness or of merchantability, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond Unocal's control prevent it from eliminating all risks in connection with the use of this material. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for uses stated herein and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage, and use of this material (except those assumed by Unocal in (1) above). Unocal shall not be liable for incidental or consequential damages.

GAVICIDE is a registered trademark of Union Oil Company of California.

Manufactured by/for



Unocal Petroleum Products & Chemicals Division
Union Oil Company of California dba Unocal
P.O. Box 2390 • Brea, CA 92622-2390 • (800) 825-0076

EPA Reg. No. 612-10

EPA Est. No. _____

Net Contents: _____ Gallons

GENERAL DIRECTIONS

GAVICIDE SUPER 90 is a straight-cut, narrow range petroleum oil designed specifically for use on fruit and nut crops.

DILUTION: Dilute recommended dosage in 50-200 gallons water per acre for ground equipment and in 1-30 gallons for aircraft as needed for coverage.

Fill spray tank with 1/2 the required amount of water. Start agitator. Add total amount of spray oil. Continue to agitate while adding balance of water.

COMPATIBILITY: GAVICIDE SUPER 90 may be tank mixed with insecticides or fungicides. To assure compatibility of tank mixtures, mix the correct proportion of the tank mix components in a small container. After thorough mixing, let stand for five minutes. If the combination remains mixed or can be readily remixed, the tank mixture is compatible.

CHEMIGATION: DO NOT APPLY this product through any type of irrigation system.

APPLICATIONS: GAVICIDE SUPER 90 is formulated as a "fast break" spray oil and will not stay in solution unless agitated constantly. Keep agitator running at all times during spray operations. Apply so as to give full coverage to all parts of crop. Thorough drenching of trunks, limbs, twigs and leaves is essential for optimum results.

DONOT COMBINE this product with sulfur nor within 7 days before or after an application of sulfur unless allowed by the RECOMMENDATIONS of this product and the Directions for Use of the sulfur product.

DO NOT APPLY to trees weakened by disease, drought, drying winds, or high nitrogen applications.

DO NOT USE GAVICIDE SUPER 90 when temperatures are above 90°F Fahrenheit.

RECOMMENDATIONS

Apply as pests appear. Adjust dosage within recommended range according to pest infestation, canopy size, crop condition, etc.

**ALMONDS, APRICOTS, APPLES,
CHERRIES, NECTARINES, PEACHES,
PEARS, PLUMS, PRUNES:**

Dormant/Delayed Dormant Applications:

To control Scales such as Black Scale, Brown Apple Scale, Frosted Scale, Italian Pear Scale, Lecan Scale, Oleander Scale, Olive (parietaria) Scale, Jose Scale, Soft Brown Scale, etc., apply 5-8 g GAVICIDE SUPER 90 per acre.

ACCEPTED with COMMENTS In EPA Letter Dated

MAR 24 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 612-10

612-10
1202-178-3
Transpnd n

Ridomil PC

11G

50 Pounds
Net Weight

GRANULAR FUNGICIDE

For the control of diseases of beans, cotton, and
peanuts caused by *Pythium* and *Rhizoctonia*

Active Ingredients: Pentachloronitrobenzene
(PCNB) 10.0%
Metalaxyl: *N*-(2,6-dimethylphenyl)-*N*-(methoxyacetyl)
alanine methyl ester 1.0%

Inert Ingredients: 89.0%

Total: 100.0%

Ridomil PC 11G is a granular fungicide.

EPA Reg. No. 100-664

EPA Est. 5905-GA-01

Product ID: **54810**

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and direc-
tions for use on back of tag.

cibaTM

BEST AVAILABLE COPY

CGA 130-718A1

444

BEST AVAILABLE COPY

GAVICIDE® SUPER 90

**"440 SPRAY OIL"
— NARROW RANGE —
FOR FRUIT & NUT CROPS**

DIRECTIONS FOR USE

GENERAL DIRECTIONS

is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GAVICIDE SUPER 90 is a straight-cut, narrow boiling range petroleum oil designed specifically for use on fruit and nut crops.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only certified handlers may be in the area during application. Follow any requirements specific to your State or Tribe, and the agency responsible for pesticide regulation.

DILUTION: Dilute recommended dosage in 50-200 gallons water per acre for ground equipment and in 15-30 gallons for aircraft as needed for coverage.

Agricultural Use Requirements

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR part 155. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, safety, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Fill spray tank with 1/2 the required amount of water. Start agitator. Add total amount of spray oil. Continue to agitate while adding balance of water.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

COMPATIBILITY: GAVICIDE SUPER 90 may be tank mixed with insecticides or fungicides. To assure the compatibility of tank mixtures, mix the correct proportions of the tank mix components in a small container. After thorough mixing, let stand for five minutes. If the combination remains mixed or can be readily remixed, the tank mixture is compatible.

Protective clothing is required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

CHEMIGATION: DO NOT APPLY this product through any type of irrigation system.

- Coveralls
- Waterproof gloves
- Shoes plus socks

APPLICATIONS: GAVICIDE SUPER 90 is formulated as a "fast break" spray oil and will not stay in solution unless agitated constantly. Keep agitator running at all times during spray operations. Apply so as to give total coverage to all parts of crop. Thorough drenching of trunks, limbs, twigs and leaves is essential for optimum results.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

DO NOT COMBINE this product with sulfur nor within 30 days before or after an application of sulfur unless allowed by the RECOMMENDATIONS of this product and the Directions for Use of the sulfur product.

SPRAYER: Store only in original container in locked storage areas. If a spill occurs, visible puddles or drips should be covered with a dry absorbent such as clay, kieselguhr, lime or soda ash. Sweep-up and dispose of in an approved disposal facility. Avoid cross contamination with other pesticides and fertilizers during handling and storage. Do not store near heat or open flame. Protect temperatures below 0°F. Not for use or storage in or around the home.

DO NOT APPLY to trees weakened by disease, drought, drying winds, or high nitrogen applications.

CONTAINER DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

DO NOT USE GAVICIDE SUPER 90 when temperatures are above 90°F Fahrenheit.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not reuse container.

RECOMMENDATIONS

Apply as pests appear. Adjust dosage within recommended range according to pest infestation, crop canopy size, crop condition, etc.

ALMONDS, APRICOTS, APPLES, CHERRIES, NECTARINES, PEACHES, PEARS, PLUMS, PRUNES:

Dormant/Delayed Dormant Applications:
To control Scales such as Black Scale, Brown Apricot Scale, Frosted Scale, Italian Pear Scale, Leucanium Scale, Oleander Scale, Olive (parlatoria) Scale, San Jose Scale, Soft Brown Scale, etc., apply 5-8 gals. GAVICIDE SUPER 90 per acre.

Active Ingredient:	Unulfonated Residue:
Mineral Oil 99.0%	(ASTM D-483) Min. 93%
Inert Ingredients 1.0%	Petroleum Oil Classification:
Total 100.0%	Unclassified

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON THE LEFT PANEL

STATEMENT OF PRACTICAL TREATMENT

If swallowed do not induce vomiting, call a physician immediately.
If on skin, immediately remove contaminated clothing and flush skin with plenty of water.
If inhaled, remove victim to fresh air and get medical attention if symptoms persist.
If in eyes, rinse with plenty of water. Get medical attention if irritation persists.

WARRANTY LIMITATIONS AND DISCLAIMER

- Unocal warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated when used in accordance with the use directions under normal conditions. Unocal neither makes, nor authorizes any agent or representative to make, any other warranty of fitness or of merchantability, guarantee or representation, express or implied, concerning this material.
- Critical and unforeseeable factors beyond Unocal's control prevent it from eliminating all risks in connection with the use of this material. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for uses stated herein and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability resulting from the handling, storage, and use of this material (except those assumed by Unocal in (1) above). Unocal will not be liable for incidental or consequential damages.

GAVICIDE is a registered trademark of Union Oil Company of California.

Manufactured by/for



Unocal Petroleum Products & Chemicals Division
Union Oil Company of California dba Unocal
P.O. Box 2390 • Brea, CA 92622-2390 • (800) 825-0076

ACCEPTED
with COMMENTS
In EPA Letter Dated
12-24-78

Under the Federal Insecticide Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 1202-178

EPA Reg. No. 612-10 EPA Est. No. _____

Net Contents: _____ Gallons

1. Time to cover 1,000 ft. at:
 - 2 mph = 5 min. 40 sec.
 - 3 mph = 3 min. 54 sec.
 - 4 mph = 2 min. 50 sec.
 - 5 mph = 2 min. 16 sec.
 - 6 mph = 1 min. 57 sec.

2. Ounces (wt.) of granules to approximate ounces (volume):

6 oz. wt. = 8 fl. oz. (1 level cup)

When calibrating granular applicators, follow the recommendations below:

1. To reduce granule flow variability, adjust orifice openings by moving the controls in one direction only, such as from the closed position to open.
2. Granule flow through an orifice is not always proportional to ground speed, so make calibration checks at the same speed at which you plant and maintain that ground speed as constantly as possible when planting.
3. Granular pesticide products vary considerably in size, density, and particle shape, so recalibrate applicators whenever you change brands and/or formulations. The same brand can even change from year to year.
4. Changing humidity and temperature during the day can change the flow rate of granules through the applicator's orifice. Recalibrate when conditions change and periodically readjust orifice settings to compensate.
5. Know the recommended rate. For band applications, the area actually is the area covered by the band, not total cropland area planted. Some row-crop recommendations are based on treating in-the-row and these rates generally are specified as amounts (ounces) or granules per certain row length (often 1,000 ft.).
6. Calibrate frequently. There are several acceptable calibration methods. Learn one. Do the arithmetic first, then complete the material collection phase under field conditions.

Calculate the amount of Ridomil PC 11G needed for band treatment as follows:

$$\frac{\text{Band Width in Inches}}{\text{Row Spacing in Inches}} \times \text{Broadcast Rate Per Acre} = \text{Band Rate Per Acre}$$

Beans

Ridomil PC 11G will control damping-off and seed and seedling rots of dry and green beans (bush and pole types) caused by *Pythium* and *Rhizoctonia* when used at planting.

Apply $\frac{3}{4}$ lb. (12 oz.) of Ridomil PC 11G/1,000 linear ft. of row at the time of planting. Adjust the application equipment so the granules are mixed with the soil surrounding the seed.

Note: To avoid possible illegal residues, do not allow the feeding of vines or grazing of foliage by livestock.

Cotton

Ridomil PC 11G will control damping-off, seed and seedling rot diseases caused by *Pythium* spp. and *Rhizoctonia solani*.

For best results, use high quality seed and plant into a properly prepared seedbed.

Apply 7-10 lbs. of Ridomil PC 11G/13,000 linear ft. of row (8.6-12.3 oz./1,000 linear ft.) at the time of planting. Adjust the application equipment so the granules are mixed with the soil surrounding the seed. See Chart 1 for suggested applicator settings and the **General Information** section of this label for calibration instructions.

Note: To avoid possible illegal residues, do not allow the feeding or grazing of cotton foliage by livestock.

BEST AVAILABLE COPY

Ridomil PC® 11G

4 7 6

Chart 1: Cotton — Suggested Applicator Settings

Applicator Brand	Speed MPH	40" Rows In-Furrow Lbs./A			
		10	9	8	7
		Equivalent Oz./1,000 ft.			
		12.3	11.1	9.8	8.6
Gandy	4	28	26.8	25	23.5
	5	31	30	28	26.3
	6	34.5	32.5	30.5	29
Allis Chalmers 1979-1985 International Harvester 1980 and Before	4	2.4	2.1	1.9	1.6
	5	3	2.8	2.4	2.05
	6	3.6	3.2	2.9	2.5
Noble 1972 and Before	4	19	17	15	13
	5	23	21	19	17
	6	27	25	22	20
Noble (Remcor) 1973-Present White 5100 Series Deutz Allis 1985-Present	4	14	13	12 or 11	10
	5	17 or 16	15	14	13 or 12
	6	19	18	16	15 or 14
John Deere Metal Box	4	2/12	2/10	2/8	2/5
	5	2/18	2/15	2/12	2/9
	6	2/22	2/19	2/16	2/13
John Deere 7000 and 7100 Max-Emerge® Meter Scale 0...10...20	4	28	27	24	22
	5	31	30	28	26
	6	33	32	31	29
John Deere 7000 and 7100 Max-Emerge® Meter Scale 0...5...15	4	18	17	16	15
	5	21	20	18	17
	6	23	22	21	19
International Harvester 1981-1985 Case IH 1985-Present	4	3.1	2.9	2.6	2.35
	5	3.6	3.4	3.1	2.8
	6	4.5	4	3.55	3.2

Peanuts

Ridomil PC 11G applied at planting will provide control of seed and seedling rots of peanuts caused by Pythium and Rhizoctonia. Applications of Ridomil PC 11G during early pegging will provide control of early pod rot caused by Pythium and Rhizoctonia and will provide suppression of *Sclerotium rolfsii* during early pegging. These treatments will not provide late season control of these diseases.

Seed and Seedling Rots: Apply 12.5-25 lbs. of Ridomil PC 11G/14,520 linear ft. (36-inch row) in a 4-inch band at the time of planting and mix in the covering soil. Incorporate with mechanical equipment to a depth of 2-4 inches. For double-drilled or double-row plantings, apply 12.5-25 lbs./14,520 linear ft. in a 4-inch band centered on each drill or row. Use the higher rate in fields with a history of seedling disease problems or when weather conditions are favorable for infection and disease development.

Pod Rot: Apply 50 lbs./14,520 linear ft. (36-inch row) at early pegging. For banded applications, adjust the equipment so the granules are distributed uniformly over the row(s) and in the pegging zone. For mid- to late-season control of pod rot, use other EPA-registered fungicides.

Sclerotium rolfsii: When Ridomil PC 11G is used for pod rot control, some suppression of *Sclerotium rolfsii* will be obtained. In areas where *Sclerotium rolfsii* is a problem, use other fungicides that are labeled for this disease. Use other products labeled for mid- and late-season control of southern blight/white mold infections.

BEST AVAILABLE COPY

Ridomil PC[®] 11G

5 of 6

Notes: To avoid possible illegal residues, (1) Do not allow the feeding or grazing of peanut foliage or harvest residues by livestock. (2) Do not apply more than 75 lbs. of Ridomil PC 11G/14,520 linear ft. of row per season. (3) Do not apply within 75 days of harvest.

Rotational Crops

These crops represent those which may be planted (rotated to) according to the interval listed following an application of Ridomil PC 11G to beans, cotton, or peanuts. Do not plant any crop not registered for use with metalaxyl in soil treated with this product for a period of 12 months. Do not plant root crops in fields treated with this product within 12 months of application.

Crop	Rotational Interval
Alfalfa, Almonds, Apples, Asparagus, Avocados	0 days
Beans (Dry and Snap), Blueberries, Broccoli	
Cabbage, Cauliflower, Citrus, Cotton, Cucurbit Vegetables	
Head Lettuce, Hops	
Legume Vegetables Other than Beans	
Onions	
Peanuts, Peppers, Pineapples	
Raspberries	
Soybeans, Spinach, Stone Fruits, Strawberries	
Tomatoes	
Walnuts	
Brussels Sprouts	12 months
Corn	
Garlic	
Root Crops	
Wheat	
All Other Crops Intended for Food or Feed	

Storage and Disposal

Pesticide

Store in a dry place. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide is a violation of federal law. Pesticide that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Disposal

Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

BEST AVAILABLE COPY

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust.

Statement of Practical Treatment

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention.

If inhaled: Move to fresh air.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Cover or incorporate all granules. Do not contaminate water when disposing of equipment wash waters.

Ridomil PC[®] trademark of Ciba-Geigy Corporation

Max-Emerge[®] trademark of John Deere Company

©1994 Ciba-Geigy Corporation

Ciba Plant Protection

Ciba-Geigy Corporation

Greensboro, North Carolina 27419

CGA 81L3K

014

BEST AVAILABLE COPY