

100-669

3-6-2003

1/22

(Booklet for 15 gal container)

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY
 CERTIFIED APPLICATORS OR PERSONS
 UNDER THEIR DIRECT SUPERVISION, AND ONLY
 FOR THOSE USES COVERED BY THE
 CERTIFIED APPLICATOR'S CERTIFICATION.
 DUE TO HIGH TOXICITY.

Curacron® 8E

Insecticide-Miticide

Closed System

For insect and mite control in cotton

ACCEPTED

MAR 6 2003

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 100-669

Active Ingredient:

Profenofos (CAS No. 41198-08-7).....	73.0%
Other Ingredients*:	27.0%
Total:	100.0%

*Contains petroleum distillates

Curacron 8E contains 8 lbs. active ingredient per gal.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entienda la etiqueta busque a alguien para se la explique en detalle. (If you do not understand the label find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-669

EPA Est.

SCP 669A-L5 Pending

3/05/03

15 gallons
U. S. Standard Measure

2/22

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if absorbed through skin. Causes eye irritation. Do not get in eyes, on skin, or on clothing. Avoid breathing vapors or spray mist. Harmful if swallowed or inhaled.

FIRST AID	
This product contains an organophosphate that inhibits cholinesterase	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 – 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible • Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN	
Contains petroleum distillate – vomiting may cause aspiration pneumonia.	
This product contains profenofos, an organophosphate that inhibits cholinesterase. Atropine, only by injection, is the preferable antidote. cyanosed patients must be given oxygen first; atropine must not be given to anoxic patients. Oximes such as 2-PAM or toxogonin are also antidotal and may be administered, but only in conjunction with atropine. Avoid the use of respiratory depressants e.g. barbiturates and opiates.	
Have the product container or label with you when calling a poison control center or doctor or when going for treatment.	
HOT LINE NUMBER	
For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

3/22

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category F on an EPA chemical-resistant category selection chart.

Mixers, loaders, applicators, flaggers, and other handlers using engineering controls (see requirements below) must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber or viton and chemical-resistant apron when mixing and loading
- Protective eyewear

Handlers, for which use of an engineering control is not possible or when engaged in activities such as cleaning up a spill or leak or cleaning or repairing contaminated equipment must wear:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber or viton,
- Chemical-resistant footwear plus socks,
- Chemical-resistant apron
- Protective eyewear

All mixers, loaders, applicators, and flaggers (except aerial applicators) must be provided and must have immediately available for use in an emergency, such as a spill, or equipment failure or if exiting a cab in a treated area, the following personal protective equipment (PPE)

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, or viton.
- Chemical-resistant footwear plus socks,
- Chemical resistant apron
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

4/22

Engineering Control Statements

Closed System

- Mixers and loaders must use a mechanical transfer system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)] for providing dermal protection.
- The system must be capable of removing the pesticide from the shipping container and transferring it into mixing tanks and/or application equipment.
- At any disconnect point, , the system must be equipped with a dry disconnect or dry couple shut-off device that is warranted by the manufacturer to minimize drippage to not more than 2 ml. per disconnect point.
- Persons using a closed system that operates under pressure shall wear protective eyewear.

Enclosed Cab/Cockpit

Pilots must:

- use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)];

Ground equipment applicators and flaggers must:

- use an enclosed cab/cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5)].

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ADVISORY TO CERTIFIED CROP ADVISORS: Users should inform Certified Crop Advisors [as defined by the Worker Protection Standard (WPS)] that people engaged in scouting activities should wear coveralls, shoes and socks, and chemical-resistant gloves made of any waterproof material when entering treated areas during the first 48 hours following application (72 hours in areas where the average rainfall is less than 25 inches per year).

5/22

Environmental Hazards

This pesticide is acutely toxic to fish and wildlife. Use with care when applying to areas frequented by wildlife or adjacent to any body of water. Due to the hazard to endangered fish, the application of this product is prohibited in Reeves County, TX and within one mile of the Dexter National Fish Hatchery in NM. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposing of equipment wash water. Do not apply to saturated soil and do not treat while precipitation is occurring or while conditions favor runoff from the treated area due to the potential for surface water runoff that may cause fish kills. Do not apply with aircraft within 300 ft. or with ground boom equipment within 100 ft of any body of water including impounded waters, rivers, streams, lakes, or oceans. Do not apply with aircraft when wind speed is greater than 10 mph. Use vegetative buffers

This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply this product or allow it to drift to crops or weeds on which bees are foraging. Additional information may be obtained from your Cooperative Agricultural Extension Service.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

6/22

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

7/22

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this pesticide only as specified on this label.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. The REI is 72 hours in areas where average rainfall is less than 25 inches per year.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes and socks
- Protective eyewear

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

General Application Restrictions:

- 1) Cotton treated with this product must be mechanically harvested. Hand harvesting is prohibited.
- 2) Pilots are prohibited from participating in mixing or loading on the day of application.
- 3) Ground boom and aerial applicators must use enclosed cabs and cockpits.
- 4) Do not allow this product to drift.
- 5) Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.
- 6) Chemigation: Do not apply this product through any type of irrigation system.

**FAILURE TO FOLLOW ALL DIRECTIONS AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN CROP INJURY, POOR CONTROL, AND/OR ILLEGAL RESIDUES.**

8/22

GENERAL INFORMATION

Curacron 8E is an emulsifiable insecticide-miticide which is effective for control of cotton bollworm, tobacco budworm, certain other insects, and mites on cotton.

Best results can be expected from fields under Pest Management/ Scouting programs.

When using Curacron 8E as a larvicide, best results will be obtained when application is timed to control first to third instar larvae. For information on use practices, refer to the Cotton Council website: <http://www.carefulbynature.org>

Curacron 8E is an organophosphate insecticide which has a specific mode of action and could be subject to development of resistant strains of insects. Development of resistance cannot be predicted. Therefore, Syngenta cannot assume liability for crop damage resulting from resistant strains of insects. If the treatment is ineffective due to the presence of resistant strains, neither Curacron 8E nor any other insecticide with similar action will effectively control those pests. Consideration should then be given to the prompt use of other types of suitable insecticides. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and information on resistance management strategies in your area.

ROTATIONAL CROPS

Fields treated with Curacron 8E may be rotated to other crops 30 days after the last application.

SPRAY DRIFT

Aerial Spray Drift Management

A. Responsibility

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment - and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering these factors when making decisions about aerial application

B. Drift Management Requirements

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops.

- 1) Nozzles must always point backwards parallel with the air stream and never be pointed downwards more than 45 degrees.
- 2) The distance of the outermost nozzles on the boom must not exceed ¼ of the length of the wingspan or rotor.
- 3) Do not apply with aircraft when wind speed is greater than 10 mph.

9/22
3

When states have more stringent regulations, they should be followed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information section that follows.

C. Aerial Drift Reduction Advisory

This section is advisory in nature and does not take the place of the mandatory requirements in the **Aerial Spray Drift Management** section

1. Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or made under unfavorable environmental conditions (see Wind, Temperature, and Humidity and Temperature Inversions).

2. Controlling Droplet Size

- a. Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher-rated flows produce larger droplets.
- b. Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- c. Number of nozzles – Use the minimum number of nozzles that provide uniform coverage.
- d. Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the air stream produces larger droplets than other orientations and is recommended. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- e. Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the least drift.

3. Boom Length

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

4. Application Height

Applications should not be made at a height greater than 10 feet above the top of the tallest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

10/22

5. Swath Adjustment

When applications are made during a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

6. Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

7. Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are hot and dry.

8. Temperature Inversions

Applications should not be made during a temperature inversion because drift potential is high.

Temperature inversions:

- a. restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light, variable winds that are common during inversions,
- b. are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind,
- c. begin to form as the sun sets and often continue into the morning,
- d. can be indicated by ground fog; however if fog is not present, inversions can be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind condition) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

9. Sensitive Areas

Curacron 8E should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or non-target crops) is minimal (for example, when wind is blowing away from the sensitive area).

MIXING INSTRUCTIONS

- 1) Prepare no more spray mixture than is needed for the immediate operation.
- 2) Thoroughly clean spray equipment before using this product.
- 3) Agitate thoroughly before and during application. Good bypass agitation is sufficient to keep Curacron 8E in emulsion.
- 4) Do not let spray mixture stand overnight in the spray tank.
- 5) Rinse spray tank at the end of the day.

Curacron 8E Alone

- 1) Add 2/3 of the required amount of water to the spray tank with the agitator running; then add the desired amount of Curacron 8E.
- 2) Continue agitation while adding the remainder of the water.

Curacron 8E + Tank Mixtures

NOTE: Tank mixtures are permitted only in those states where the tank mix partner is registered.

Add 2/3 of the required amount of water to the spray tank. With the agitator running, if possible, add the desired amount of the other products recommended on this label. Then add the desired amount of Curacron 8E. Continue agitation while adding the remainder of water.

Curacron 8E + Foliar Nutrients

Some foliar fertilizers can increase chances for phytotoxicity or decrease performance when tank mixed with Curacron 8E. Syngenta cannot be held responsible for the performance of Curacron tank mixed with foliar fertilizers.

APPLICATION

Buffer Zone Restrictions

Follow the restrictions listed below when applying Curacron 8E near any body of water, including impounded waters, rivers, streams, lakes, and oceans.

- 1) Do not apply with aircraft within 300 ft of any body of water.
- 2) Do not apply with ground boom equipment within 100 ft. of any body of water.

12/22
***Maximum Use Rates**

- 1) Do not apply more than 5 pts (5 lb ai)/A per season.
- 2) Do not apply more than $\frac{3}{4}$ pt (0.75 lb ai/A) per application.

NOTE: For lepidopteran pests only (cotton bollworm, tobacco budworm, beet armyworm, fall armyworm), you may use 1 pt (1.0 lb ai/A) per application with a maximum of 2 applications per season.

For information on use practices refer to the Cotton Council website:
<http://www.carefulbynature.org/>

CROP USE DIRECTIONS**Pre Harvest Interval (PHI)**

Do not apply Curacron 8E within 14 days of harvest.

Curacron 8E Alone

Apply Curacron 8E at the specified rates below in a minimum of 5 gals. of water/A with ground equipment, or in a minimum of 1 gal. of water/A by air. In late season, under heavy infestations and high temperatures, apply Curacron 8E in a minimum of 3 gals. of water/A to ensure maximum coverage. Avoid spray overlap because crop injury may result.

If control of boll weevil is needed while using Curacron 8E, Guthion® or methyl parathion may be tank mixed with Curacron 8E. Refer to the Guthion or methyl parathion labels for directions, limitations, and precautions.

13/22

PEST	LOCATION	AMOUNT OF CURACRON 8E TO APPLY	HOW OFTEN TO APPLY	INSTRUCTIONS
Aphids	All cotton growing states	½ pint /A	5-7 day intervals	Begin application when economic thresholds are reached and continue at 5-7 day intervals. Economic thresholds will vary by locality. Consult your pest control advisor or cotton consultant to determine economic thresholds in your area.
Plant bugs, lygus bugs*	All cotton growing states. *In AZ and CA Curacron offers suppression of lygus bugs	¼ to ½ pint/A		
Cotton Bollworm	All cotton growing states except AZ and CA	<u>Contact Ovicide Activity:</u> 1/8 to ¼ pint/A	5-7 day intervals	Use the higher rate if plant bugs are also present.
		<u>Larvae Control Only:</u> ¼ to 1 pint/A NOTE: Do not make more than 2 applications per season at the 1 pt/A rate	5-7 day intervals 5-6 day intervals may be necessary when infestation is heavy	Apply 1 pt/A on large larvae and when infestation is heavy. Scout fields thoroughly and repeat application at a 5-7 intervals. 5-6 day intervals may be needed when infestation is heavy.
Tobacco Budworm	All cotton growing states except AZ and CA	<u>Contact Ovicide Activity:</u> ¼ pint/A	5-7 day intervals	
		<u>Larvae Control Only:</u> ¼ to 1 pint/A NOTE: Do not make more than 2 applications per season at the 1 pt/A rate	5-7 day intervals 5-6 day intervals may be necessary when infestation is heavy	Apply 1 pt/A on large larvae and when infestation is heavy. Scout fields thoroughly and repeat application at a 5-7 intervals. 5-6 day intervals may be needed when infestation is heavy.

14/22

Cotton bollworm, Tobacco budworm	AZ and CA	½ to ¾ pint/A	5-7 day intervals	Use this application rate for light to moderate infestations
		¾ to 1 pint/A NOTE: Do not make more than 2 applications per season at the 1 pt/A rate.	5-7 day intervals	Use this application rate for moderate to heavy infestations
Cotton leafperforator, Beet armyworm	All cotton growing states	1 pint/A NOTE: Do not make more than 2 applications per season at the 1 pt/A rate.	5-7 day intervals 5-6 day intervals may be necessary when infestations are heavy	Begin application when egg masses and larvae first appear. Scout fields thoroughly to determine if repeat application is needed.
Fall armyworm	All cotton growing states	¾ to 1 pint/A NOTE: Do not make more than 2 applications per season at the 1 pt/A rate	5-7 day intervals	Begin application when larvae first appear. For best results, apply to young larvae still feeding on leaf surfaces and before the larvae migrate to blooms and bolls.
Mites	All cotton growing states	½ to ¾ pint/A	5-7 day intervals	Begin when mites first appear.
Whitefly Suppression	All cotton growing states	¾ pint/A	5-7 day intervals	Curacron 8E will aid in the reduction of whitefly populations. Begin treatment when whiteflies first appear.

15/22

Curacron 8E - Tank Mixtures (All cotton growing states)

If multiple pests, including boll weevil, are present, it may be useful to tank mix Curacron 8E with other EPA-registered cotton pesticides to broaden the spectrum of control.

When using Curacron 8E in tank mixtures, observe all limitations, precautions, and use rates that appear on the labels for these products. Curacron 8E is compatible in tank mixtures with the following pesticides:

- | | |
|----------------|------------------|
| Ammo® 2.5EC | Lorsban® 4E |
| Asana® XL | Malathion |
| Baythroid 2® | Meta-Systox R® |
| Capture® 2EC | Monitor® 4 Spray |
| Condor® | Mustang™ |
| Danitol® 2.4EC | Orthene® 75S |
| Dipel® ES | Ovasyn® |
| Dipel® 2X | Parathion |
| Fury® 1.5EC | Phaser® 3EC |
| Guthion® 2L | Pounce® 3.2EC |
| Javelin® WG | Provado™ |
| Karate® | Scout X-TRA® |
| Lannate® | Supracide® |
| Larvin® 3.2 | Thiodan® 3EC |
| Lock-On® | Vydate® L |

NOTE: Applications to cotton may cause spotting or reddening of leaves.

Curacron 8E + Oil (All cotton growing states)

Apply Curacron 8E following use directions above, selecting the amount to use and the application intervals according to the pest problem.

MIXTURE	AMOUNT TO USE
For ULV application, dilute Curacron 8E + once refined vegetable oil	Apply using 1-2 qts of finished spray per acre
Curacron 8E + water + emulsified vegetable oil	Add 1-2 pts of emulsified vegetable oil as an additive. Apply using a minimum of 1 gal total mixture per acre

Important: When applying Curacron 8E in oil:

- 1) If the use rate permits 3 or more applications per season, make no more than 3 applications of Curacron 8E in oil per season.
- 2) Do not apply within 30 days of harvest..

16/22

GRAZING RESTRICTIONS

- 1) Do not allow livestock to graze treated cotton.
- 2) Do not feed gin trash or foliage from treated cotton plants to livestock.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

Emergency Response

In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Curacron®, Karate® and the Syngenta logo are trademarks of a Syngenta Group Company
U.S. Patent No. 3,992,533

Ammo®, Capture®, Fury®, Mustang®, Pounce®, and Thiodan® trademarks of FMC Corporation

Asana®, Lannate®, and Vydate® trademarks of E. I. duPont de Nemours and Company

Baythroid®, Guthion®, and Provado® trademarks of Bayer Corporation

Condor® and Javelin® trademarks of Certis USA

Danitol®, Dipel®, Monitor®, and Orthene® trademarks of Valent USA Corporation

Larvin®, Ovasyn®, Phaser®, and Scout-XTRA® trademarks of Aventis Group

Lock-On® and Lorsban® trademarks of Dow AgroSciences

Meta-Systox R® trademark of Bayer Corporation, licensed to Gowan Company

Curacron® trademark of Curacron, Inc. licensed to Gowan Company

17/22

©2003 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 669A-L5 (Pending)

CFB 2/27/03

18/22

(container label 15 gal. - PSL)

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY
 CERTIFIED APPLICATORS OR PERSONS
 UNDER THEIR DIRECT SUPERVISION, AND ONLY
 FOR THOSE USES COVERED BY THE
 CERTIFIED APPLICATOR'S CERTIFICATION.

Curacron® 8E

Insecticide-Miticide

Closed System

For insect and mite control in cotton

Active Ingredient:

<u>Profenofos (CAS No. 41198-08-7)</u>	<u>73.0%</u>
<u>Other Ingredients*</u>	<u>27.0%</u>
Total:	100.0%

*Contains petroleum distillates

Curacron 8E contains 8 lbs. active ingredient per gal.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-669

EPA Est.

SCP 669A-L6 Pending (15 gal. Container label)

15 gallons
U. S. Standard Measure

19/22

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if absorbed through skin. Causes eye irritation. Do not get in eyes, on skin, or on clothing. Avoid breathing vapors or spray mist. Harmful if swallowed or inhaled.

FIRST AID	
This product contains an organophosphate that inhibits cholinesterase	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 – 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible • Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN	
Contains petroleum distillate – vomiting may cause aspiration pneumonia.	
This product contains profenofos, an organophosphate that inhibits cholinesterase. Atropine, only by injection, is the preferable antidote. Cyanosed patients must be given oxygen first; atropine must not be given to anoxic patients. Oximes such as 2-PAM or toxogonin are also antidotal and may be administered, but only in conjunction with atropine. Avoid the use of respiratory depressants e.g. barbiturates and opiates.	
Have the product container or label with you when calling a poison control center or doctor or when going for treatment.	
HOT LINE NUMBER	
For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

20/22
1

Environmental Hazards

Apply this pesticide only as specified on this label. This pesticide is acutely toxic to fish and wildlife. Use with care when applying to areas frequented by wildlife or adjacent to any body of water. Due to the hazard to endangered fish, the application of this product is prohibited in Reeves County, TX and within one mile of the Dexter National Fish Hatchery in NM. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposing of equipment wash water. Do not apply to saturated soil and do not treat while precipitation is occurring or while conditions favor runoff from the treated area due to the potential for surface water runoff that may cause fish kills. Do not apply with aircraft within 300 ft. or with ground boom equipment within 100 ft of any body of water including impounded waters, rivers, streams, lakes, or oceans. Do not apply with aircraft when wind speed is greater than 10 mph. Use vegetative buffers

This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply this product or allow it to drift to crops or weeds on which bees are foraging. Additional information may be obtained from your Cooperative Agricultural Extension Service.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

21/22
3

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

22/22
3

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

Curacron® and the Syngenta logo are trademarks of a Syngenta Group Company
U.S. Patent No. 3,992,533

©2003 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 669A-L6 Pending