10/2

Supplemental Labeling Dual Herbicide EPA Reg. No. 100-673

4 Rodenticide Act.

Active Ingredient:

Metolachlor: 2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-

methylethyl) acetamide

Inert Ingredients: 13.6% 100.02

Total:

Keep Out of Reach of Children

WARNING/AVISO

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

General Information

Observe all precautions and limitations on the labels of each product used in tank mixtures.

Dual is a selective herbicide recommended as a preplant surface-applied, preplant incorporated, or preemergence treatment in water or fluid fertilizer for control of most annual grasses and certain broadleaf weeds in nonbearing citrus, corn, cotton, nonbearing grapes, peanuts, pod crops, potatoes, safflowers, grain or forage sorghum, soybeans, stone fruits, tree nuts, and woody ornamentals.

Rotational Crops: (1) If crop treated with Dual alone is lost, any crop on this label may be replanted immediately. make a second broadcast application of Dual. If the original application was banded and the second crop is planted in the untreated row middles, a second banded treatment may be applied. (2) Barley, oats, rye, or wheat may be planted 4 1/2 months following treatment; alfalfa may be planted 4 months following application. (3) Any crop on this label in addition to root crops, barley, buckwheat, milo, oats, rice, rye, or wheat may be planted in the spring following treatment. Clovers may be seeded 9 months rullowing application. Do not graze or feed forage or fodder from cotton to livestock. All other rotational crops may be planted 18 months after application. (3) Following a lay-by treatment or multiple treatments applied the previous season, any crop on this label may be planted in the spring. All other rotational crops may be planted 18 months after a lay-by application.

Important Notes: To avoid injury to rotational alfalfa or clover, (1) Do not apply more than 3 lbs. ai of metolachlor (3 pts. Dual) preemergence (including preplant-surface, preplant-incorporated, postplant incorporated, etc.). (2) Do not make lay-by or other postemergent applications of Dual. (3) For Rotational Crops restrictions for Dual used in tank mixtures, refer to the statements/restrictions above for Dual and to the respective product labels of any mixing partner(s) for additional statements/restrictions.

Follow all applicable directions, restrictions, and precautions on the EPA-registered label.

This label must be in the possession of the user at the time of application.

Dual trademark of CIBA-GEIGY Corporation for metolachlor. U.S. Patent No. 3,937,730

Agricultural Division CIBA-GEIGY Corporation P. O. Box 18300 Greensboro, NC 27419

l

February 8, 1988

(LABELS-D.3 - DUAL/SUPPF1- DRAFT)