

Handwritten notes and signatures

Supplemental Labeling
Dual® Herbicide
EPA Reg. No. 100-673

Rotational Tomatoes

Active Ingredient:

Metolachlor: 2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl) acetamide 86.4%

Inert Ingredients: 13.6%

Total: 100.0%

Dual contains 8 lbs. active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING


See additional precautionary statements inside booklet.

AVISO

PRECAUCIÓN AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

All applicable directions, restrictions, and precautions on the EPA-registered Dual label are to be followed.

This label must be in the possession of the user at the time of application.



DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Rotational Crops: Dual Alone: (1) If crop treated with Dual alone is lost, any crop on this label may be replanted immediately. Do not make a second broadcast application of Dual. If the original application was banded and the second crop is planted in the untreated row middles, a second banded treatment may be applied. (2) Barley, oats, rye, or wheat may be planted 4 1/2 months following treatment; alfalfa may be planted 4 months following application. Tomatoes may be planted 6 months following application. (3) Any crop on this label in addition to root crops, tobacco, barley, buckwheat, milo, oats, rice, rye, wheat, cabbage, peppers, stonefruits, or tree nuts may be planted in the spring following application. Clover may be seeded 9 months following application. Do not graze or feed forage or fodder from cotton to livestock. All other rotational crops may be planted 12 months after application. (4) Following a lay-by treatment or multiple treatments applied the previous season, any crop on this label in addition to tobacco, cabbage, peppers, stone fruits or tree nuts may be planted in the spring. All other rotational crops may be planted 12 months after a lay-by application. Dual Tank Mixtures: For Rotational Crops restrictions for Dual used in tank mixtures, refer to the statements/restrictions above for Dual and to the respective product labels of any mixing partner(s) for additional statements/restrictions.

Important Notes: To avoid injury to rotational alfalfa or clover, (1) Do not apply more than 3 lbs. ai of metolachlor/A (3 pts. Dual) preemergence (including preplant-surface, preplant-incorporated, postplant incorporated, etc.). (2) Do not make lay-by or other postemergent applications of Dual.

Dual® trademark of CIBA-GEIGY Corporation for metolachlor.

©1992 CIBA-GEIGY Corporation

CIBA-GEIGY Corporation
Agricultural Division
P. O. Box 18300
Greensboro, NC 27419

July 12, 1989
April 30, 1992

[GANNONC.LABELD - DUAL_SUPPF2] - 4/29/92