



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
100-717

Date of Issuance:
APR 06 1992

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, 3as amended)

Term of Issuance:
Conditional

Name of Pesticide Product:
Subdue II Fungicide

Name and Address of Registrant (include ZIP Code):

Ciba-Geigy Corporation
P.O. Box 18300
Greensboro, North Carolina 27419-8300

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Submit 12 month storage stability data at ambient warehouse conditions. This should be received by the Agency within 15 months after registration of this product.
3. Copies of the raw analytical data samples representing five batches of this product should be provided to the Agency within 12 months after registration of this product.

Signature of Approving Official:

Date:

APR 06 1992

Page 2

EPA Reg. No. 100-717

4. Make the following label change before you release the product for shipment: Add the phrase "EPA Registration No. 100-718".

5. Submit five copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,



Susan T. Lewis
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

plantPak™

ACCEPTED
with COMMENTS
in EPA Letter Dated:

LABELMASTER BOOKLET

APR 06 1992

Subdue® II

Fungicide

Under the Fungicide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 100-717

For control of certain diseases in ornamentals, vegetable plant beds, nonbearing citrus, conifers, and nonbearing deciduous fruit and nut trees in nurseries and landscape plantings.

Active Ingredient:

Metalaxyl: <u>N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester</u>	25.0%
<u>Inert Ingredients:</u>	75.0%
Total:	100.0%

Subdue II is a wettable powder.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-
EPA Est.

This outer protective container contains Subdue II in eight inner water soluble packets. These inner packets dissolve in water, allowing contents to wet. To open, remove lid, then pull tab back to expose packets. Immediately dump the required number of unopened inner packets into the partially filled sprayer or mix tank. Do not handle the soluble packets or expose them to moisture, since this may cause rupturing. When not using the entire number of packets in this container, reclose the container by replacing the protective seal and lid.

Eight X One Ounce Soluble Packets
Total One Half Pound
Net Weight

Do not offer individual packets for sale.

Subdue® trademark of CIBA-GEIGY Corporation.

CIBA-GEIGY

©1991 CIBA-GEIGY Corporation

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, NC 27419

CGA 114L1

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of CIBA-GEIGY or the Seller. All such risks shall be assumed by the Buyer.

CIBA-GEIGY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. CIBA-GEIGY makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall CIBA-GEIGY or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. CIBA-GEIGY and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CIBA-GEIGY.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR PLANT INJURY.

Reentry Statement

Do not apply this product in such a manner as to directly or through drift expose workers or other persons, except those knowingly involved in the application. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various sites treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must inform workers of areas or fields that may not be entered without specific protective clothing until sprays have dried, and appropriate actions to take in case of accidental exposure, as described under Precautionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "CAUTION. Area treated with Subdue II on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental eye contact, flush eyes with plenty of water. Get medical attention. In case of contact with skin, wash thoroughly with soap and water. Call physician if irritation occurs. Remove and wash contaminated clothing before reuse."

General Information

Subdue II is a systemic fungicide for use on ornamentals, vegetable plant beds, nonbearing citrus, conifers, and nonbearing deciduous fruit and nut trees in nurseries and landscape plantings.

BEST AVAILABLE COPY

NOTE: 1) Subdue II is a systemic fungicide having a specific mode of action. Use of Subdue II could result in development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, CIBA-GEIGY cannot assume liability for plant damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Subdue II resistant strains of fungi which may occur. 2) Do not use Subdue II for control of downy mildew diseases. 3) Use Subdue II only as a soil application for control of soil-borne diseases with the exception of Phytophthora blights of rhododendrons (including azaleas).

Mixing Instructions

Prepare no more spray mixture than is required for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate thoroughly before and during application. Good bypass agitation is sufficient to keep Subdue II thoroughly mixed and in suspension. Flush spray tank thoroughly with water daily after use and dispose of pesticide rinsate by application to an already treated area.

Note: Do not use with oils or adjuvants.

Subdue II Alone: Add 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble packets of Subdue II into the tank. Continue agitation while adding the remainder of the water.

Subdue II + Tank Mixtures: Subdue II is usually compatible with Banner®, Benlate®, D·z·n, Tersan 1991®, Daconil 2787®, Tersan LSR®, Fore®, and Dyrene®. Add 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble packets of Subdue II into the tank. Then add the desired amount of the selected tank mix partner(s). Continue agitation while adding the remainder of the water.

Precaution: Soluble packets of Subdue II must be completely dissolved in water before any other tank mix partner, including micronutrients or other liquid or dry fertilizers, is added to the spray solution.

If using Subdue II in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate should be exceeded and the most restrictive label precautions and limitations should be followed. This product should not be mixed with any product which prohibits such mixing.

BEST AVAILABLE COPY

Application Instructions for Use in Irrigation Systems

Overhead, Microjet or Drip Irrigation Application: Subdue II alone or in tank mixture with other pesticides which are registered for overhead, microjet, or drip application may be applied in irrigation water at rates recommended on this label. Apply this product only through an overhead, microjet, or drip irrigation system. Do not apply this product through any other type of irrigation system. Plant injury or lack of effectiveness can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Operation Instructions:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment. Avoid spray overlap as injury may result.

BEST AVAILABLE COPY

8. Prepare a mixture with a minimum of 1 gal. of water to 1 packet of Subdue II and inject this mixture into the system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the fungicide in suspension.
9. Meter into irrigation water during the beginning of the irrigation cycle.
10. With overhead or microjet systems, apply additional water after irrigation is completed to remove residues from the foliage. Apply enough supplemental water to wet the root zones of the plants.

Precaution for irrigation applications: Where distribution patterns do not overlap sufficiently, unacceptable control may result. Where distribution patterns overlap excessively, injury to desirable plants may result.

Ornamentals

Subdue II is for use on container, bench, or bed-grown ornamentals in greenhouses or outdoor nurseries, and for ornamentals grown for indoor and outdoor landscaping. Subdue II controls damping-off and root and stem rot diseases caused by Pythium and Phytophthora. Use Subdue II as a soil drench on ornamentals at the time of seeding, transplanting or to established plants. Subdue II may be applied as a foliar spray on azaleas. Within a rate range given for a specific group of ornamentals, use the lower rate for the shortest interval listed and the higher rate for the longest interval. Under severe disease conditions, use the highest rate and the shortest interval specified.

For use on ornamentals in fewer than 4 inches of soil, apply the specified amount of Subdue II in 100 gals. of water. Apply this solution at a rate of 1 pt./sq. ft. For soil depth greater than 4 inches, apply 1 1/2-2 pts./sq. ft. Do not exceed recommended rates or apply more frequently than directed, since plant injury may occur.

NOTICE TO USER: Plant tolerances to Subdue II have been found to be acceptable for the specific genera and species listed on this label. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for tolerance to Subdue II. Neither the manufacturer nor the seller has determined whether or not Subdue II can be used safely on ornamental and nursery plants not specified on this label. The professional user should determine if Subdue II can be used safely prior to commercial use. In a small area, test the recommended rates for a particular group of unlabeled plants, i.e., bedding plants, foliage, etc., for phytotoxicity prior to widespread use.

<p><u>Foliage Plants</u></p> <p>Aglaonema Aphelandra Dieffenbachia Peperomia Philodendron* Pothos Schefflera Sedum Sempervivum Zygocactus</p>	<p><u>SOIL DRENCH:</u> Mix 1 packet of Subdue II with 100 gals. of water. Apply 1 pt. solution/sq. ft. For soil depth greater than 4 inches, apply 1 1/2-2 pts. solution/sq. ft. Repeat applications at two to three month intervals if necessary.</p> <p>*On Philodendron, use 1-2 packets of Subdue II/100 gals.</p> <p><u>Precaution:</u> To minimize the potential for injury to Pothos, do not use more than 1 packet of Subdue II/100 gals. and do not apply more frequently than once every 3 months.</p>
<p><u>Bedding Plants</u></p> <p>Ageratum Algerian Ivy Artemisia Aster Begonia Caladium Carnation Chrysanthemum Coleus Daisy, Shasta English ivy Foxglove Gaillardia Geranium Impatiens Marigold Pansy Petunia Phlox Pinks Primrose Prostrate Rosemary Salvia Snapdragon Verbena Vinca Zinnia</p>	<p><u>At Transplanting (Soil 2-3 inches deep):</u> Mix 1-2 packets of Subdue II with 100 gals. of water and apply 1 pt. solution/sq. ft. For soil depth greater than 4 inches, apply 1 1/2-2 pts. solution/sq. ft. Repeat applications at one to two month intervals if necessary. Do not apply rates of 1-2 packets of Subdue II/100 gals. more often than once every six weeks.</p> <p><u>Precaution:</u> Do not apply to English ivy more than once every 6 months since injury may occur.</p>

<p><u>Flowers</u></p> <p>African violet Anthurium Baby's breath Carnation Chrysanthemum Columbine Delphinium Easter lily Geranium Gloxinia Poinsettia Rose</p>	<p><u>SOIL DRENCH:</u> Mix 1-2 packets of Subdue II with 100 gals. of water and apply 1 pt. solution/sq. ft. For soil depth greater than 4 inches apply 1 1/2-2 pts. solution sq. ft. Repeat application at one to two month intervals if necessary. Do not apply rates of 1-2 packets of Subdue II/100 gals. more often than once every six weeks.</p> <p><u>Precaution:</u> Do not apply more than one packet of Subdue II/100 gals. water to Easter lily and only make one at-planting application.</p>
<p><u>Rhododendrons and Azalea</u></p>	<p><u>SOIL DRENCH:</u> Phytophthora root and crown rot - Mix 1-2 packets of Subdue II with 100 gals. of water and apply 1 pt. solution/sq. ft. For soil depth greater than 4 inches, apply 1 1/2-2 pts. solution/sq. ft. Repeat applications at two to four month intervals if necessary.</p> <p><u>FOLIAR SPRAY:</u> Phytophthora shoot blight - Mix at 1-2 packets of Subdue II with 100 gals. of water. Spray to runoff. Repeat at two to three month intervals if necessary.</p> <p><u>Precautions:</u> (1) To minimize the potential for injury to azaleas, do not apply repeat soil applications of 2 packets of Subdue II/100 gals. closer than every 3 months, and do not exceed a total of 4 packets of Subdue II in 6 months. (2) Use the lower rate for "Coral Bell" variety.</p>

<u>Woody Ornamentals Other Than Azalea</u> <u>Aucuba japonica</u> Arborvitae Boxwood Ceanothus Cotoneaster Dogwood Ficus "Halls" Honeysuckle Ilex <u>Juniperus spp.</u> Photinia <u>Pieris japonica</u> <u>Pinus spp.</u> Pittosporum Rhododendron White cedar White pine Yew	<u>SOIL DRENCH:</u> Mix 1-4 packets of Subdue II with 100 gals. of water and apply 1 pt. solution/sq. ft. For soil depth greater than 4 inches, apply 1 1/2-2 pts. solution/sq. ft. Repeat applications at two to three month intervals if necessary. Do not apply rates greater than 3 packets of Subdue II/100 gals. more often than once every ten weeks. <u>Precaution: Do not apply to Euonymus or injury may occur.</u>
--	--

Nonbearing Citrus

Use Subdue II on nonbearing citrus for control of citrus foot rot, root rot, and trunk canker caused by Phytophthora spp. Apply to the soil as a drench or as a spray in a banded application.

Make the first application of Subdue II at the time of planting. Make repeat applications at three-month intervals during the period when trees are actively growing.

Soil Drench: Mix 4-6 packets of Subdue II/100 gals. of water and apply as a drench over the row at the rate of 100-250 gals./1,000 feet of row. The width of the drench treatment should be wide enough to cover the root systems of the plants.

Soil Surface Spray: Apply 6 packets of Subdue II/1,000 sq. ft. in a broadcast or banded surface spray to seedbeds, liners, or bedded stock in sufficient water to obtain uniform coverage. If applications are banded, the treated area should be wide enough to cover the root systems of the plants. Follow with a 1/2 inch irrigation.

BEST AVAILABLE COPY

Conifers in Nurseries

Subdue II provides control of Phytophthora root rot of conifers.

<u>Seedbeds and Plug-Plantings</u>	Apply 1 packet of Subdue II/1,000 sq. ft. in at least 2 gals. of water/1,000 sq. ft. in the spring and again in the fall.
<u>2-0 Transplants</u>	Apply 2 packets of Subdue II/1,000 sq. ft. in at least 2 gals. of water/1,000 sq. ft. in the spring and again in the fall.

Nonbearing Deciduous Fruits and Nuts (Nurseries and Landscape Plantings)

Subdue II provides control of Pythium root rot and Phytophthora root, crown, and collar rot of nonbearing deciduous fruits and nuts.

Apply 6-12 packets of Subdue II/1,000 sq. ft. in sufficient water to obtain thorough coverage of the soil under the canopy of the trees. Treat sufficient surface area in nurseries to cover the root zone of the plants. Use the high rate in areas known to be infested with Phytophthora. Additional applications may be made as necessary at three-month intervals during the growing season.

Note: Do not apply to trees that will bear harvestable fruit within 12 months of application, since illegal residues may result.

Vegetable Bedding Plants (Broccoli, Cabbage, Cauliflower, Eggplant, Tomatoes, Cucumbers, Lettuce, Melons, Peppers, Spinach and Squash)

Pythium Damping-Off of Broccoli, Cabbage, Cauliflower, Eggplant, Tomatoes, Cucumbers, Lettuce, Melons, Peppers, Spinach and Squash

Apply 1 packet of Subdue II/1,000 sq. ft. as a broadcast spray in sufficient water to provide uniform coverage before or at the time of seeding. Lightly incorporate or follow with 1/2 inch sprinkler irrigation water.

Note: Do not use Subdue II for disease control in vegetable plants grown in an enclosed environment, i.e., greenhouse, glasshouse, etc.

BEST AVAILABLE COPY

Storage and Disposal

Storage - Store in a cool, dry place. Do not store this product under wet conditions. When stored at temperatures less than 50°F, handle outer container carefully to avoid breakage of inner soluble packets.

Pesticide Disposal - Do not contaminate water, food or feed by storage or disposal, or cleaning of equipment. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Disposal - Do not reuse outer container. Dispose of empty outer container in a sanitary landfill or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Causes eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

Statement of Practical Treatment

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

BEST AVAILABLE COPY

If on skin: Wash with plenty of soap and water. Get medical attention.

If inhaled: Remove victim to fresh air. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Physical and Chemical Hazards

Do not use or store near heat or open flame.

Benlate® and Tersan 1991® trademark of E. I. duPont de Nemours and Co., Inc. for benomyl

Daconil 2787® trademark of ISK Biotech for chlorothalonil

Dyrene® trademark of Mobay Chemical Corporation for anilazine

D·z·n® trademark of CIBA-GEIGY

Fore® trademark of Rohm and Haas Co. for mancozeb

plantPak™ trademark of CIBA-GEIGY

Subdue® trademark of CIBA-GEIGY for metalaxyl
U.S. Patent No. 4,151,299

Tersar LSR® trademark of E. I. duPont de Nemours and Co., for maneb

©1991 CIBA-GEIGY Corporation

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419

CGA

Initial Registration

[GANNONC.LABELS]SUBDUEII - 6/25/91

This booklet manufactured using post-consumer, recycled paper.

BEST AVAILABLE COPY

LABELMASTER BOOKLET
BACK LABEL

Subdue®

Fungicide

For control of certain diseases in ornamentals, vegetable plant beds, nonbearing citrus, conifers, and nonbearing deciduous fruit and nut trees in nurseries and landscape plantings.

Active Ingredient:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester	25.0%
<u>Inert Ingredients:</u>	<u>75.0%</u>
Total:	100.0%

Subdue II is a wettable powder.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See directions for use in attached booklet.

EPA Reg. No. 100-

EPA Est.

This outer protective container contains Subdue II in eight inner water soluble packets. These inner packets dissolve in water, allowing contents to wet. To open, remove lid, then pull tab back to expose packets. Immediately dump the required number of unopened inner packets into the partially filled sprayer or mix tank. Do not handle the soluble packets or expose them to moisture, since this may cause rupturing. When not using the entire number of packets in this container, reclose the container by replacing the protective seal and lid.

Eight X One Ounce Soluble Packets
Total One Half Pound
Net Weight

Do not offer individual packets for sale.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed, inhaled or absorbed through the skin.
Causes eye irritation. Avoid contact with eyes, skin, or cloth-

BEST AVAILABLE COPY

ing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

Statement of Practical Treatment

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention.

If inhaled: Remove victim to fresh air. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Physical and Chemical Hazards

Do not use or store near heat or open flame.

Subdue® trademark of CIBA-GEIGY Corporation.

©1991 CIBA-GEIGY Corporation

Agricultural Division
CIBA-GEIGY Corporation
Greensboro, North Carolina 27419

CGA 114L1

[GANNONC.LABELS]SUBDUEII - 06/24/91