

Reg # 100-724

PM-10

192

# TORUS®

## Insect Growth Regulator For Extended Insect Control

(Insect Growth Regulator for Use in Homes, Apartments, and Other Buildings, in Non-Food Preparation Areas of Food Handling Establishments, and Outdoors Around Buildings and in Pet Areas.)

Active Ingredient: Fenoxycarb: (Ethyl [2-(4-phenoxyphenoxy) ethyl] carbamate) ..... 24.37%

Inert Ingredients\*: ..... 75.63%

Total: ..... 100.00%

\*Contains petroleum solvents

EPA Reg. No. 100-724 / EPA Est. 602-MO-1

**KEEP OUT OF REACH OF CHILDREN.**

### WARNING

See additional precautionary statements at end of label.

#### DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

#### Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. **Ciba-Geigy makes no other express or implied warranty of fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.** Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL OR ILLEGAL RESIDUES.**

#### General Information

Torus will control cockroaches, fleas and certain stored food pests in homes; apartments and non-food preparation areas of industrial, institutional and commercial buildings including but not limited to: hotels, hospitals, restaurants, stores, manufacturing plants, warehouses, transportation equipment, and ships.

Torus may also be used outdoors as a perimeter treatment around buildings to control cockroaches and in animal runs, resting areas and kennels to control fleas.

**BEST AVAILABLE COPY**

15 x 0.33 fl. oz. (15 x 10 ml)

U. S. Standard Measure



Note: Do not apply in dining areas of restaurants while food is exposed. Avoid contamination of food, feedstuffs or water supply.

Torus kills flea eggs, and treated larvae fail to emerge as adult fleas. Torus has excellent residual effect indoors, lasting six months or more depending on the material treated, frequency of cleaning and environmental conditions. Outdoor treatments should be repeated as needed.

Torus acts on the immature growth stages of cockroaches, killing young instar nymphs by disrupting molting. Torus prevents the development of last stage nymphs into fertile adults.

Torus acts on stored food pests as an ovicide and prevents the larvae from becoming adults. Torus will control: granary weevil, rice weevil, sawtoothed grain beetle, red flour beetle, confused flour beetle, lesser grain borer, cigarette beetle, Indian meal moth, angoumois grain moth, warehouse moth and mill moth.

Torus will not kill adult insects. If adult insects are present, Torus should be combined with an insecticide registered for adult control of the target insect(s). Torus complements the rapid activity of the insecticide by providing long-term control.

#### Torus System - Application Rates

Pest	Type of Application	Amount of Torus/gal. of water	Specific Directions
Cockroaches	Clean out	20-30 ml	For high infestations use 30 ml; for light infestation, 20 ml.
	Maintenance	20 ml	
	Perimeter	20 ml	
Fleas	Indoor	10 ml	Outdoors, apply 10 ml Torus to approximately 300-350 sq. ft. in sufficient water to cover.
	Outdoor	10 ml	
Stored food pests	Maintenance	20 ml	

#### Mixing Procedure

Make sure sprayer is clean before mixing. To mix Torus partially fill the sprayer with water, add the correct amount of Torus for the use specified in the table and agitate. Add the labeled rate of the insecticide selected and fill with water to the final volume. Agitate before each spray. Do not allow spray mixture to stand overnight.

#### Application Precautions

Do not allow spray mist to contact food, feedstuffs or water supplies. Dishes and food handling utensils should be thoroughly washed in soap and water if exposed to contamination by the application of the product. Do not allow children or pets to contact treated surfaces until spray has dried. **DO NOT USE IN FOOD PREPARATION AREAS OF COMMERCIAL FOOD HANDLING ESTABLISHMENTS.**

#### Torus System Application Procedure

##### Cockroaches

###### Clean Out

Apply as a crack and crevice spray and as a spot treatment with a low pressure sprayer equipped with a multi-tip nozzle. Apply to all areas that harbor cockroaches, including behind cupboards, pantries, under shelves and drawers, behind and under stoves, refrigerators and other large appliances, under sinks, baseboards, window frames, plumbing and other utility installations.

1. Initial application: Use the clean out rate. Empty all cabinets and drawers prior to application.
2. Second application: Apply the maintenance rate 6-8 weeks after the initial application, depending on the residual insecticide use.
3. Subsequent applications: Apply the maintenance rate every 3-4 months.

###### Perimeter Treatments

Apply in a band 2 to 4 feet adjacent to and around the structure and apply to a height of 2 to 3 feet on the foundation where pests may be active or may find entrance. Apply as a residual spray around window frames, doors, garage doors, patios, carports, shutters, around garbage areas and other areas where cockroaches may congregate. Avoid spraying into the air.

###### Food Serving Areas

Torus can be used where prepared foods are served, such as dining rooms, but excluding areas where food is being held or prepared.

Apply as a crack and crevice and/or spot treatment to infested areas or areas likely to be infested and between and beneath elements of construction only. Dishes and food handling utensils should be thoroughly washed in soap and

...er if exposed to contamination. Do not apply in ( ng areas of restaurants while food is exposed.

**Fleas**

Prior to treatment, carpets should be vacuumed and the vacuum cleaner bag disposed of in an outdoor trash receptacle. Pet bedding should be replaced or washed in hot water. To control the source of infestation, pets should be treated with a product registered for on-animal flea control.

**Indoor**

Apply one gallon of finished spray on approximately 1,500 sq. ft. of surface area, using a low pressure sprayer suitable for indoor applications. Treat all areas that may harbor fleas, including carpets, furniture, pet sleeping areas and throw rugs. Treat under cushions of upholstered furniture. Avoid contact with treated areas until spray has dried. Remove pets and cover aquariums prior to treatment.

**Outdoor**

To control fleas outdoors apply Torus to animal runs and areas where the animals rest. Also, if animals are kenneled, apply Torus in and around the kennel area. Apply the recommended rate to 300 to 350 square feet in sufficient water to obtain good coverage. Repeat as needed.

**Stored Food Pests**

**For Use in the Home**

Apply as a crack and crevice and a spot treatment to pantry and cupboard shelves where food containers are stored. Prior to spraying remove all food containers and food stuffs, any shelving paper and utensils. Do not replace food or containers until spray has dried. Destroy infested food packages.

**For Use in Warehouses and Commercial Food Storage Areas**

Prior to application, sweep all floors of debris and pick up all litter. Application should be made as a crack and crevice and spot treatment. Apply to all areas that may harbor pests. Apply under and between pallets.

**Storage and Disposal**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage**

Store in a locked storage area away from children. Store away from heat sources or open flame.

**Pesticide Disposal**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal**

Do not reuse container. Triple rinse container, then dispose of in a sanitary landfill or dispose of by other approved state and local procedures.

Note: Soak up spills with absorbent material and dispose of absorbent in accordance with pesticide disposal instructions. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372 day or night.

**Precautions, Statements**

**Hazards to Humans and Domestic Animals**

**WARNING**

Concentrate causes substantial but temporary eye injury. Harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing vapor or spray mist. Wear goggles, face shield or safety glasses when handling concentrate. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**Statement of Practical Treatment**

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water. Get medical attention.

If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Do not induce vomiting.

If on skin: Wash with plenty of soap and water. Get medical attention.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**Environmental Hazards**

This product is toxic to fish and aquatic invertebrates. Drift and runoff from treated areas may be hazardous to aquatic invertebrates in adjacent aquatic sites. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

**Physical or Chemical Hazards**

To avoid electrical shock or short circuits, do not apply in areas such as conduits, motor housings, junction and switch boxes, and other electrical equipment.

Do not use or store near heat or open flame.

Torus® trademark of Ciba-Geigy Corporation  
U.S. Patent No. 4,215,139

© 1993 Ciba-Geigy Corporation

Ciba-Geigy Corporation  
Animal Health Division  
Greensboro, North Carolina 27419  
CGA 1L120L1D 093

**ACCEPTED**  
18 FEB 1994  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under  
EPA Reg. No. 100-729



**BEST AVAILABLE COPY**