



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 14 1994

Theresa Kamicker
CIBA-GEIGY CORP.
P.O. Box 18300
Greensboro, NC 27419

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

**Subject: Label Amendment Submission of 07/26/93 in Response to PR Notice 93-7
EPA Reg. No. 100-736
BANNER GL FUNGICIDE**

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- **BEFORE** selling or distributing any product bearing the final printed labeling
- AND**
- **WITHIN** one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy-Cerium ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,


Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

WPS
MASTER LABEL BOOKLET
(For Update Purposes Only)

BANNER® GL

FUNGICIDE

Broad spectrum and systemic
disease control for turf and
ornamentals

ACCEPTED
with COMMENTS
by EPA Letter Dated
JAN 14 1994
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
100-736

This outer protective container holds Banner GL in 6 individual
water soluble packets which are designed to dissolve in water.
Do not handle the soluble packets excessively or expose them to
moisture, since this may cause them to rupture.

Active Ingredient:

Propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-
1,3-dioxolan-2yl]methyl]-1H-1,2,4-

triazole	41.8%
Inert Ingredients:	58.2%
Total:	100.0%

EPA Reg. No. 100-736

EPA Est. 5905-GA-01

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se
la explique a usted en detalle. (If you do not understand the
label, find someone to explain it to you in detail.)

See additional precautionary statements
and directions for use inside booklet.

CGA 131L1A 053

6 X 7 oz. Soluble Packets
Total 42 Ounces Net Weight

Ciba-Geigy

[GANNONC.LABELB] - [BNRGLWPS] - 7/14/93

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. Ciba-Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

General Information

Banner GL is a systemic fungicide for use on turfgrasses for the control of dollar spot (*Sclerotinia homoeocarpa*), brown patch (*Rhizoctonia solani*), anthracnose (*Colletotrichum graminicola*), red thread (*Corticium fuciforme*), rust (*Puccinia graminis*), powdery mildew (*Erysiphe graminis*), stripe smut (*Ustilago striiformis* and *Urocystis agropyri*), summer patch (*Magnaporthe poae*), spring dead spot (*Leptosphaeria korrea*), gray leafspot (*Pyricularia grisea*), pink snowmold (*Fusarium nivale*), gray snowmold (*Typhula* spp.), and zoysia patch.

Banner GL also controls numerous diseases on ornamentals and other landscape and nursery plantings. It controls powdery mildews, rusts, leafspots, and blights. Refer to the appropriate section for specified diseases and plants.

Note: Do not apply this product through any type of irrigation system. Do not use Banner GL as a tree injection.

Mixing Procedures

Thoroughly clean spray equipment before using this product to remove contaminants which may cause crop injury or sprayer clogging. Prepare no more spray mixture than is required for the immediate operation. Agitate thoroughly before and during application such that a rippling or rolling action is created at the liquid surface. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already-treated area.

Precautions: 1) Soluble packets of Banner GL must be dissolved completely in water before any other tank mix partner, including micronutrients or other liquid or dry fertilizer, is added to the spray solution. 2) Do not use with oils or adjuvants.

Banner GL Alone: Add 1/2 to 2/3 of the required amount of water

to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble packets of Banner GL into the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the packets have dissolved and the material has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Banner GL + Tank Mixtures: For broader spectrum control, Banner GL can be tank mixed with other fungicides. For example, Subdue® or Pace® may be tank mixed with Banner GL or used alone when conditions are favorable for Pythium blight. Banner GL is usually compatible with Chipco® 26019, Daconil 2787®, Fore®, Pace, Pennant®, Sprint® 330, Subdue 2E, Tersan® 1991 and Triumph® 4E. Do not mix Banner GL and Tersan 1991 in a fertilizer solution. Contact your dealer or a Ciba-Geigy sales representative for additional information on compatibility of other tank mix partners.

Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble packets of Banner GL into the tank. Continue agitation while adding the remainder of the water. Tank mix partners may only be added when 1) the Banner GL packets and their contents have dissolved and dispersed in the tank, and 2) when the tank has filled with water. Continue agitation when adding the tank mix partners and while spraying the solution onto the crop.

When using two or more products packaged in water-soluble packets, add the Banner GL to the tank first. Allow the packet to completely dissolve and the product to completely disperse into the mix water. Then add the other product packaged in a water-soluble packet and allow it to fully dissolve and disperse. Maintain rolling agitation throughout the entire mixing and spraying process.

If using Banner GL in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions and limitations which appear on the tank mix partner label. No label dosage rate should be exceeded and the most restrictive label precautions and limitations should be followed. This product should not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Turfgrass and Dichondra Disease Control

1. USE BANNER GL IN A PREVENTATIVE DISEASE CONTROL PROGRAM.
2. Apply in sufficient water to ensure thorough coverage.
3. Apply after mowing OR allow sprayed area to completely dry before mowing.

4. For control of foliar diseases, allow sprayed area to completely dry before irrigation.
5. For control of soil-borne diseases, Banner GL can be watered in after application.
6. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
7. For optimum turf quality and disease control, use Banner GL in conjunction with turf management practices that promote good plant health and optimum disease control.
8. Addition of surfactants may result in phytotoxicity.
9. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

CAUTION: Bermudagrass can be sensitive to Banner GL. Do not apply more than 1 packet per 5,000 sq. ft. every 30 days on any variety of bermudagrass. In Florida, do not apply Banner GL to Bermudagrass golf course greens when temperatures exceed 90°F.

NOTE: (1) To avoid possible illegal residues, do not graze animals on treated areas and (2) do not feed clippings from treated areas to livestock or poultry.

Turfgrass - Specific Diseases, Rates and Application Timing

<u>Disease</u>	<u>Area to be Treated per Packet (sq. ft.)</u>	<u>Application Interval/ Timing</u>	<u>Instructions</u>
Dollar Spot	40,000	7-14 days	Apply when conditions are favorable for disease development. If the 40,000 sq. ft. rate is used, tank mix with the lowest label rate of another EPA-registered fungicide having a different mode of action. If using the 10,000-20,000 sq. ft. rate, make no more than 3 consecutive applications for dollar spot control before rotating to an alternate EPA-registered fungicide having a different mode of action.
	10,000-20,000	14-28 days	

<u>Disease</u>	<u>Area to be Treated per Packet (sq. ft.)</u>	<u>Application Interval/ Timing</u>	<u>Instructions</u>
Brown Patch	5-10,000	10-21 days	Begin applications in May or June before disease is present. If disease is present, mix 1 packet Banner GL per 10,000 sq. ft. with a registered contact fungicide at the labeled rate. Under conditions of high temperature and high humidity, use the higher rates and shorter intervals.
Powdery Mildew, Rust, thracnose	10-20,000	14-28 days	Apply when conditions are favorable for An-disease development.
Red Thread	10,000	14-21 days	Apply when conditions are favorable for disease development.

<u>Disease</u>	<u>Area to be Treated per Packet (sq. ft.)</u>	<u>Application Interval/ Timing</u>	<u>Instructions</u>
Stripe Smut	10-20,000	Fall or Spring	Apply once in the fall after grass becomes dormant or in the early spring before grass starts to grow.
Gray Leafspot (St. Augustinegrass sod production)	10,000	14 days	Apply when conditions are favorable for disease development.
Melting Out, Leaf Spot	5-10,000	14 days	Under light to moderate pressure, apply Banner GL to reduce the severity of leaf spot and melting out caused by Helminthosporium type pathogens.
Summer Patch, Poa Patch	5,000	30 days	Apply Banner GL at monthly intervals beginning in April. Make 2-3 applications depending on local recommendations.
Spring Dead Spot (California Only)	5,000	30 days	Apply Banner GL at monthly intervals beginning in August. Make three applications.

<u>Disease</u>	<u>Area to be Treated per Packet (sq. ft.)</u>	<u>Application Interval/ Timing</u>	<u>Instructions</u>
Snowmold			
Gray - <u>Typhula</u> spp.	5,000	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow.
Pink - <u>Fusarium</u> <u>nivale</u>			
Zoysia patch, large patch of zoysia	3,500- 5,000	Early Fall	Make one application in the early fall (mid-September to mid-October) prior to development of disease symptoms. Consult local turfgrass extension experts to determine the optimum application timing for your area.

Dichondra - Specific Diseases, Rate and Application Timing

<u>Disease</u>	<u>Area to be Treated per Packet (sq. ft.)</u>	<u>Application Interval/ Timing</u>	<u>Instructions</u>
Dichondra Rust	10,000	14-21 days	Apply when conditions are favorable for disease development.

Establishment of Cool Season Turfgrass

When used in a preventative disease control program, Banner GL applied to cool season grass seedlings or sod will improve the rate of establishment.

New Seedlings: Apply 1 packet/20,000 sq. ft. at the two to three leaf stage of growth for faster root development and top growth.

Sod: Apply 1 packet/20,000 sq. ft. 2-6 weeks before cutting for increased sod knitting and rate of establishment after resetting.

Disease Control in Nurseries (Field) and Landscape Plantings

1. USE BANNER GL IN A PREVENTATIVE DISEASE CONTROL PROGRAM. INITIATE APPLICATIONS ACCORDING TO LOCAL RECOMMENDATIONS OR AT FIRST SIGNS OF DISEASE.
2. Apply in enough water to thoroughly cover foliage to the point of runoff.
3. Allow spray to dry before overhead irrigation is applied.
4. Optimum benefit of Banner GL is obtained when used in conjunction with sound disease management practices.

Note: Banner GL has been shown to be safe on the species of plants and effective in controlling the diseases listed in the Directions For Use. Other plant species may be sensitive to Banner GL and other diseases may not be controlled. Before using Banner GL on plants or for diseases that are not listed in the Directions For Use, test Banner GL on a small scale basis first. Apply the recommended rates for a particular type of disease, i.e., rust, powdery mildew, etc., and evaluate for phytotoxicity and disease control prior to widespread use. Do not apply Banner GL to African violets, begonias, Boston fern, or geraniums.

Woody Plants

<u>Species</u>	<u>Disease</u>	<u>Recommended Rates and Timing</u>
Crape Myrtle	Powdery Mildew (<u>E. lagerstroemia</u>)	Apply 1 packet per 130 gals. every 21 days.
Douglas Fir	Needle Rust (<u>Melampsora occidentalis</u>)	Apply 1 packet per 130 gals. Apply once in May.
Jeffrey Pine	Tip blight (<u>Sirococcus strobolinus</u>)	Apply 1 packet per 130 gals. Begin applications in mid-late winter (Nov.-Jan.) Apply 3 times at 2-month intervals.
Oaks	Leafspot (<u>Tubakia dryina</u>)	Apply 1 packet per 130 gals. every 14 days.
Red Tip Photinia	Leafspot (<u>Fabraea maculata</u> syn. <u>Entomosporium maculatum</u>)	Apply 1 packet per 100-150 gals. every 7-14 days. Reductions of plant height may occur with Banner GL applications, especially on plants 6 inches or smaller.
Shasta Fir	Needle Rust (<u>Pucciniastrum goeppertianum</u>)	Apply 1 packet per 130 gals. in May-June prior to anticipated infection periods.

Non-Bearing Fruits and Nuts
(Nurseries and Landscape Plantings)

<u>Species</u>	<u>Disease</u>	<u>Recommended Rates and Timing</u>
Cherry	Powdery Mildew Leafspot (<u>Coccomyces hiemalis</u>)	Apply 1 packet per 1.5-3.5 acres on a 14-21 day interval. Apply 1 packet per 1.5 acres on a 14-21 day interval.
	Blossom Blight (<u>Monilinia fructicola</u>)	Apply 1 packet per 1.5-2.25 acres. Apply at 5-10% bloom and again at 70-100% bloom.
Citrus	Greasy Spot (<u>Mycosphaerella citri</u>)	Apply 1 packet per 100 gals. during June-August.
Peach Plum Nectarine	Blossom Blight (<u>M. fructicola</u> or <u>M. laxa</u>)	Apply 1 packet per 1.5-2.25 acres. Apply at 5-10% bloom and again at 70-100% bloom.
Pecan	Scab (<u>Cladosporium</u>), Powdery Mildew, Downy Spot, Zonate Leafspot, Fungal Leaf Scorch, Brown Spot	Apply 1 packet per 3.75-5 acres beginning at bud break. Apply 3 times on a 14-day interval.

Note: To avoid possible illegal residues, do not apply to cherry, citrus, peach, plum, nectarine, or pecan trees that will bear harvestable fruit within 12 months of application.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage

Store in a cool, dry place. Do not store this product under wet conditions. Handle outer packaging carefully to avoid breakage of inner soluble packets.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instruc-

tions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Disposal

Do not reuse outer container. Dispose of empty outer container in a sanitary landfill, or by incineration, or by burning if allowed by state and local authorities. Stay out of smoke from burning containers.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372 day or night.

Precautionary Statements**Hazards to Humans and Domestic Animals****WARNING/AVISO**

Causes substantial but temporary eye injury. Do not get in eyes. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin or clothing. Avoid breathing vapors or spray mist.

Statement of Practical Treatment

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water. Get medical attention.

If swallowed: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, large quantities of water. Never give anything by mouth to an unconscious person. Call a physician or Poison Control Center.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth and get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

Notes to Physician: If ingested, lavage stomach to avoid aspiration. Give a saline laxative and supportive therapy.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Physical or Chemical Hazards

Do not use or store near heat, open flame, or hot surfaces.

Banner® trademark of Ciba-Geigy
U.S. Patent No. 4,079,062

Chipco® 26019 trademark of Rhône-Poulenc, Inc.

Daconil 2787® trademark of ISK Biotech Corp.

Fore® trademark of Rohm & Haas Co.

Pace®, Pennant®, Sprint®, Subdue® and Triumph® trademarks of Ciba-Geigy Corporation

Tersan® 1991 trademark of E. I. duPont de Nemours and Co., Inc.

Unite® trademark of Hopkins Chemical Co.

©1993 Ciba-Geigy Corporation

Ciba Plant Protection
Turf and Ornamental Products
Ciba-Geigy Corporation
Greensboro, North Carolina 27419

6 X 7 oz. Soluble Packets
Total 42 Ounces Net Weight

*This booklet manufactured using post-consumer, recycled paper
and soy-based inks.*

[GANNONC.LABELB]BNRGLWPS - 7/14/93

Booklet Back Panel

BANNER® GL

FUNGICIDE

Broad spectrum and systemic disease control for turf and ornamentals.

Active Ingredient:

Propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2yl]methyl]-1H-1,2,4-triazole	41.8%
<u>Inert Ingredients:</u>	<u>58.2%</u>
Total:	100.0%

See directions for use in attached booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-736
EPA Est. 5905-GA-01

Banner® GL trademark of Ciba-Geigy
U.S. Patent No. 4,079,062

©1992 Ciba-Geigy Corporation

6 X 7 oz. Soluble Packets
Total 42 Ounces Net Weight.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes substantial but temporary eye injury. Do not get in eyes. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin or clothing. Avoid breathing vapors or spray mist.

Statement of Practical Treatment

- If in eyes: Hold eyelids open and flush with a steady, gentle stream of water. Get medical attention.
- If swallowed: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution or, if these are not available, large quantities of water. Never give anything by mouth to an unconscious person. Call a physician or Poison Control Center.
- If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth and get medical attention.
- If on skin: Wash with plenty of soap and water. Get medical attention.

Notes to Physician: If ingested, lavage stomach to avoid aspiration. Give a saline laxative and supportive therapy.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Physical or Chemical Hazards

Do not use or store near heat, open flame, or hot surfaces. In the event of spill, fire, or other emergency, call 1-800-888-8372 day or night.

Storage

Store in a cool, dry place. Do not store this product under wet conditions. Handle outer packaging carefully to avoid breakage of inner soluble packets.

Chemigation

Do not apply this product through any type of irrigation system.

Ciba Plant Protection
Turf and Ornamental Products
Ciba-Geigy Corporation
Greensboro, North Carolina 27419

CGA 131L1A 053

Product ID: 59890

Ciba-Geigy

[GANNONC.LABELB]BNRGLWPS - 7/14/93