

Banner® GL

Fungicide

Broad spectrum and systemic disease control for turf and ornamentals

Active Ingredient:	
Propiconazole: (CAS No. 60207-90-1).....	41.8%
Other Ingredients:	58.2%
Total:	100.0%

Banner GL contains a nominal 3.6 pounds of active ingredient per gallon.

This outer protective container holds Banner GL in 6 individual water-soluble packets which are designed to dissolve in water. Do not handle the soluble packets excessively or expose them to moisture, since this may cause them to rupture.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-736

EPA Est. 5905-GA-01


SCP 736A-M2

6 X 7.2 ounces

Water Soluble Packets

43.2 ounces

Total Net Contents



FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN	
If ingested, induce emesis or lavage stomach. Treat symptomatically.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER	
For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

Causes substantial, but temporary eye injury. Do not get in eyes. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin or clothing. Avoid breathing vapors or spray mist.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Physical and Chemical Hazards

Do not use or store near heat, open flame, or hot surfaces.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and of Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

6 8 21

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage

Store in a cool, dry place. **Do not store below 40°F.** Do not store this product under wet conditions. Handle outer packaging carefully to avoid breakage of inner soluble packets.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Disposal

Do not reuse outer container. Dispose of empty outer container in a sanitary landfill, or by incineration, or by burning, if allowed by state and local authorities. Stay out of smoke from burning containers.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

GENERAL INFORMATION

Banner GL is a systemic fungicide for use on turfgrasses for the control of dollar spot (*Sclerotinia homoeocarpa*), brown patch (*Rhizoctonia solani*), anthracnose (*Colletotrichum graminicola*), red thread (*Corticium fuciforme*), rust (*Puccinia graminis*), powdery mildew (*Erysiphe graminis*), stripe smut (*Ustilago striiformis* and *Urocystis agropyri*), summer patch (*Magnaporthe poae*), spring dead spot (*Leptosphaeria korrea*), gray leafspot (*Pyricularia grisea*), pink snowmold (*Fusarium nivale*), gray snowmold (*Typhula* spp.), and zoysia patch.

Banner GL also controls numerous diseases on ornamentals and other landscape and nursery plantings. It controls powdery mildews, rusts, leafspots, and blights. Refer to the appropriate section for specified diseases and plants.

Note: Do not apply more than 37 packets of Banner GL/acre/year.

Note: Do not apply this product through any type of irrigation system. Do not use Banner GL as a tree injection.

Mixing Procedures

Thoroughly clean spray equipment before using this product to remove contaminants which may cause crop injury or sprayer clogging. Prepare no more spray mixture than is required for the immediate operation. Agitate thoroughly before and during application, such that a rippling or rolling action is created at the liquid surface. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already-treated area.

Precautions: (1) Soluble packets of Banner GL must be dissolved completely in water before any other tank mix partner, including micronutrients, or other liquid or dry fertilizer, is added to the spray solution. (2) Do not use with oils or adjuvants.

Banner GL Alone: Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble packets of Banner GL into the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the packets have dissolved and the material has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Banner GL + Tank Mixtures: For broader spectrum control, Banner GL can be tank mixed with other fungicides. For example, Subdue® may be tank mixed with Banner GL, or used alone when conditions are favorable for Pythium blight. Banner GL is usually compatible with Chipco® 26019, Daconil 2787®, Fore®, and Pennant®. Contact your dealer or a Syngenta sales representative for additional information on compatibility of other tank mix partners.

Add $\frac{1}{2}$ – $\frac{2}{3}$ of the required amount of water to the spray or mixing tank. With the agitator running, drop the required number of unopened soluble packets of Banner GL into the tank. Continue agitation while adding the remainder of the water. Tank mix partners may only be added when 1) the Banner GL packets and their contents have dissolved and dispersed in the tank, and 2) when the tank has filled with water. Continue agitation when adding the tank mix partners and while spraying the solution onto the crop.

When using two or more products packaged in water-soluble packets, add the Banner GL to the tank first. Allow the packet to completely dissolve and the product to completely disperse into the mix water. Then add the other product packaged in a water-soluble packet and allow it to fully dissolve and disperse. Maintain rolling agitation throughout the entire mixing and spraying process.

If using Banner GL in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix partner label. No label dosage rate should be exceeded and the most restrictive label precautions and limitations should be followed. This product should not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

TURFGRASS AND DICHONDRA DISEASE CONTROL

1. USE BANNER GL IN A PREVENTATIVE DISEASE CONTROL PROGRAM.
2. Apply in sufficient water to ensure thorough coverage.
3. Apply after mowing OR allow sprayed area to completely dry before mowing.
4. For control of foliar diseases, allow sprayed area to completely dry before irrigation.
5. For control of soil-borne diseases, Banner GL can be watered in after application.
6. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
7. For optimum turf quality and disease control, use Banner GL in conjunction with turf management practices that promote good plant health and optimum disease control.
8. Addition of surfactants may result in phytotoxicity.

10 8 21

9. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.
10. Do not apply more than 37 packets/acre.

PRECAUTION: Bermudagrass can be sensitive to Banner GL. Do not apply more than 8 packets/acre every 30 days on any variety of bermudagrass. In FL, do not apply Banner GL to bermudagrass golf course greens when temperatures exceed 90°F.

NOTES: (1) To avoid possible illegal residues, do not graze animals on treated areas, and (2) do not feed clippings from treated areas to livestock or poultry.

Turfgrass - Specific Diseases, Rates, and Application Timing

Disease	Area to be Treated per Packet (sq. ft.)	Application Interval/ Timing	Instructions
Dollar Spot	43,560 (1 acre)	7-14 days	Apply when conditions are favorable for disease development. If the 43,560 sq. ft. (1 acre) rate is used, tank mix with the lowest label rate of another EPA-registered fungicide having a different mode of action. If using the 10,890-21,780 sq. ft. (0.25-0.5 acre) rate, make no more than 3 consecutive applications for dollar spot control before rotating to an alternate EPA-registered fungicide having a different mode of action.
	10,890-21,780 (0.25-0.50 acre)	14-28 days	

Turfgrass - Specific Diseases, Rates, and Application Timing

Disease	Area to be Treated per Packet (sq. ft.)	Application Interval/ Timing	Instructions
Brown Patch	10,890-21,780 (0.25-0.50 acre)	14-21 days	Begin applications in May or June before disease is present. Tank mix 1 packet Banner GL per 10,890-21,780 ft. ² (0.25-0.5 acre) with a registered contact fungicide at the labeled rate. Under conditions of high temperature and high humidity, use the higher rates and shorter intervals.
Powdery Mildew Rust Anthracnose	10,890-21,780 (0.25-0.50 acre)	14-28 days	Apply when conditions are favorable for disease development.
Red Thread	10,890 (0.25 acre)	14-21 days	Apply when conditions are favorable for disease development.

Turfgrass - Specific Diseases, Rates, and Application Timing

Disease	Area to be Treated per Packet (sq. ft.)	Application Interval/ Timing	Instructions
Stripe Smut	10,890-21,780 (0.25-0.50 acre)	Fall or Spring	Apply once in the fall after grass becomes dormant or in the early spring before grass starts to grow.
Gray Leafspot (St. Augustine-grass sod production)	10,890-21,780 (0.25-0.50 acre)	14-21 days	Apply when conditions are favorable for disease development. Tank mix with a registered contact fungicide at label rate.
Melting Out Leaf Spot	10,890-21,780 (0.25-0.50 acre)	14 days	Under light to moderate pressure, apply Banner GL to reduce the severity of leaf spot and melting out caused by Helminthosporium type pathogens.
Summer Patch Poa Patch	5,445 (0.125 acre)	28 days	Apply Banner GL at monthly intervals beginning in April.
	10,890 (0.25 acre)	14 days	Make 2-3 applications depending on local recommendations. For optimum control, tank mix with contact fungicide at label rates.
Spring Dead Spot (California Only)	5,445 (0.125 acre)	30 days	Apply Banner GL at monthly intervals beginning in August. Make 3 applications.

Turfgrass - Specific Diseases, Rates, and Application Timing

Disease	Area to be Treated per Packet (sq. ft.)	Application Interval/ Timing	Instructions
Snowmold Gray - <i>Typhula</i> spp. Pink - <i>Fusarium</i> <i>nivale</i>	5,445 (0.125 acre)	Late Fall	Apply one application in the late fall before snow cover. Do not apply on top of snow.
Zoysia patch large patch of zoysia	5,445 (0.125 acre)	Early Fall	Make one application in the early fall (mid-September to mid-October) prior to development of disease symptoms. Consult local turfgrass extension experts to determine the optimum application timing for your area.

Dichondra - Specific Disease, Rate, and Application Timing

Disease	Area to be Treated per Packet (sq. ft.)	Application Interval/ Timing	Instructions
Dichondra Rust	10,890 (0.25 acre)	14-21 days	Apply when conditions are favorable for disease development

Establishment of Cool Season Turfgrass

When used in a preventative disease control program, Banner GL applied to cool season grass seedlings or sod will improve the rate of establishment.

New Seedlings: Apply 1 packet/21,780 sq. ft. (0.5 acre) at the 2 to 3-leaf stage of growth for faster root development and top growth.

Sod: Apply 1 packet/21,780 sq. ft. (0.5 acre) 2-6 weeks before cutting for increased sod knitting and rate of establishment after resetting.

DISEASE CONTROL IN NURSERIES (FIELD) AND LANDSCAPE PLANTINGS

1. USE BANNER GL IN A PREVENTATIVE DISEASE CONTROL PROGRAM. INITIATE APPLICATIONS ACCORDING TO LOCAL RECOMMENDATIONS OR AT FIRST SIGNS OF DISEASE.
2. Apply in enough water to thoroughly cover foliage to the point of runoff.
3. Allow spray to dry before overhead irrigation is applied.
4. Optimum benefit of Banner GL is obtained when used in conjunction with sound disease management practices.
5. You can apply up to 37 packets of Banner GL/A/crop.

NOTE: Banner GL has been shown to be safe on the species of plants and effective in controlling the diseases listed in the **Directions for Use**. Other plant species may be sensitive to Banner GL and other diseases may not be controlled. Before using Banner GL on plants or for diseases that are not listed in the **Directions for Use**, test Banner GL on a small scale basis first. Apply the recommended rates for a particular type of disease, i.e., rust, powdery mildew, etc., and evaluate for phytotoxicity and disease control prior to widespread use. **Do not apply Banner GL to African violets, begonias, Boston fern, or geraniums.**

Woody Plants

Species	Disease	Recommended Rates and Timing
Crape Myrtle	Powdery Mildew (<i>E. lagerstroemia</i>)	Apply 1 packet/135 gals. every 21 days.
Douglas Fir	Needle Rust (<i>Melampsora occidentalis</i>)	Apply 1 packet/135 gals. Apply once in May.
Jeffrey Pine	Tip blight (<i>Sirococcus strobolinus</i>)	Apply 1 packet/135 gals. Begin applications in mid-late winter (Nov.-Jan.). Apply 3 times at 2-month intervals.
Oaks	Leafspot (<i>Tubakia dryina</i>)	Apply 1 packet/135 gals. every 14 days.
Red Tip Photinia	Leafspot (<i>Fabraea maculata</i> syn. <i>Entomosporium maculatum</i>)	Apply 1 packet/110-180 gals. every 7-14 days. Reductions of plant height may occur with Banner GL applications, especially on plants 6 inches or smaller.
Shasta Fir	Needle Rust (<i>Pucciniastrum goeppertianum</i>)	Apply 1 packet/135 gals. in May-June, prior to anticipated infection periods.

Non-Bearing Fruits and Nuts (Nurseries and Landscape Plantings)

Species	Disease	Recommended Rates and Timing
Cherry	Powdery Mildew Leafspot (<i>Coccomyces hiemalis</i>)	Apply 1 packet/1.75-3.75 acres on a 14 to 21-day interval. Apply 1 packet/1.5 acres on a 14 to 21-day interval.
	Blossom Blight (<i>Monilinia fructicola</i>)	Apply 1 packet/1.75-2.5 acres. Apply at 5-10% bloom and again at 70-100% bloom.
Citrus	Greasy Spot (<i>Mycosphaerella citri</i>)	Apply 1 packet/100 gals. during June- August.
Peach Plum Nectarine	Blossom Blight (<i>M. fructicola</i> or <i>M. laxa</i>)	Apply 1 packet/1.75-2.5 acres. Apply at 5-10% bloom and again at 70-100% bloom.
Pecan	Scab (<i>Cladosporium</i>) Powdery Mildew Downy Spot Zonate Leafspot Fungal Leaf Scorch Brown Spot	Apply 1 packet/1.25-1.5 acres beginning at bud break. Apply 3 times on a 14-day interval.

Note: To avoid possible illegal residues, do not apply to cherry, citrus, peach, plum, nectarine, or pecan trees that will bear harvestable fruit within 12 months of application.

Banner®, Daconil 2787®, Pennant®, Subdue and the Syngenta logo are trademarks of a Syngenta Group Company
 Chipco® 26019 trademark of Aventis CropScience
 Fore® trademark of Rohm and Haas Company
 Unite® trademark of HACO, Inc.

©200X Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

6 X 7.2 ounces
Water-Soluble Packets

43.2 ounces
Total Net Contents

SCP 736A-M2

BAN GL SCP 736A-M2-AMD DRAFT2 nsp 08-13-04

Booklet Base

Banner® GL

Fungicide

Broad spectrum and systemic disease control for turf and ornamentals

Active Ingredient:	
Propiconazole: (CAS No. 60207-90-1).....	41.8%
Other Ingredients:	58.2%
Total:	100.0%

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See directions for use in attached booklet.

<p>AGRICULTURAL USE REQUIREMENTS</p> <p>Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.</p>

EPA Reg. No. 100-736

EPA Est. 5905-GA-01

6 X 7.2 ounces
Water-Soluble Packets

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
NOTE TO PHYSICIAN	
If ingested, induce emesis or lavage stomach. Treat symptomatically.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372	

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes substantial, but temporary eye injury. Do not get in eyes. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin or clothing. Avoid breathing vapors or spray mist.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Physical and Chemical Hazards

Do not use or store near heat, open flame, or hot surfaces. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Storage

Store in a cool, dry place. **Do not store below 40°F.** Do not store this product under wet conditions. Handle outer packaging carefully to avoid breakage of inner soluble packets.

Chemigation

Do not apply this product through any type of irrigation system.

Banner® and the Syngenta logo are trademarks of a Syngenta Group Company

©200X Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 736A-M2

BAN GL SCP 736A-M2-AMD DRAFT2 nsp 08-13-04
000100-00736.20040818.Banner GL Amend.doc
000100-00736.20040818.Banner GL Amend.pdf