

JUN 4 1994

Dr. Gregory R. Watson
 Ciba Plant Protection
 Ciba-Geigy Corporation
 P.O. Box 18300
 Greensboro, NC 27419-8300

Dear Dr. Watson:

Subject: Label Amendment - Microinjector Container
 Alamo Fungicide
 EPA Registration Number 100-741
 Your Submission Dated February 28, 1994

The amendment referred to above, which revised the directions for use in microinjector container, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration or reregistration of your product under FIFRA sec. 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the following labeling changes;
 - a) On page 3 of the proposed label in the first sentence of the second full paragraph, change "Alamo: water solution" to "Alamo" since there appears to be no additional water used for this treatment.
 - b) Delete the chemigation statement. This statement is not required for products solely intended for injection into plants.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subjected to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

CONCURRENCES

SYMBOL	7505C						
SURNAME	Robbins						
DATE	6/14/94						

20712

[2]

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Steve Robbins *SR*
Acting Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

30412
Highlighted
copy

Microinjection Unit - 1994
Master Label Booklet

ALAMO®
FUNGICIDE

A trunk-injected systemic fungicide for control of certain wilt diseases in oaks and elms.

Active Ingredient:

Propiconazole:

1-[[2-(2,4-Dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]-methyl]-1H-1,2,4-triazole 14.3%

Inert Ingredients: 85.7%

Total: 100.0%

EPA Reg. No. 100-741

EPA Est. 64014-NY-1

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

PRECAUCIÓN AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

See additional precautionary statements and directions for use inside booklet.

10 mls.

U. S. Standard Measure

ACCEPTED
with COMMENTS
In EPA Letter Dated

JUN 4 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

100-741

CGA 136L2

[GANNONC.LABELA]ALAMOF42 - 2/23/94

BEST AVAILABLE COPY

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. Ciba-Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

General Information

Alamo is a systemic fungicide for use as a root flare injection for prevention and treatment of (1) oak wilt (Ceratocystis fagacearum) of oak (Quercus spp.), and (2) Dutch elm disease (Ceratocystis ulmi) of elms (Ulmus spp). These fungi infect the vascular system and cause plugging throughout the tree. It is recommended that Alamo be administered by trained arborists or others trained in injection techniques and in the identification

BEST AVAILABLE COPY

of oak wilt and Dutch elm disease. Alamo should be injected into flare roots to ensure distribution throughout the vascular system of the tree.

Important: Oak trees exhibiting less than 20% crown loss from oak wilt will have the best chance of responding to treatment by Alamo. Preventive application is more effective than therapeutic treatment into trees showing disease symptoms. Trees in advanced stages of disease development may not respond to treatment.

Uninfected trees will generally absorb the full amount of Alamo:water solution within 1 hour. Trees exhibiting specific symptoms or symptomless trees immediately adjacent to a diseased tree should be considered infected. Symptomless trees separated by a primary plow line from diseased trees may be at less risk of infection. Infected trees will absorb the material more slowly due to the vascular plugging caused by the disease. If Alamo is not absorbed within 24 hours, the tree is considered high risk and has a poor chance of survival.

Correct Location for Microinjection Placement

The root flare area is the transitional zone between the trunk and the root system. Uptake and distribution of Alamo is more effective when injections are made into the flare roots. In addition, wounds created in the root flare area close more rapidly in comparison to wounds above the root flare area.

Tree Measurement

Measure the diameter of the tree using a tree diameter tape (D tape) at 4 1/2 ft. above the ground. This is the diameter at breast height (DBH). Use the DBH measurement to determine the number of microinjection units to inject into each tree.

Tree Preparation

1. Heavy, thick, or loose outer bark may be carefully shaved to form a smoother injection point and to ensure the operator that the drill hole penetrates through the bark to the xylem.
2. If root flares are not clearly exposed, carefully remove 2-4 inches of soil from the base of the tree to uncover the top of the root flares. Brush away loose soil.
3. Do not space microinjection units further than 5 inches apart. A slight downward angle hole is recommended for more complete drainage of the microinjection unit.

BEST AVAILABLE COPY

Microinjection Procedure

1. Using an electric drill at 600 to 800 rpm, with a sharp, clean 11/64 inch (0.4 cm) drill bit, the installer should drill evenly spaced holes to a depth of 3/8-1/2 inch (0.90-1.3 cm) through the bark into the xylem.
2. After reaching the proper depth range, the drill bit should be withdrawn carefully to avoid dislodging bark fragments around the exterior opening of the hole.
3. Disinfect the drill bit between trees with Lysol or a 20% solution of household bleach. Rinse bit with clean water after treatment with Lysol or the bleach solution.
4. Each hole should be drilled and a microinjection unit installed as soon as possible after the hole is drilled. A small amount of water squirted into the drill hole will help form a tight seal.
5. While wearing the appropriate protective clothing and eye wear, manually insert the dispenser tube portion of the microinjection unit in the hole.
6. Placing the plastic installation cap over the rear barrel end, strike the cap with a plastic hammer to seat the microinjection unit firmly in the hole.
7. When the microinjection unit is positioned correctly in the tree, remove the cap and push the rear barrel portion of the unit downward until it is flush with the edge of the locking mechanism. This pressurizes the microinjection unit and assists in the movement of Alamo into the vascular system of the tree.
8. When properly installed, the microinjection unit generates internal pressure resulting in the flow of Alamo solution through the dispenser tube. The microinjection unit should never be activated unless installed correctly and securely in the tree to be treated.
9. Microinjection units containing Alamo may require up to several minutes or more to empty depending on the health of the treated tree and local weather conditions. Microinjection units should be removed when the evacuation of Alamo is completed. Never assume that microinjection units have depressurized completely because they appear empty.
10. When removing the microinjection unit, wear eye protection and the gloves recommended in the Precautionary Statements portion of this label. Cover the microinjection unit with one hand near the point of insertion into the root flare while grasping the barrel end on the microinjection unit with

BEST AVAILABLE COPY

the other hand. The microinjection unit should be twisted slightly as it is slowly withdrawn from the tree.

11. After units are removed from the trees, they must be discarded into the heavy-duty plastic disposal bag included in each case of injector units. The bag should be properly sealed and placed in the original carton.
12. Sealed cartons should be returned freight prepaid for disposal to: Tree Technology System, 1014 Rein Road, Cheektowaga, NY 14225.

Oak Wilt of Oaks, Dutch Elm Disease of Elms

For treatment of oak wilt and Dutch elm disease, use one microinjector unit of Alamo per each inch DBH.

Retreatment

At the initial injection of Alamo, take notes on the level of disease in each tree. Reevaluate disease level in trees at 12 month intervals after treatment for the potential need for retreatment with Alamo. After this 12-month period, continue to evaluate trees for the need for retreatment on an annual basis. Preventive applications should be considered 24 months after the initial injection. Trees in high disease risk areas or high value trees should be evaluated for possible retreatment 12 months after each treatment.

Follow application procedures described above for repeat injections; new drill holes will be needed for subsequent treatments.

Notes: (1) Accurate diagnosis of oak wilt and Dutch elm disease is important since Alamo only provides control of the diseases listed on this label. (2) Alamo will be most effective when used in conjunction with other cultural practices recommended for management of oak wilt and Dutch elm disease (removal of dead elm trees, pruning of diseased tree limbs and branches, control of bark beetles, etc.). For further information on the proper diagnosis and control of oak wilt and Dutch elm disease, consult your local extension agent.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of unused pesticide is a violation of federal law. If these wastes cannot be used according to label instructions, contact your local

BEST AVAILABLE COPY

X0712

State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Disposal

Do not reuse empty microinjector container. Used microinjector units should be placed in the heavy-duty plastic bag which accompanies each case of injector units. The bag must be properly sealed, placed into the original shipping carton, and returned freight prepaid for disposal to: Tree Technology Systems, Inc., 1014 Rein Road, Cheektowaga, NY 14225.

For minor spills, leaks, etc., follow all precautions listed on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

BEST AVAILABLE COPY

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING/AVISO

Causes substantial, but temporary eye injury. Wear goggles or face shield. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapor. Wear rubber gloves and a long-sleeved shirt when mixing, handling, and applying the product. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

Statement of Practical Treatment

If in eyes: Immediately flush eyes with a steady, gentle stream of water. Get medical attention.

If on skin: Wash thoroughly with soap and water. Get medical attention if irritation occurs.

If swallowed: Do not induce vomiting. Drink plenty of water and contact a physician, hospital, or local Poison Control Center.

If inhaled: Move victim to fresh air.

Notes to Physician: If ingested, lavage stomach to avoid aspiration. A slurry of activated charcoal in water can be left in the stomach. Give a saline laxative and supportive therapy.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

Chemigation

Do not apply this product through any type of irrigation system.

BEST AVAILABLE COPY

108712

Alamo® trademark of Ciba-Geigy Corporation

U.S. Patent No. 4,079,062

©1994 Ciba-Geigy Corporation

Ciba Plant Protection
Turf and Ornamental Products
Ciba-Geigy Corporation
Greensboro, NC 27419

CGA 136L2

Product ID:

) *This booklet manufactured using post-consumer, recycled paper.*

)
BEST AVAILABLE COPY

Booklet Back Panel

ALAMO®

FUNGICIDE

A trunk-injected systemic fungicide for control of certain wilt diseases in oaks and American elm.

Active Ingredient:

Propiconazole:

1-[[2-(2,4-Dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]-methyl]-1H-1,2,4-triazole 14.3%

Inert Ingredients: 85.7%

Total: 100.0%

See directions for use in attached booklet.

EPA Reg. No. 100-741

EPA Est. 64014-NY-1

Alamo® trademark of Ciba-Geigy Corporation

U.S. Patent No. 4,079,062

©1994 Ciba-Geigy Corporation

10 mls.

U.S. Standard Measure

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

PRECAUCIÓN AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes substantial, but temporary eye injury. Wear goggles or face shield. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapor. Wear rubber gloves and a long-sleeved shirt when mixing, handling, and applying the product. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

BEST AVAILABLE COPY

12/12

Statement of Practical Treatment

- If in eyes: Immediately flush eyes with a steady, gentle stream of water. Get medical attention.
- If on skin: Wash thoroughly with soap and water. Get medical attention if irritation occurs.
- If swallowed: Do not induce vomiting. Drink plenty of water and contact a physician, hospital, or local Poison Control Center.
- If inhaled: Move victim to fresh air.

Notes to Physician: If ingested, lavage stomach to avoid aspiration. A slurry of activated charcoal in water can be left in the stomach. Give a saline laxative and supportive therapy.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Physical or Chemical Hazards

Do not use or store near heat or open flame. In the event of a spill, fire, or other emergency, call 1-800-888-8372, day or night.

Chemigation

Do not apply this product through any type of irrigation system.

Ciba Plant Protection
Turf and Ornamental Products
Ciba-Geigy Corporation
Greensboro, NC 27419

CGA 136L2

Product ID:

[GANNONC.LABELA]ALAMOF42 - 2/23/94

BEST AVAILABLE COPY