

GROWTH REGULATOR For control of selected insects, including whitefiles, soft scales, fungue gnats, and shore files in landscape ornamentals, nurseries, and greenhouse ornamental crops

ve ingredient: Fencinycarb: (Ethyl (2-(4-phenoxyphenoxy) ethyl carbernaie)

inen Ingredients:

100%

75%

Ketal:

Precision is a weitable powder.

EPA Reg. No. 100-750

EPA Est. 10163-AZ-1

KEEP OUT OF REACH OF CHILDREN. CAUTION

booklet on bag

This outer prosective bag contains Precision in 8 small, inner water soluble packets. Entire inner packets and their contents dissolve in water. After opening outer bag, immediately dump the required number of unopened packets into the parketly filled sprayer or mot tank. On not handle the soluble packets of expose them to Mosture, since this may hadge the southle packets of expose them to mostly since that it cause augmand. If not using all water soluble packets, press to the outer projective bag for storage, Chba-Geigy and the Seler storage to the commend against using parket water soluble packets of Practs Such practice may result in injury and/or telure of performance.

8 x 10 Ounce Water Soluble Packets Total 8.0 Ounces Net Weight



DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and Condition of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, metlectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks reterred to above. Ciba-Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it subject to the foregoing Conditions of Sale and irranty, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy

DIRECTIONS FOR USE

It is a violation of Inderal law to use this product in a manner inconsistent with its labeling. Observe all precautions and limitations on this label and on the labels of each product used in tank mixtures with this product.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAU-TIONS ON THIS LABEL MAY RESULT IN CROP MUURY, KLERAL RESIDUES, AMOVOR POOR MISECT CONTROL.

Do not apply this product in such a manner as to directly, or A C C E P T E D through drift, expose workers or other persons. The area being through drift, expose workers or other persons. The area being treated must be vecated by unprotected persons.

Do not enter treated area without protective clothing until sprays have dired.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given which inform workers of areas or fields that may not be antered without specific protective clothing, period of time field must be vacated, and appropriate actions to take in case of accidental exposure as described under Procuulionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "CAUTION. (Area or held description) treated with Precision on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, flush eyes or skin with plenty of water. Call a physician if irritation persists. Remove and wash confaminated clothing before

General Internation

For control of selected insects, such as whiteflies (sweet potato and greenhouse), soft scales, fungus gnats and shore flies, in landscape ornamentals, container grown ornamentals, non-bearing fruit and nut trees in nursenes. Christmas tree plantations, ground covers, greenhouse, lath and shade house ornamental crops, and interiorscapes.

Precision should be used in conjunction with IPM practices including the early detection of target insect populations, threshold treatment levels, cultural control practices, and other procedures to manage target pest populations. Refer to local extension or university personnel or brochures for information on IPM practices.

Application

For control of whiteflies and soft scale insects (wax, brown soft, euonymus, etc.), apply 2-4 oz. Precision per 100 gals, water as soon as the presence of target insects are detected. Sprays for control of scale should be directed to the crawler stage. Repeat applications at 7-14 day intervals to control the immature stages of the insect pests. Apply the spray mixture uniformly to all surfaces of the plants, particularly the stems and the undersides of the leaves. Spray the plants to the point of runoff. Follow the manufacturer's recommendations for proper use of application equipment.

For the control of fungus gnats (Bradysia spp.) and shore flies (Scatella spp.) apply 4 oz. Precision in 100 gals, of water as a spray or dranch to all surfaces where the insect pests may breed. This includes the potting media surfaces, bench tops, undersides of benches, and the areas under the benches that may be contaminated with potting media or algae. Repeat applications on a 10-14 day schedule to minimize reinfestations.

Precision will not control the adult stages of the insect pests. For control of adults, insecticides labeled for adult insect control should be used

Chemigation: Do not apply this product through any type of irngation system

Notes: (1) Precision has been evaluated for safety to many greenhouse, nursery, butb, and ornamental plants as a drunch for control of fungus gnats. No phytotoxicity has been seen on amarylis. begonia, chrysanthemum, geranium, gerbera, impabens, poinsettra, azalea, or Boston fern. (2) Leaf stunbing has been observed after multiple applications of tokar sprays of the high rate of Precision on several pointsettra varieties. Varieties should be evaluated for phytotoxicity prior to making applications to the entire crop Applications should not be made after bract formation, (3) Slight phytotoxic symptoms have occasionally been seen on tokage of

1 OCT 1993

Under the Federal Insecticide. Fungicide, and Accenticide Act C) autoriod, his the pesticide registared under EPA Log. No. /00-75 0



ageratum, celosia, coleus, dianthus, dusty miller, impatiens, mangold, pansy, petunia, snapdragon, and verbena after multiple tokar applications. The grower should determine if Precision can be used safely on ornamental crops prior to use. To evaluate possible phytoticic effects, before treating ornamental crops, the grower should test Precision, on a small group of plants, under the same condisions in which they will be grown, for at least 2 weeks.

Mixiga lectracions

Prepare no more spray mixture than is necessary for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate thoroughly before and during application. "Rolling" agitation is suggested to heep Precision thoroughly mined and in suspension. Flush spray tank thoroughly with clean water daily after use and dispose of pesticide rinsate by application to a previously treated area.

Precision Alone: Add ½ to 2/s of the required amount of water to the spray or mix tank. With the agitator running, drop the required number of unequent soluble paciets of Precision into the tank all at once. Continue agitation while adding the remander of the water. Begin application of the spray solution after the packets have dissolved, and the material has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Precision + Tank Mixturne: Precision is compatible with most commonly used insecticides, fungicides, and spray adjuvants used in the production of bedding plants, ornamentals, and other crops grown in nurseries. When using Precision in tank mixture with other pesticides, observe all directions for use, and precautions on the respective tank mix label.

Add $\frac{1}{2}$ to $\frac{2}{3}$ of the required amount of water to the spray or mix tank. With the agitator running, drop the required number of unopened soluble packets of Precision into the tank all at once. Continue agitation while adding the remainder of the water. Tank mix partners may be added only when: (1) the Precision packets and their contents have dissolved and dispersed into the mix water, and (2) when the tank has filled with water. Maintain agitation when adding the tank mix partners and while spraying the solution onto the crop.

Precautions: (1) Water soluble paciets of Precision must be completely dissolved and dispersed in water before any other tank mix partner, including micronutrients or other liquid or dry tertilizer, is added to the spriny solution. (2) When more than one product is packaged in water soluble packets, add all of those products at once to the mix or soray tank. Apply any other products after the soluble packets have completely dissolved, and the material has completely dispersed into the mix water.

Note: Do not let spray mixture stand overnight in the spray tank. Rinse spray tank at the end of the day.

Table 1: Application to Control WhiteHies and Scale

Area to be treated (sq. ft)	20,000	10,000	5.000
Water Volume	100 gals.	50 gals.	25 gals.
Number of Water Soluble Packets	2-4	1-2	1

Table 2: Application to Control Fungus Gnats and Shore Flies

Drench* Area to be treated (sq. ft.)	20,000	10,000	5,000	1 000
Number of Water Soluble Packets	80	40	20	4

^{*100} gals, will drench 1,000 sq. ft.

Spray Area to be treated (sq. ft.)	20,000	10,000	5,000
Water Volume	100 gals.	50 gals	25 gals
Number of Water Soluble Packets	4	2	1

Electrostatic Spraying Systems, PulsFog Φ Systems, or Other Law Volume Systems

To calculate the amount of product to be applied, use the appropriate amount of Precision for the square footage to be treated with spray as listed. The amount of carrier (water) is dependent on the amount needed for adequate coverage.

Storage and Disposal

Pesticide Sterage – Do not store this product under wet conditions; store in a cool, dry place. Handle outer container carefully to avoid breakage of inner soluble packets.

Penticide Dispesal - Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal — Do not reuse ower bag. Dispose of empty outer bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, dispose of by burning. If bag is burned, stay out of smole.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Precautionary Statements Hazards to Humans and Domestic Animals CALITION

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

Statement of Practical Tradement

If on skin: Wash with plenty of soap and water, Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if writation persists.

If inhaled: Move victim to fresh air.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Oriff and runoff from treated sites may be hazardous to aquatic invertebrates in adjacent aquatic sites. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not confarmate water when disposing of equipment wash waters.

Accu-Pak™ and Precision™ trademarks of Ciba-Geigy Corporation U. S. Patent No. 4,215,139

©1993 Ciba-Geigy Corporation

Ciba Plant Protection Ciba-Geigy Corporation Greensboro, North Carolina 27419 CGA 137L1 063

> This booklet manufactured using post-consumer recycled paper





Precision

INSECT **GROWTH** REGULATOR

For control of selected insects, including white-flies, soft scales, fungus gnots, and shure flies in landscape ornamentals, nurseries, and greenhouse emamental crops

Inert Ingredients: 75%

Total: Precision is a wettable

powder See directions for use in attached backlet on bag.

EPA Rog. No. 188-758 EPA Est. 10163-AZ-1

Acco-Pah¹⁰ and Procision¹⁰ tradomarks of Cibe-Golgy Corporation U. S. Patent No. 4.215,139

8 x 1.0 Ounce **Water Soluble Packets**

Total 8.0 Ounces **Not Weight**



KEEP OUT OF REACH OF CHILDREN.

CAUTION

Precautionary Statements

Hammis to Humans and Demonis

Additions
Harmful if inhaled or absorbed through
the slon. Causes moderate eye intoson.
Audi breathing spray mist. Audi conject
with eyes, sloth, or clothing. Remove con-taminated clothing and wash before
reuse. Wash thoroughly with soap and
waster stater franchism. water after handling

Statement of Practical Treatment

If on aidn: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water Call a physician if irritation persists

If inheled: Move victim to fresh air

Environmental Hazards

This product is time to fish and aquatic invertebrates. Drift and runoff from treated sites may be hazardous to aquatic invertebrates in adjacent aquatic sites for terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contam-nate water when disposing of equipment wash waters

On not apply this product through any type of irrigation system

©1993 Cibo-Golgy Corporation

Cito Plant Protection Cito-Galgy Corporation Granactions, North Carolina 27419 **CGA 137L1**

リッチル

8 X 1.0 Ounce Water Soluble Packets

PrecisionTM

٠,

Insect Growth Regulator

For control of selected insects, including whiteflies, soft scales, fungus gnats, shore flies, and lepidopterous leafminers in landscape ornamentals, nurseries, greenhouse ornamental crops, and other areas as specified on the label.

Active Ingredient:

Fenoxycarb: (Ethyl [2-(4-phenoxyphenoxy) ethyl] carbamate)...25% Inert Ingredients: <u>75%</u> Total: 100%

Precision is a wettable powder.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet on bag.

EPA Reg. No. 100-750

EPA Est. 10163-AZ-1

This outer protective bag contains Precision in 8 small, inner water soluble packets. Entire inner packets and their contents dissolve in water. After opening outer bag, immediately dump the required number of unopened packets into the partially filled sprayer or mix tank. Do not handle the soluble packets or expose them to moisture, since this may cause rupturing. not using all water soluble packets, press to reseal the outer protective bag for storage. Ciba-Geigy and the Seller strongly recommend against using partial water soluble packets of Precision. Such practice may result in injury and/or failure of performance.

8 X 1.0 Ounce Water Soluble Packets

Total 8.0 Ounces Net Weight

CGA 137L1 063

50414

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire <u>Directions for Use</u> and <u>Conditions of Sale and Warranty</u> before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

)

The <u>Directions for Use</u> of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the <u>Directions for Use</u> subject to the inherent risks referred to above. <u>Ciba-Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it subject to the foregoing <u>Conditions of Sale and Warranty</u>, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.</u>

600+14

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Observe all precautions and limitations on this label and on the labels of each product used in tank mixtures with this product.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, ILLEGAL RESIDUES, AND/OR POOR INSECT CONTROL.

Do not apply this product in such a a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

Reentry Statement

Do not enter treated area without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated, and appropriate actions to tak? in case of accidental exposure as described under <u>Precautionary Statements</u> on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "CAUTION.

(Area or field description) treated with Precision on (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, flush eyes or skin with plenty of water. Call a physician if irritation persists. Remove and wash contaminated clothing before reuse."

General Information

For control of selected insects, such as whiteflies (sweet potato and greenhouse), soft scales, fungus gnats and shore flies, in landscape ornamentals, container grown ornamentals, non-bearing fruit and nut trees in nurseries, Christmas tree plantations, ground covers, greenhouse, lath and shade house ornamental crops, and interiorscapes.

For control of lepidopterous leafminers (e.g., citrus leafminer and azalea leafminer) on nonbearing citrus and woody ornamentals. Do not apply to trees that will bear fruit within one year of application.

Precision should be used in conjunction with IPM practices including the early detection of target insect populations, threshold treatment levels, cultural control practices, and other procedures to manage target pest populations. Refer to local extension or university personnel or brochures for information on IPM practices.

Application

For control of whiteflies and soft scale insects (wax, brown soft, euonymus, etc.), apply 2-4 oz. Precision per 100 gals. water as soon as the presence of target insects are detected. Sprays for control of scale should be directed to the crawler stage. Repeat applications at 7-14 day intervals to control the immature stages of the insect pests. Apply the spray mixture uniformly to all surfaces of the plants, particularly the stems and the undersides of the leaves. Spray the plants to the point of runoff. Follow the manufacturer's recommendations for proper use of application equipment.

For the control of fungus gnats (<u>Bradysia</u> spp.) and shore flies (<u>Scatella</u> spp.) apply 4 oz. Precision in 100 gal. of water as a spray or drench to all surfaces where the insect pests may breed. This includes the porting media surfaces, bench tops, undersides of benches, and the areas under the benches that may to contaminated with potting media or algae. Repeat applications on a 10-14 day schedule to minimize reinfestations.

For control of lepidopterous leafminers, apply 6.0 to 8.0 oz. Precision when oviposition begins on the new flush of growth. A second application should be made 14-21 days later. Application should be made in full coverage sprays.

Precision will not control the adult stages of the insect pests. For control of adults, insecticides labeled for adult insect control should be used.



80444

<u>Chemigation</u>: Do not apply this product through any type of irrigation system.

(1) Precision has been evaluated for safety to many Notes: greenhouse, nursery, bulb, and ornamental plants as a drench for control of fungus gnats. No phytotoxicity has been seen on amaryllis, begonia, chrysanthemum, geranium, gerbera, impatiens, poinsettia, azalea, or Boston fern. (2) Leaf stunting has been observed after multiple applications of foliar sprays of the high rate of Precision on several poinsettia varieties. ties should be evaluated for phytotoxicity prior to making applications to the entire crop. Applications should not be made after bract formation. (3) Slight phytotoxic symptoms have occasionally been seen on foliage of ageratum, celosia, coleus, dianthus, dusty miller, impatiens, marigold, pansy, petunia, snapdragon, and verbena after multiple foliar applications. grower should determine if Precision can be used safely on ornamental crops prior to use. To evaluate possible phytotoxic effects, before treating ornamental crops, the grower should test Precision, on a small group of plants, under the same conditions in which they will be grown, for at least 2 weeks.


Mixing Instructions

Prepare no more spray mixture than is necessary for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate thoroughly before and during application. "Rolling" agitation is suggested to keep Precision thoroughly mixed and in suspension. Flush spray tank thoroughly with clean water daily after use and dispose of pesticide rinsate by application to a previously treated area.

Precision Alone: Add 1/2 to 2/3 of the required amount of water to the spray or mix tank. With the agitator running, drop the required number of <u>unopened</u> soluble packets of Precision into the tank all at once. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the packets have dissolved, and the material has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

<u>Precision + Tank Mixtures</u>: Precision is compatible with most commonly used insecticides, fungicides, and spray adjuvants used in the production of bedding plants, ornamentals, and other crops grown in nurseries. When using Precision in tank mixture with other pesticides, observe all directions for use, and precautions on the respective tank mix label.

Add 1/2 to 2/3 of the required amount of water to the spray or mix tank. With the agitator running, drop the required number of unopened soluble packets of Precision into the tank all at once. Continue agitation while adding the remainder of the water. Tank mix partners may be added only when: (1) the



POFIL

Precision packets and their contents have dissolved and dispersed into the mix water, and (2) when the tank has filled with water. Maintain agitation when adding the tank mix partners and while spraying the solution onto the crop.

Precautions: (1) Water soluble packets of Precision must be completely dissolved and dispersed in water before any other tank mix partner, including micronutrients or other liquid or dry fertilizer, is added to the spray solution. (2) When more than one product is packaged in water soluble packets, add all of those products at once to the mix or spray tank. Apply any other products after the soluble packets have completely dissolved, and the material has completely dispersed into the mix water.

Note: Do not let spray mixture stand overnight in the spray tank. Rinse spray tank at the end of the day.

Table 1: Application to Control Whiteflies and Scale

	(sq. feet)			
Area to be Treated (sq. ft.)	20,000	10,000	5,000	
Water Volume	100 gals.	50 gals.	25 gals.	
Number of Water Soluble Packets	2-4	1-2	1	

Table 2: Application to Control Fungus Gnats and Shore Flies

<pre>Drench* Area to be Treated (sq. ft.)</pre>	20,000	10,000	5,000	1,000
Number of Water Soluble Packets	80	40	20	4

*100 gal. will drench 1,000 sq. ft.

Spray Area to be Treated (sq. ft.)	20,000	10,000	5,000	
Water Volume	100 gals.	50 gals.	25 gals.	
Number of Water Soluble Packets	4	2	1	

Table 3: Application to Control Lepidopterous Leafminers

Spray Amount Per Acre	6 oz.	8 oz.
Number of Water Soluble Packets	6	8

Electrostatic Spraying Systems, PulsFog® Systems or Other Low Volume Systems:

To calculate the amount of product to be applied, use the appropriate amount of Precision for the square footage to be



treated with spray as listed. The amount of carrier (water) is dependent on the amount needed for adequate coverage.

Storage and Disposal

<u>Pesticide Storage</u> - Do not store this product under wet conditions; store in a cool, dry place. Handle outer container carefully to avoid breakage of inner soluble packets.

<u>Pesticide Disposal</u> - Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

<u>Container Disposal</u> - Do not reuse outer bag. Dispose of empty outer bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, dispose of by burning. If bag is burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

Statement of Practical Treatment

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If inhaled: Move victim to fresh air.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Drift and runoff from treated sites may be hazardous to aquatic invertebrates in adjacent aquatic sites. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Accu-PakTM and PrecisionTM trademarks of Ciba-Geigy Corporation U. S. Patent No. 4,215,139

©1993 Ciba-Geigy Corporation

Ciba Plant Protection Ciba-Geigy Corporation Greensboro, North Carolina 27419

CGA 137L1 063

[GANNONC.LABELP]PRECWSP - 9/9/93

Adhesive Back Label

Precision TM

Insect Growth Regulator

For control of selected insects, including whiteflies, soft scales, fungus gnats, and shore flies in landscape ornamentals, nurseries, and greenhouse ornamental crops

Active Ingredient:

Fenoxycarb: (Ethyl [2-(4-phenoxyphenoxy) ethyl] carbamate)...25% Inert Ingredients: 75% Total:

Precision is a wettable powder.

See directions for use in attached booklet on bag.

EPA Reg. No. 100-750

EPA Est. 10163-AZ-1

Accu-PakTM and PrecisionTM trademarks of Ciba-Geigy Corporation U. S. Patent No. 4,215,139

8 X 1.0 Ounce Water Soluble Packets

Total 8.0 Ounces Net Weight

KEEP OUT OF REACH OF CHILDREN.

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

Statement of Practical Treatment

If on skin: Wash with plenty of soap and water. Get medical
attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If inhaled: Move victim to fresh air.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Drift and runoff from treated sites may be hazardous to aquatic invertebrates in adjacent aquatic sites. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Chemigation

Do not apply this product through any type of irrigation system.

@1993 Ciba-Geigy Corporation

Ciba Plant Protection Ciba-Geigy Corporation Greensboro, North Carolina 27419

CGA 137L1 063

[GANNONC.LABELP]PRECWSP - 9/8/93