

100-752

10/20/2003

1/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Ms. Michele A. Schulz
NAFTA Fungicide Team Leader
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419

OCT 20 2003

Dear Ms. Schulz:

Subject: Primo® WSB

EPA Registration No. 100-752
Application and Your Letter Dated September 10, 2003,
Request to Amend Registration by Revising "First Aid
Statements" To Comply with EPA PR Notice 2001-1, Revise
the REI to 0 Days and Other Minor Changes; and
Resubmission of Labeling October 20, 2003

The proposed amendments to the subject registration reflected on the submitted labeling have been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) as amended, provided that you:

1. Revise the claim that reads: "Primo WSB should be used only in accordance with recommendations on this label or in separately published Syngenta supplemental labeling recommendations for this product." to read: "Primo WSB must be used only in accordance with the directions of use on this label or in supplemental labeling accepted by the US Environmental Protection Agency."
2. Submit one copy of the final printed labeling prior to shipment of this product under the revised labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of this product under the revised labeling constitutes acceptance of these conditions. A stamped copy of the accepted labeling is enclosed for your records.

Sincerely yours,

Joanne I. Miller
Joanne I. Miller

Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

Primo® WSB

For Turf Growth Management

For managing growth, improving quality, and edging of warm- and cool-season grasses

Active Ingredient:	
Trinexapac-ethyl (CAS No. 95266-40-3).....	25.0%
Other Ingredients:	75.0%
Total:	100.0%

Primo WSB is a wettable powder packaged in a water-soluble bag. This outer protective container contains Primo WSB in an inner water-soluble bag. Entire inner bag and contents dissolve in water. After opening outer container, immediately dump entire unopened inner bag into the partially filled sprayer or mix tank. Do not handle the soluble bag or expose it to moisture, since this may cause rupturing. Using partial bags of Primo WSB voids any warranty by Syngenta Crop Protection, Inc. on its safety and efficacy.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements on back of container. See directions for use below.

EPA Reg. No. 100-752

EPA Est.

EPA Est.

(Superscript is first letter of batch code on container)

SCP

4 ounces

Net Weight

_____ Total Net Weight

ACCEPTED
with COMMENTS
In EPA Letter Dated
OCT 20 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

100-752

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER	
For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing and breathing dust or spray mist.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material

- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Do not apply when weather conditions favor drift from treated areas.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Primo WSB should be used only in accordance with recommendations on this label or in separately published Syngenta supplemental labeling recommendations for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter treated areas without footwear until sprays have dried. The restricted entry interval (REI) is 0 days.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without footwear until sprays have dried.

Observe all precautions and limitations on this label and on the labels of each product used in tank mixtures with this product.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN UNEVEN GROWTH REDUCTION OR SEVERELY STUNTED TURF.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage:

Store in a cool, dry place. Do not store this product under wet conditions. Handle outer container carefully to avoid breakage of inner soluble bag.

Pesticide Disposal:

Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, they must be disposed of according to federal, state, or local procedures. Contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Container Disposal

Plastic Containers: Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or alternatives allowed by State and local authorities.

FOR BULK AND MINIBULK CONTAINERS:

Container Disposal and Precautions

When this container is empty, replace the cap or cover and seal all openings that have been opened during use; and return the container to the point of purchase or to a designated location named at time of purchase of this product. The container must only be refilled with this pesticide product. **DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE.** Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, contact Syngenta Crop Protection, Inc. at 1-800-888-8372. If not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

GENERAL INFORMATION

Primo WSB for turf growth management reduces the frequency of mowing and the amount of grass clippings by reducing the growth of warm- and cool-season grasses. Other benefits such as increased turf density, increased color, and increased turf quality are frequently observed after Primo WSB applications.

Primo WSB can be applied to well-maintained, quality turfgrass areas such as residential and commercial lawns, golf courses, sod farms, sports fields, cemeteries, and similar areas. Primo WSB is useful in the management of difficult-to-mow areas and can be used to minimize the need for edging turfgrass along sidewalks, curbs, parking lots, driveways, flower beds, fences and around posts, storage sheds, and trees.

Primo WSB reaches the growing point by foliar uptake and is rainfast from rainfall or irrigation after one hour. Watering-in is not necessary for activation.

NOTICE TO USER: Plant tolerances to Primo WSB have been found to be acceptable for the grasses listed on this label. Due to the large number of species and cultivars of grasses, it is impossible to test every one for tolerance to Primo WSB. Neither the manufacturer nor the seller has determined whether or not Primo WSB can be used safely on grasses not specified on this label. The professional user should determine if Primo WSB can be used safely prior to commercial use. Before using Primo WSB for grasses not listed in the application table, test Primo WSB on a small scale first. Apply the lower recommended rate for the turf setting (lawn, fairway, etc.) and evaluate for phytotoxicity and growth inhibition prior to widespread use.

APPLICATION INFORMATION

Environmental conditions, management, and cultural practices that affect turf growth and vigor will influence the response of the turf to Primo WSB applications. These include fertility level, moisture availability, plant vigor, height, and frequency of mowing.

The rates presented in the Application Rate Table provide approximately 50% growth inhibition over a 4-week period with little or no discoloration of turf growing under favorable conditions. Excessive turf growth, which may occur under high fertilization program or during spring flushes, may require higher rates of Primo WSB. Under these conditions, Primo WSB rates may have to be increased up to 50% to provide an adequate length of control. For extended growth suppression up to 8 weeks, where temporary discoloration can be tolerated, a maximum of twice the recommended Primo WSB rate from the Application Rate Table may be applied. Primo WSB use rates may need to be reduced up to 50% for turfgrass grown under conditions of low fertility, compaction, or other factors which stress the turf.

Multiple applications can be made each growing season, but do not exceed a total of 4 oz./1,000 sq. ft. (174 oz./acre) per year.

Mowing

Generally, Primo WSB provides more suppression when turfgrasses are mowed at lower mowing heights than higher mowing heights. Application rates have been selected for typical mowing heights. The application rate may need to be adjusted depending on actual mowing conditions. It is suggested that Primo WSB not be applied less than 4 hours before or 4 hours after mowing to minimize any potential injury.

Application Timing

Apply Primo WSB to actively growing turf. Reduce Primo WSB rates on turf going into dormancy due to high or low temperatures or lack of moisture.

Repeat applications of Primo WSB may be made as soon as the turf resumes growth or more suppression is desired.

Equipment

Apply Primo WSB with a sprayer capable of accurate and uniform application. Clean spray equipment thoroughly before use. Primo WSB may be applied in backpack sprayers, hand sprayers, boom sprayers, and by spray gun application devices. Rinse sprayer with clean water after use and dispose of rinsate in an approved manner.

Mixing Instructions

Primo WSB should be applied in sufficient carrier (e.g., 0.50 to 4.0 gallons of water per 1,000 sq. ft.) to provide uniform and thorough coverage.

Prepare no more mixture than is necessary for the application to be made. Agitate thoroughly before and during application. Primo WSB may be tank-mixed with many commonly used pesticides and liquid fertilizers.

If using Primo WSB in a tank mixture, observe all directions for use, sites, use rate dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate should be exceeded and the most restrictive label precautions and limitations should be followed. Primo WSB should not be mixed with any product that bears a label prohibiting such mixing.

Precautions: (1) Soluble prepacks of Primo WSB must be dissolved completely in water prior to adding other tank mix partner(s), including micronutrients or other liquid or dry fertilizer to the spray solution. (2) When Primo WSB is tank mixed with Banner® or Bayleton®, use a compatibility agent at 3 pts./100 gals. of water. (3) When preparing a

tank mixture, if more than one product is packaged in soluble prepacks, add all of those products together at once. Add any other tank mix partners after the soluble bags have dissolved, and the material has completely dispersed into the mix water.

Backpack and Hand Sprayers

Primo WSB Alone: Add $1/2$ - $2/3$ of the required water to the spray tank. Drop the required number of unopened soluble prepacks of Primo WSB into the tank all at once. Allow the mixture to stand for 3-5 minutes to let the soluble package(s) dissolve. Then, close and vigorously shake for 30 seconds. Begin application after the soluble packages have dissolved and the material has completely dispersed into the mix water.

Boom and Hand Gun Sprayers

Primo WSB Alone: Add $1/2$ - $2/3$ of the required amount of water to the spray tank. Then, while agitating, drop the required number of Primo WSB prepacks into the tank all at once. Continue agitation while adding the remainder of the water. Apply spray solution after the bags have dissolved and the material has completely dispersed into a uniform mixture. Agitate during application of the resulting mixture.

Primo WSB + Tank Mixtures: Add $1/2$ - $2/3$ of the required amount of water to the spray or mixing tank. Then, while agitating, drop the required number of unopened soluble prepacks of Primo WSB into the tank all at once. Continue agitation while adding the remainder of the water. Tank mix partners should only be added when: (1) the Primo WSB bags and their contents have dispersed into the mix water, and (2) after the tank mix has been filled with water. Stop agitation long enough to add the tank mix partner and then resume agitation while spraying the solution.

Tank Mix Partner Addition Sequence:

1. Add gel formulations to the spray tank first. Allow the packet to completely dissolve and the product to completely disperse into the mix water.
2. Next, add wettable powders to the tank allowing them to completely suspend in the tank before proceeding. This process can be hastened by premixing the product in water before adding to the tank.
3. Add flowables or suspensions next.
4. Add emulsifiable concentrates last.
5. Do not leave tank mix combinations in the spray tank for prolonged periods without agitation. Mix and apply them the same day.
6. Primo WSB should not be mixed with any product which prohibits such mixing. No label application rate should be exceeded and the most restrictive label

precautions and limitations should be followed. Refer to the tank-mix product label(s) for further information.

Dyes

A water-soluble dye to mark the sprayed area can be used to ensure uniform application, and to avoid skips and overlaps. Refer to the instructions which accompany the dye for proper concentrations and mixing procedures.

Application Near and Around Monuments and Hardscape Materials

Primo WSB at normal dilution rates will not stain brass, bronze, concrete, marble, granite, or other types of stone. Before using Primo WSB around other types of stone, test on a small scale basis first.

POA ANNUA CONVERSION/RENOVATION

Primo WSB can be applied to existing turf infested with stands of *Poa annua* as part of an overseeding/renovation program. Such an application allows better germination and seedling growth of more desirable turfgrasses. Primo WSB, in conjunction with the use of appropriate cultural practices that help to ensure the vigor and growth of new seedlings will also result in fewer clippings and thus, less maintenance traffic on new seedlings. Since Primo WSB is foliarly absorbed, seed germination is not affected. Apply Primo WSB 1–5 days before seeding; and before verticutting, scalping, spiking, or other similar operations.

Temporary initial discoloration is possible with aggressive application rates of Primo WSB to turf with *Poa annua*. Primo WSB applications the following spring should use the upper end rate for the turf type and setting listed in the Application Rate Table. Actual conversion success will also depend on growing conditions, fertilization, rainfall, and other agronomic and environmental conditions.

BERMUDAGRASS OVERSEEDING

An application of Primo WSB to bermudagrass will enhance the establishment of cool-season turfgrasses. Primo WSB, in conjunction with the use of cultural practices that help ensure new seedling vigor and growth will also result in fewer clippings and less maintenance traffic on new seedlings. Primo WSB is foliarly absorbed, so germination and seedling growth is not affected.

Apply Primo WSB before verticutting, scalping, spiking, or other similar operations to the bermudagrass. Apply Primo WSB 1 to 5 days before seeding.

Temporary initial discoloration is possible with aggressive application rates of Primo WSB. Use normal seeding rates for your area and turf setting (lawn, fairway, etc.). Actual overseeding success will also depend on growing conditions, fertilization, rainfall, and other agronomic and environmental conditions. For maintenance application see the Application Rate Table.

APPLICATION WITH TURF MARKING PAINT

Primo WSB can extend the duration of marking visibility when applied before or with marking agents. Mix Primo WSB with water first when combining with latex based marking agents. Refer to the marking agents product label for further instructions.

Apply Primo WSB at 0.50 oz./gal. of marking paint mix to treat approximately 1,000 sq. ft. of line surface area. Refer to the Primo WSB + Tank Mixtures section of this label for additional instructions.

Notes: (1) Do not apply to bare ground since Primo WSB is foliarly absorbed. (2) Turf treated with Primo WSB should continue to receive regular good maintenance practices including irrigation; fertilization; and weed, disease, and insect control when necessary, and as recommended for quality turf. As some herbicides can injure turf, tank mixes with Primo WSB should be tested on a small scale before widespread use. (3) Primo WSB may cause temporary yellowing. This usually disappears about one week after application. To minimize yellowing and to enhance the green color of turf, apply readily available nitrogen at 0.2 to 0.5 lbs. of actual nitrogen per 1,000 sq. ft. If desirable, a recommended rate of an iron source can also be used. (4) Full growth regulation by Primo begins at about 3 to 5 days after application.

Precautions: (1) Do not apply Primo WSB through any type of irrigation system. (2) Do not graze areas or feed clippings to livestock.

Table 1. Recommended Primo WSB Application Rates^{1,2}

Turf Type	Residential & Commercial Turf ³		Golf Course Fairways (Cut at 0.50" or less)		Golf Course Greens		Edging/Banding ⁴	
	oz./1,000 sq. ft.	Area/4 oz. bag (sq. ft.)	oz./1,000 sq. ft.	Area/4 oz. bag (sq. ft.)	oz./1,000 sq. ft.	Area/4 oz. bag (sq. ft.)	oz./1,000 sq. ft.	Area 4/oz. bag (sq. ft.)
Cool-Season								
Bentgrass	0.375	10,667	0.125	32,000	0.063	63,492	0.50	8,000
Fescue, Red	0.375	10,667					0.50	8,000
Fescue, Tall (Ky-31)	0.50	8,000					0.50	8,000
Fescue, Tall (Turf Types)	0.375	10,667					0.50	8,000
Kentucky Bluegrass	0.30	13,333	0.125	32,000			0.375	10,667
Mixture (Bentgrass/ <i>Poa annua</i>)			0.125 ⁵	32,000	0.075 ⁵	53,333		
Mixture (K. Bluegrass/ Fescue/Ryegrass)	0.375	10,667						
Mixture (K. Bluegrass/ Ryegrass/ <i>Poa annua</i>)			0.25 ⁵	16,000				
Ryegrass, Annual	0.50	8,000					0.50	8,000
Ryegrass, Perennial	0.50	8,000	0.25	16,000			0.50	8,000
Warm-Season								
Bahiagrass	0.50	8,000					0.50-1.0	8,000-4,000
Bermudagrass⁶								
Bermudagrass, Common	0.375	10,667	0.125	32,000			0.50-1.0	8,000-4,000
Bermudagrass Other Hybrids	0.125	32,000	0.10	40,000			0.25-0.375	16,000-10,667

/Continued ...

12/15

Table 1. Recommended Primo WSB Application Rates^{1,2} (Continued)

Turf Type	Residential & Commercial Turf ³		Golf Course Fairways (Cut at 0.50" or less)		Golf Course Greens		Edging/Banding ⁴	
	oz./1,000 sq. ft.	Area/4 oz. bag (sq. ft.)	oz./1,000 sq. ft.	Area/4 oz. bag (sq. ft.)	oz./1,000 sq. ft.	Area/4 oz. bag (sq. ft.)	oz./1,000 sq. ft.	Area 4/oz. bag (sq. ft.)
Bermudagrass, Tifdwarf	0.10	40,000	0.10	40,000	0.031	129,032	0.025-0.375	16,000-10,667
Bermudagrass, Tifgreen (328)	0.125	32,000	0.10	40,000	0.063	63,492	0.375	10,667
Bermudagrass, Tifway (419)	0.19	21,053	0.125	32,000			0.375	10,667
Buffalograss	0.50	8,000					0.50	8,000
Carpetgrass	0.125-0.20	32,000-20,000					0.25	16,000
Centipedegrass	0.25	16,000					0.50	8,000
Kikuyugrass	0.15-0.25	26,667-16,000	0.15	26,667			0.50	8,000
St. Augustinegrass	0.125-0.19	32,000-21,053					0.50-1.0	8,000-4,000
St. Augustinegrass, Texas Common	0.125	32,000					0.25	16,000
Zoysiagrass	0.125	32,000	0.063	63,492			0.375	10,667

¹These rates should provide 50% suppression of turf growth under good growing conditions for 4 weeks.

²Seedheads: At rates equal to or higher than the rates in the Table on back of container, Primo WSB provides seedhead suppression of hybrid bermudagrass, and partial seedhead suppression of annual bluegrass, bahiagrass, buffalograss, carpetgrass, common bermudagrass, Kentucky bluegrass, and tall fescue. Primo must be applied prior to seedhead formation.

³Including, but not limited to home lawns, parks, recreation areas, golf course roughs, cemeteries, business sites, sports fields, and sod farms.

⁴Primo WSB can be applied along the perimeter of lawns, sidewalks, curbs, parking lots, driveways, posts, storage buildings, pet pens, fences, or other areas. Primo WSB can be used around trees, shrubs, flower beds, and other border plants or similar areas with no injury. Apply Primo WSB in an 8 to 12-inch band with a single nozzle sprayer. Use the higher concentration to reduce the growth of the turfgrass into adjacent areas. Important: When calculating the number of bags to use, always round off to the nearest number of whole bags. Do NOT use partial bags.

⁵Where yellowing of *Poa annua* is a concern, use half this rate and apply more frequently. For *Poa annua* conversion/renovation, where temporary discoloration can be tolerated, twice this rate may be applied.

⁶Bermudagrass overseeding preparation: 0.25 oz./1,000 sq. ft.

14/15

Banner®, Primo®, and the Syngenta logo are trademarks of a Syngenta Group Company

Bayleton® trademark of Bayer Crop Science

©2003 Syngenta Crop Protection, Inc.

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP

000100-00752.20030915b.PrimoWSBFirstAid.pdf

PRI WSB752A AMD DRAFT aa-10-9-03 clean