

100-758

3/12/2004

Jacket

1/14

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Larry Zang
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

MAR 12 2004

Dear Mr. Zang:

SUBJECT: Maxim FS
EPA Registration Number 100-758
Your Submission of September 23, 2004

The amendment referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) is acceptable provided you make the following changes:

Ingredient Statement - Page 1:

Line up the percentages of fludioxonil, "other ingredients" and "total" so that the decimal points fall along the same vertical line.

General Information - Mixing Procedures - Page 4:

In the paragraph beginning "Federal law requires . . ." change the sentence "Corn and sorghum forage may not be grazed until 30 days after planting" to "Forage may not be grazed until 30 days after planting"

Please submit one copy of your final printed labeling before you release the product for shipment. A stamped copy of the label is enclosed for your records. If you have any questions, please contact Lisa Jones of my staff by phone at (703) 308-9424 or by e-mail at jones.lisa@epa.gov.

Sincerely,

Lisa Jones for

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure: Stamped Label

CONCURRENCES							
SYMBOL ▶	7505C						
SURNAME ▶	Lisa Jones						
DATE ▶	Mar 10, 2004						

Maxim® 4FS

Fungicide

A seed treatment product for protection against certain seed and seedling diseases of crop plants

Active Ingredient:	
Fludioxonil (CAS No. 131341-86-1).....	40.3%
Other Ingredients:.....	59.7%
Total:	100.0%

Maxim 4FS contains 4 lbs. active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-758

EPA Est. 34704-MS-2

Product of Switzerland

Formulated in the USA

SCP 758A-

_____ gallons
Net Content

ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 12 2004

Under the National Insecticide
Registration Act of 1947

100-758

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly. • Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. • Call poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
<p>HOT LINE NUMBER For 24 Hour Medical Emergency Assistance for Incidents Involving Human or Animal Exposure or for Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call 1-800-388-8372</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if absorbed through the skin or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers used closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsates.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

6/14

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Maxim 4FS should be used only in accordance with recommendations on this label or in separately published Syngenta supplemental labeling recommendations for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated:

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage

Store in a cool, dry place.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance in proper disposal methods.

Container Disposal

Triple rinse (or equivalent), then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or alternatives allowed by state or local authorities.

FOR BULK AND MINIBULK CONTAINERS

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS: Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.

REFILL ONLY WITH MAXIM 4FS FLOWABLE. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than Maxim 4FS will result in contamination and may weaken container. After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

GENERAL INFORMATION

Maxim 4FS is a seed treatment fungicide which provides protection against certain soil-borne and seed-borne diseases of crop plants. Maxim 4FS is active against *Fusarium*, *Rhizoctonia*, *Helminthosporium*, and weakly pathogenic fungi such as *Aspergillus* and *Penicillium*. When rate ranges are given, use higher rate when disease pressure is expected to be severe.

MIXING PROCEDURES

Apply Maxim 4FS as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the recommended amount of Maxim 4FS into the required amount of water for the slurry treater and dilution rate to be used. Follow the manufacturer application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow seed to dry before bagging.

Federal law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with fludioxonil fungicide. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs. Corn and sorghum forage may not be grazed until 30 days after planting." Use an EPA-approved dye or colorant that imparts an unnatural color to the seed.

Maxim 4FS does not control diseases caused by *Pythium* spp. or *Phytophthora* spp. If these diseases are expected to be a problem, apply Maxim 4FS tank mixed with Apron XL® LS. If Apron XL LS is not registered on that crop, apply Maxim as a tank mix with another labeled seed treatment product that has activity against these diseases. Consult the label for appropriate use rate and follow all use instructions for the products.

Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing.

BRASSICA (COLE) LEAFY VEGETABLE GROUP

Broccoli (including chinese and raab), brussels sprouts, cabbage (including chinese bok choy, chinese napa, and chinese mustard), cauliflower, cavalo broccola, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

BULB VEGETABLES

Garlic (including great headed), leek, onion (dry bulb and green), welch onion, and shallot.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed or propagating material.

CEREAL GRAINS (INCLUDES FORAGE AND SILAGE)

Barley, buckwheat, corn (field corn, popcorn, and sweet corn), millet (pearl and proso), oats, rice, rye, sorghum, teosinte, triticale, wheat, and wild rice.

Corn (Field Corn and Popcorn): For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.034 – 0.068 fl. oz. of Maxim 4FS per 80,000 kernel count unit of seed or 0.006 to 0.012 mg active ingredient per kernel. Maxim 4FS contains 14.17 grams active ingredient per ounce.

Rate of Maxim 4FS* mg active per kernel	Rate of Maxim 4FS fl. oz. per 80,000 kernel count unit of seed
0.006 mg	0.034 fl. oz.
0.012 mg	0.068 fl. oz.

*Refer to the **General Information** section for additional guidance on rate selection.

Apply Apron XL LS seed treatment in combination with Maxim 4FS for protection against *Pythium* spp. Apply Apron XL LS at 0.0425 fl. oz per 100 lb seed or up to 0.004 mg active ingredient per kernel tank mixed with the recommended rates of Maxim 4FS. Or apply Apron XL LS at 0.085 fl. oz. per 100 lbs. of seed or up to 0.008 mg active ingredient per kernel, tank mixed with the recommended rates of Maxim 4FS. Follow application and other instructions found on the Apron XL LS product label.

Grain Sorghum: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

Apply Apron XL LS in combination with Maxim 4FS for protection against damping-off due to *Pythium* spp. Apply Apron XL LS at 0.32-0.64 fl. oz. per 100 lbs. of seed tank mixed with recommended rates of Maxim 4FS.

For systemic downy mildew control on cultivars which possess sorghum downy mildew resistance, apply Apron XL LS at 0.64 fl. oz. per 100 lbs. of seed as a tank mix with recommended rates of Maxim 4FS. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew grown in areas with recent history of severe disease, apply 1.28 fl. oz. of Apron XL LS tank mixed with recommended rates of Maxim 4FS. Follow application and other instructions found on the Apron XL LS label.

Note: Seed treated with the 1.28 fl. oz. of Apron XL LS should be planted and not stored or carried over to the following year. Only high quality sorghum seed (≥90% germ) should be treated with Apron products.

Sweet Corn: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.08 fl. oz. of Maxim 4FS per 100 lbs. of seed.

Apply Apron XL LS seed treatment product in combination with Maxim 4FS for protection against damping-off due to *Pythium* spp. Apply Apron XL LS at 0.32 fl. oz. per 100 lbs. of seed tank mixed with the recommended rates of Maxim 4FS. For protection against systemic downy mildew of sweet corn, apply Apron XL LS at 1.28 fl. oz. per 100 lbs. of seed tank mixed with recommended rates of Maxim 4FS. Follow application and other instructions found on the Apron XL LS product label.

For all other cereal crops: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

Important: (1) Do not use treated seed for feed, food, or oil. (2) Cereal forage may not be grazed until 30 days after planting.

CUCURBIT VEGETABLES

Chayote, chinese waxgourd, citron melon, cucumber, gherkin, gourd edible (includes hyotan, cucuzza, hechima, chinese okra), momordica, all muskmelon, pumpkin, all squash, and watermelon.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

FLAX, COTTON, CRAMBE, JAJOBA, LESQUERELLA, RAPE, SAFFLOWER, SUNFLOWER

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

FOLIAGE OF LEGUME VEGETABLES

Any cultivar of bean (*Phaseolus* spp.), pea (*Pisum* spp.), and soybean.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

FRUITING VEGETABLES EXCEPT CUCURBITS

Eggplant, groundcherry, pepino, pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

GRASS FORAGE, FODDER, AND HAY

All grasses (including turfgrass, bermudagrass, bluegrass, and fescue or bromegrass), Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed or propagating root or tuber material.

Important: Forage may not be grazed until 30 days after planting.

HERBS AND SPICES

All spice, angelica, anise, anise (star), annatto, balm, basil, borage, burnet, camomile, caper buds, caraway, caraway black, cardamom, cassia bark, cassia buds, catnip, celery seed, chervil, chive, chinese chive, cinnamon, clary, clove buds, coriander (Chinese parsley or cilantro), costmary, culantro (seed), cumin, curry (leaf), dill seed and dill weed, common fennel and florence fennel, fenugreek, grains of paradise, horehound, hyssop, juniper berry, lavender, lemongrass, lovage, lovage seed, mace, marigold, marjoram, mustard (seed), nasturtium, nutmeg, parsley (dried), pennyroyal, pepper (black and white), poppy seed, rosemary, rue, saffron, sage, savory (summer and winter), sweet bay, tansy, tarragon, thyme, vanilla, wintergreen, woodruff, wormwood.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed or propagating root or tuber material.

LEAFY VEGETABLES EXCEPT BRASSICA

Amaranth (leafy, Chinese spinach, tampala), arugula, cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress (garden and upland), dandelion, dock, endive, fennel (finocchio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach (including New Zealand and vine), and swiss chard.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

LEAVES OF ROOT AND TUBER VEGETABLES

Beet (garden and sugar), burdock (edible), carrot, cassava (bitter and sweet), chervil (turnip rooted), chicory, dasheen (taro), parsnip, radish (including oriental), rutabaga, salsify (black), sweet potato, tanager, turnip, and true yam.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed or propagating root or tuber material.

LEGUME VEGETABLES (SUCCULENT OR DRY)

Bean (*Lupinus* spp. – includes grain lupin, sweet lupin, and white lupin), bean (*Phaseolus* spp. – includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, and wax bean), Bean (*Vigna* spp. – includes adzuki bean, asparagus bean, blackeyed pea, catjang, chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, and yardlong bean), broad bean, chickpea, guar, jackbean, lablab bean, lentils, pea (dwarf, edible-pod, English, field, garden, green, snow, sugar snap), pigeon pea, soybean, soybean (immature), and sword bean.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed or propagating root or tuber material.

NONGRASS ANIMAL FEEDS (FORAGE, FODDER, STRAW, AND HAY)

Alfalfa, bean (velvet), clover, kudzu, lespedeza, lupin, sainfoin, trefoil, vetch (including crown vetch and milk vetch).

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

Important: Forage may not be grazed until 30 days after planting.

PEANUTS

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed.

ROOT AND TUBER VEGETABLES

Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet (garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley, parsnip, potato (0.08 fl. oz. only), radish (including oriental), rutabaga, salsify (oyster, black, and spanish), skirret, sweet potato, taniel, turmeric, turnip, yam bean (jicama, maonic pea), and true yam.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08-0.16 fl. oz. of Maxim 4FS per 100 lbs. of seed or propagating root or tuber material.

Note for Application to Potato: Maxim 4FS applications to potato seed pieces must be made only through specific types of equipment. Please contact your Syngenta Crop Protection seed treatment representative to obtain information on proper equipment for application.

Note for Potatoes Intended for Seed Production: As a resistance management strategy, if Maxim 4FS is used on potatoes intended for seed production, a labeled rate of mancozeb seed treatment dust must be applied to seed tubers after the Maxim 4FS treatment. If the mancozeb dust is not used, an in-furrow application of @ at 0.40 fl. oz /1000 ft. of row or Amistar™ at 0.13 fluid oz/1000 ft. of row must be applied over the Maxim treated seed tubers.

Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing.

SEED TREATMENT - EXPORT USE

FOR TREATMENT OF THE SEED FOR EXPORT ONLY TO COUNTRIES REQUIRING TREATMENT WITH MAXIM 4FS AND APRON XL LS BEFORE IMPORTATION. User should verify actual registered rate in country of destination and/or use.

Seed treated in accordance with the export use pattern must be labeled as follows:
TREATED SEED: FOR EXPORT ONLY – NOT FOR DOMESTIC SALE OR USE.

14/19

Maxim®, Apron XL®, Amistar™, Quadris® and the Syngenta logo are trademarks of a Syngenta Group Company

©200X Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.

Product of Switzerland
Formulated in USA

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP

Product ID.

HISTORIC FILE NAME: Z:LABELM/MAXIM 4FS 758A-L7G 0202 - wg - 3/01/02
MAX 4FS 758A L7G 0202 HILITED.doc - jda - 09/05/2002
MAX 4FS 758A-L7H 0103 nsp 1/10/03 new bklt size
MAX 4FS Draft aa-1-13-04 clean