

100-770

10/3/2001

1/19

D·z·n[®] diazinon

Lawn & Garden WBC
Water-Based Concentrate

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

Protects: Lawns, Vegetables, Fruits and Nuts, Roses, Flowers, Trees, and Shrubs

Controls: Ants, Aphids, Chinch Bugs, Cutworms, Deer Ticks (which may carry Lyme Disease), Fire Ants¹, Flea Beetles, Grubs, Mites, Sod Webworms... and many others as listed in the directions for use.

¹Aids in control.

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate..... 22.4%

Other Ingredients:..... 77.6%

Total: 100.0%

EPA Reg. No. 100-770 EPA Est. 70404-AL-001

**KEEP OUT OF REACH
OF CHILDREN.
CAUTION**

See additional precautionary statements and directions for use inside booklet.

SCP 770A-L1F 0901

1 quart
Net Contents

syngenta

ACCEPTED
OCT - 3 2001
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 100-770



Low odor -
Contains Syngenta Diazinon

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Repackaging and reformulation of this product is prohibited.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

GENERAL INFORMATION

D-z-n diazinon Lawn & Garden WBC is a water-based concentrate product which must be diluted with water before application. To avoid having spray mixture left over, mix only the amount needed. Refer to the section **How to Mix and Apply**. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. D-z-n Lawn & Garden WBC works fast and application should be repeated only as directed to maintain control.

Notes: (1) Do not use inside the home. (2) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia, or orchids, since injury may occur. (3) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (4) Do not apply in greenhouses.

Precautions: (1) *Wear a long-sleeved shirt and long-legged pants when spraying.* (2) *Spray with the wind to your back.* (3) *Do not spray on windy days.* (4) *If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.*

How to Mix and Apply

Step 1: Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Application should be repeated only as directed to maintain control.

Step 2: Use clean sprayer. Carefully measure and mix the amount of D-z-n diazinon Lawn & Garden WBC and water as indicated in the rate chart.

Step 3: Spray as directed in rate chart. Thorough coverage is important.

See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Step 4: Flush sprayer with clean water after each use.

D-z-n® diazinon Lawn & Garden WBC

3/19

Measurement Conversions

- 1 tablespoon = 3 teaspoons
- 1 fl. oz. = 6 teaspoons or 2 tablespoons
- 6 fl. oz. = $\frac{3}{4}$ cup
- $1\frac{1}{2}$ tablespoons in 3 gals. of water = $1\frac{1}{2}$ teaspoons in 1 gal. of water
- 2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

LAWN PEST CONTROL

Precautions

1. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl.
4. Avoid overlapping of sprays.
5. Where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least $\frac{1}{4}$ inch of water immediately after spraying this product; however, stop watering before puddling occurs.

Use on Bentgrass, Bermudagrass, Bluegrass, Dichondra, Fescue, Merion, and St. Augustinegrass.

To control pests listed below, apply recommended rates.

| Pests | Rate | | Remarks |
|---|--|-------------------------|---|
| | D-z-n | Water | |
| Deer Ticks (which may carry Lyme Disease) | $1\frac{1}{2}$ teaspoons per 125 sq. ft. or 2.25 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Spray grass and under shrubbery, particularly near house. Make the first application in spring when ticks are first noticed. Contact your local Cooperative Extension Service or Health Department for additional information about controlling ticks. |
| Lone Star Ticks | 1 teaspoon per 125 sq. ft. or 1.5 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Note: Do not apply this product to animals. |
| Lawn Chinch Bugs | $1\frac{1}{2}$ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Thoroughly wet down grass a few hours before applying. In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gals. of water covers 125 sq. ft. or 4 fl. oz. in 24 gals. of water per 1,000 sq. ft. |
| Ants, Armyworms, Bermudagrass Mites, Clover Mites, Spring-tails (<i>Collembola</i>), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs) | $1\frac{1}{2}$ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Thoroughly wet down grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active. Billbug control: Apply when activity is first observed or when chewed or brown grass is present, indicating damage from the insect. For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry. |
| Brown Dog Ticks, Chiggers, Fleas | $1\frac{1}{2}$ tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Note: Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house. |

D-z-n® diazinon Lawn & Garden WBC

4/19

| Pests | Rate | | Remarks |
|--|--|-------------------------|--|
| | D-z-n | Water | |
| Hyperodes Weevils | 1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application. |
| Millipedes | 1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Thoroughly wet grass a few hours before applying. |
| Rhodesgrass Scales | 1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Apply when crawlers first emerge. |
| White Grubs of Japanese Beetle, European Chafer, Southern Chafer | 1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft. | 3 gals. 24 gals. | Apply any time between late July and early October. Water grass thoroughly after application. |

| Nuisance Pests Outside the Home | Rate | | Remarks |
|--|-----------------------------------|---------|---|
| | D-z-n | Water | |
| Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (<i>Collembola</i>) | 6 fl. oz. or 12 tablespoons | 3 gals. | Thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide. To prevent entry into houses, spray a 5-ft. band of soil around the house next to the foundation, and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary. |
| Deer Ticks (which may carry Lyme Disease) | 1 1/2 teaspoons | 3 gals. | Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate. |
| Lone Star Ticks | 1 teaspoon | 3 gals. | |
| Fire Ants | 3 fl. oz. or 6 tablespoons | 3 gals. | To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness. Note: For best results, apply in cool weather 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear. |

1 fl. oz. = 6 teaspoons or 2 tablespoons

1 tablespoon = 3 teaspoons

1 teaspoon = 0.17 fl. oz.

D-z-n® diazinon Lawn & Garden WBC

5/19

INSECT CONTROL ON ORNAMENTAL TREES, SHRUBS, AND FLOWERS

Spray when listed pests first appear. Repeat as necessary.

| Trees and Shrubs | Insects | Rate per 1 Gal. of water | Timing/Special Directions |
|--|--|--------------------------|--|
| Arborvitae, Azalea, Birch, Boxwood, Camellia, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Scotch Pine, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellowwood, Yew | Aphids, Apple and Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly Oak), Hemlock Chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oak Loopers, Oblique-banded Leafrollers, Obscure Root Weevils, Omnivorous Leaf-tiers, Pear Slugs, Scale Crawlers (Cottony-cushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft), Tent Caterpillars, Webworms (Cotton-easter, Fall, Juniper, Mimosa), Whiteflies | 2 teaspoons | Apply as a thorough foliar spray. Wet leaves & branches to the dripping point. Try to penetrate dense foliage with spray. Mimosa Webworms: Apply when adults first appear. Make 2 additional applications at 4 to 5-week intervals. |
| Roses and Flowers | Insects | Rate per 1 Gal. of water | Timing/Special Directions |
| Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses | Aphids, Dipterous Leafminers, Euonymus Scale Crawlers, Flea Beetles, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies | 2 teaspoons | Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage. |

D-z-n® diazinon Lawn & Garden WBC

6/19

FRUITS AND NUTS

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

| Crop | PHI* |
|---------------|------|
| Apples | 21 |
| Apricots | 21 |
| Blackberries | 7 |
| Blueberries | 7 |
| Boysenberries | 7 |
| Cherries | 21 |
| Dewberries | 7 |
| Grapes | 28 |
| Loganberries | 7 |
| Nectarines | 21 |
| Peaches | 21 |
| Pears | 21 |
| Plums | 21 |
| Prunes | 21 |
| Raspberries | 7 |
| Strawberries | 5 |
| Walnuts | 45 |

*PHI = Pre-Harvest Interval, or the number of days to wait between the last application and harvest.

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

| Crop | Pests | Rate | Timing/Special Directions |
|----------|---|--------------------------------|---|
| Apples | Aphids, Apple Maggots, Codling Moths, Eyespotted Budmoths, Fruit Tree Leafrollers, Leafhoppers, Mealybugs, Scale Crawlers (San Jose Forbes), Spider Mites | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots may not be controlled by diazinon or similar type insecticides. |
| Apricots | Aphids, Apricot Mealybugs, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. |

D-z-n® diazinon Lawn & Garden WBC

7/19

| Crop | Pests | Rate | Timing/Special Directions |
|--|--|------------------------------------|--|
| Blueberries | Aphids, Blueberry Maggots, Cherry Fruitworms, Cranberry Fruitworms, Thrips, Two-spotted Spider Mites | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Allow a minimum of 14 days between applications. Do not apply more than 4 gals. per 500 sq. ft. of blueberry patch. Do not make more than 5 applications per season, a minimum of 14 days apart. Allow a minimum of 7 days between the last application and harvest. |
| Caneberries (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries | Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 5 gals. per 500 sq. ft. of caneberry patch. Do not make more than 5 applications per season, a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberries only in CA, OR, and WA. Do not use this product on caneberries grown in other states. |
| Cherries | Black Cherry Aphids, Cherry Fruitflies, Cherry Fruitworms (Maggots), Fruittree Leafrollers, Mites (Cherry Rust), Scale Crawlers (San Jose) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 7 days apart. Do not apply within 21 days of harvest. |
| Grapes | Aphids, Leafhoppers, Grape Berry Moths, Grape Leafrollers | 1 1/2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of home vineyard. Do not make more than 5 applications per season, a minimum of 7 days apart. Do not apply within 28 days of harvest. |
| Nectarines | Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. |
| Peaches | Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. |

D-z-n® diazinon Lawn & Garden WBC

8/19

| Crop | Pests | Rate | Timing/Special Directions |
|-------------------|--|----------------------------------|--|
| Pears | Aphids, Apple Maggots, Codling Moths, Mealybugs, Scale Crawlers, San Jose Forbes, Spider Mites, Tentiform Leafminers | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots will not be controlled by diazinon or similar type insecticides. |
| | Pear Psylla | 1-2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. |
| Plums and Prunes | Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red) | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 7 days apart. Do not apply within 21 days of harvest. |
| Strawberries | Aphids, Two-spotted Spider Mites, Strawberry Leafrollers | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of strawberry patch. Do not make more than 3 applications per season, a minimum of 7 days apart. Do not apply within 5 days of harvest. |
| Walnuts (CA only) | Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers | 2 teaspoons in 1 gal. of water | Apply as a full cover spray. Do not apply more than 6 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states. |

D-z-n® diazinon Lawn & Garden WBC

9/19

VEGETABLES

For foliar or soil applications, D-z-n Lawn & Garden WBC should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply in greenhouses.

Pre-Harvest Intervals (PHI*) for Foliar Applications

| Crop | PHI* (Days) |
|--|-------------|
| Beans ¹ (succulent varieties only: Lima, Pole, and Snap) | 7 |
| Broccoli | 7 |
| Cabbage | 21 |
| Cantaloupes | 3 |
| Carrots | 14 |
| Casabas | 3 |
| Cauliflower | 7 |
| Chinese Broccoli | 10 |
| Chinese Cabbage | 10 |
| Collards | 10 |
| Crenshaws | 3 |
| Cucumbers | 7 |
| Endive (Escarole) | 14 |
| Honeydew Melons | 3 |
| Kale | 10 |
| Lettuce | 14 |
| Muskmelons | 3 |
| Onions (Bulb and Green) | 14 |
| Parsnips | 14 |
| Peas ¹ | 7 |
| Peppers | 5 |
| Persian Melons | 3 |
| Potatoes (Irish) | 35 |
| Radishes | 14 |
| Rutabagas | 14 |
| Spinach | 14 |
| Squash, Summer | 7 |
| Squash, Winter | 3 |
| Sweet Corn | 7 |
| Tomatoes | 1 |
| Turnips (Roots and Tops) | 14 |
| Watermelons | 3 |

*PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

¹ Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas, or dried black-eyed peas.

D-z-n® diazinon Lawn & Garden WBC

10/19

VEGETABLES

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

| Crop | Pests | Rate | Timing/Special Directions |
|--|---|---|---|
| Beans: Lima, Pole, and Snap (succulent varieties only) | Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 5 days, up to 3 applications per season. Do not apply within 7 days of harvest. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms (snap beans only) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |

Notes: (1) Do not apply this product to dried bean varieties such as pinto beans or dried limas. (2) To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.

| | | | |
|--------------------|---|---|---|
| Beets, Red (Table) | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| Broccoli | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage-worms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Cabbage | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage-worms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 21 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |

D-z-n® diazinon Lawn & Garden WBC

4/19

| Crop | Pests | Rate | Timing/Special Directions |
|--|---|---|---|
| Cabbage (continued) | Root Maggots | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| Carrots | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| Cauliflower | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage-worms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Chinese Broccoli | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage-worms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Chinese Cabbage (Bok Choy and Napa) | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage-worms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |

D-z-n® diazinon Lawn & Garden WBC

12/19

| Crop | Pests | Rate | Timing/Special Directions |
|-----------------------|---|---|---|
| Collards | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage-worms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Corn (see Sweet Corn) | | | |
| Cucumbers | Aphids, Thrips, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest. |
| Endive (Escarole) | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Kale | Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage-worms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Lettuce | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |

D-z-n® diazinon Lawn & Garden WBC

13/19

| Crop | Pests | Rate | Timing/Special Directions |
|---|--|---|---|
| Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these) and Watermelons | Aphids, Thrips, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 3 days of harvest. |
| Onions (Bulb and Green) | Onion Thrips | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. |
| Parsnips | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Peas (succulent only) | Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 5 days, up to 3 applications per season. Allow at least 7 days between the last application and harvest. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| Peppers | Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 5 days of harvest. |
| Potatoes | Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 35 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |

Notes: (1) Do not apply this product to dried pea varieties such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom.

D-z-n® diazinon Lawn & Garden WBC

14/19

| Crop | Pests | Rate | Timing/Special Directions |
|----------------|--|---|---|
| Radishes | Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 3 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| Rutabagas | Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| Spinach | Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| Squash, Summer | Aphids, Thrips, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest. |
| Squash, Winter | Aphids, Thrips, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 3 days of harvest. |

D-z-n® diazinon Lawn & Garden WBC

15/19

| Crop | Pests | Rate | Timing/Special Directions |
|------------|--|---|---|
| Sweet Corn | Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Corn may be picked 7 days following last application. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Mole Crickets | 2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |

Note: To protect bees, do not apply to corn foliage during the pollen shed period.

| | | | |
|-------------------------|---|---|--|
| Tomatoes | Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (<i>Drosophila</i> spp.) | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 1 day of harvest. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |
| | Wireworms | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 4-8 inches. |
| Turnips and Turnip Tops | Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers | 1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden | Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides. |
| | Cutworms (surface, subterranean) | 6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water | Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms. |

STORAGE AND DISPOSAL

Storage

Store in original container in an area inaccessible to children and pets. If product freezes, place container in a warm room (68°F or above) inaccessible to children and pets and allow to thaw. Shake well before using.

Disposal

To avoid harming aquatic organisms in rivers and other surface waters, do not pour this product or rinse water into sanitary drains (for example, toilets, floor drains, and sinks) or into storm water sewers (for example, street drains). Do not reuse empty container. Triple rinse the empty container, pour the rinse water into the sprayer, then spray the area to be treated. Securely wrap empty container in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before smoking or eating. Remove contaminated clothing and wash before reuse. Do not use inside the home or inside greenhouses. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

| FIRST AID | |
|---|--|
| This product contains diazinon, an organophosphate insecticide that inhibits cholinesterase. | |
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious or convulsing person. |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| In the event of a major fire, spill, or other emergency, call 1-800-888-8372, day or night. | |
| NOTE TO PHYSICIAN | |
| This product contains diazinon, an organophosphate that inhibits cholinesterase. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should be administered only in conjunction with atropine. | |

D-z-n® diazinon Lawn & Garden WBC

17/19

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash water.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

D-z-n® and the Syngenta logo are trademarks of a Syngenta Group Company

©2001 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com
SCP 770A-L1F 0901

D·z·n® diazinon

Lawn & Garden WBC
Water-Based Concentrate

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

Protects: Lawns, Vegetables, Fruits and Nuts, Roses, Flowers, Trees, and Shrubs

Controls: Ants, Aphids, Chinch Bugs, Cutworms, Deer Ticks (which may carry Lyme Disease), Fire Ants¹, Flea Beetles, Grubs, Mites, Sod Webworms... and many others as listed in the directions for use.

¹Aids in control.

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 22.4%

Other Ingredients: 77.6%

Total: 100.0%

See directions for use in attached booklet.

EPA Reg. No. 100-770

EPA Est. 70404-AL-001

D·z·n® and the Syngenta logo are trademarks of a Syngenta Group Company
©2001 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, NC 27409
www.syngenta-us.com
SCP 770A-L1F 0901

KEEP OUT OF REACH OF CHILDREN.

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before smoking or eating. Remove contaminated clothing and wash before reuse.

Refer to label on back of container for Precautionary Statements (continued).

1 quart
Net Contents



Low odor -
Contains Syngenta Diazinon

Precautionary Statements (continued)

Do not use inside the home or inside greenhouses. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

First Aid

This product contains diazinon, an organophosphate insecticide that inhibits cholinesterase.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious or convulsing person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Hot Line Number: In the event of a major fire, spill, or other emergency, call 1-800-888-8372, day or night.

Note to Physician: This product contains diazinon, an organophosphate that inhibits cholinesterase. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should be administered only in conjunction with atropine.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas

may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash water.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Storage and Disposal**Storage**

Store in original container in an area inaccessible to children and pets. If product freezes, place container in a warm room (68°F or above) inaccessible to children and pets and allow to thaw. Shake well before using.

Disposal

To avoid harming aquatic organisms in rivers and other surface waters, do not pour this product or rinse water into sanitary drains (for example, toilets, floor drains, and sinks) or into storm water sewers (for example, street drains). Do not reuse empty container. Triple rinse the empty container, pour the rinse water into the sprayer, then spray the area to be treated. Securely wrap empty container in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.