

Ms. Carolyn Bussey
Ciba-Geigy Corporation
Ciba Plant Protection
P.O. Box 18300
Greensboro, NC 27419-8300

OCT - 8 1996

Dear Ms. Bussey:

Subject: Follow-up to Agency letter dated October 7, 1996
D.z.n diazinon AG600 WBC
EPA Reg. No. 100-784
Your submission dated September 20, 1996

As we discussed by telephone, when preparing the finished label for this product, please delete the pest "Flea Beetles" from the crop "Sweet Potatoes", since soil incorporation of this product is not appropriate as a control measure for this foliar pest. Thank you for your cooperation in this matter.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide Rodenticide Branch
Registration Division (7505C)

277

512636

OCT - 7 1996

Ms. Carolyn Bussey
Ciba-Geigy Corporation
Ciba Plant Protection
P.O. Box 18300
Greensboro, NC 27419-8300

Dear Ms. Bussey:

Subject: Amendment- Addition of low odor claim and logo and
"water based" logo

D.z.n diazinon AG600 WBC

EPA Reg. No. 100-784

Your submission dated September 20, 1996

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. The "LOW ODOR CIBA DIAZINON" logo may appear on the label in addition to (or in place of) the claim "Low Odor - Contains Ciba Diazinon", which is based upon use of the new process reduced odor diazinon manufacturing-use product, EPA Reg. No, 100-524. The "CIBA DIAZINON WATER BASED" logo may appear on this and any of your other water based diazinon formulations. Me-too registrants of water based diazinon products formulated using the new process reduced odor diazinon manufacturing-use product, EPA Reg. No, 100-524, may also add the claim "Low Odor - Contains Ciba Diazinon", with or without the associated logo, as well as the "water based" logo by Notification. The Agency

understands that Ciba will assist these me-too registrants, if necessary, by providing them a copy of Ciba's Agency approved labeling with the claim and logo(s) to accompany their state registration applications, since the me-too registrants' labeling will not have been stamped, in accordance with the Agency's Notification procedures.

- b. Under the PHI tables for Fruits and Nuts, for Vegetables-Foliar Application, and for Field Crops, in the definition of PHI revise "or the number of days between the last application and harvest" to read "or the number of days to wait between the last application and harvest."
 - c. On page 26, the pests Leafhoppers and Spider Mites are supposed to be associated with the crop Lima Beans. Add the missing horizontal line in the Crop column and add the crop name "Beans, Lima (succulent only)".
 - d. For the crops Broccoli, Brussels Sprouts, and Cabbage for consistency with the rest of the label, revise the Timing/Special Directions for Root Maggots to read "Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil."
 - e. For the crop Onions, for consistency with the rest of the label, revise the Timing/Special Directions for Onion Maggots to read "Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil."
 - f. For the crop Peas, please ensure that the printer does not add the extra carry over entries in the Crop, Pest, and Rate columns along with the Timing/Special Directions NOTES which did not fit on the previous page.
 - g. On page 63, correct the alignment problems with the column headings and the associated text for each column.
 - h. For both the jug label and the booklet label, in the Environmental Hazards add "On lawns," as the beginning of the sentence that now begins "If waterfowl (ducks or geese)".
3. Submit three (3) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section

6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide Rodenticide Branch
Registration Division (7505C)

5/77

RESTRICTED USE PESTICIDE
DUE TO AVIAN AND AQUATIC TOXICITY
FOR RETAIL SALE TO AND USE BY CERTIFIED APPLICATORS
OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND
ONLY FOR THOSE USES COVERED BY THE CERTIFIED
APPLICATOR'S CERTIFICATION.

(Booklet for 2 1/2 Gallon Container)

D•z•n® diazinon AG600 WBC

INSECTICIDE

Water-Based Concentrate

For control of certain insects on fruits, nuts, vegetables, field crops,
lawns, and ornamentals

This product must not be used on golf courses and sod farms.

Low Odor - Contains Ciba Diazinon

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-
methyl-4-pyrimidinyl) phosphorothioate56.0%

Inert Ingredients:44.0%

Total:100.0%

Shake contents well before using.

EPA Reg. No. 100-784

EPA Est. 100-AL-1

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside
booklet.

2 1/2 Gallons
U.S. Standard Measure

CGA 1L145 (DRAFT)

[QUARK/DZN DIAZINON AG600 WBC/DZN DIAZ AG600WBC-A] - CCG - 9/20/96

ACCEPTED
with COMMENTS
in EPA Letter Dated

OCT - 7 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

100-784

96 SEP 24 P 2:40

RECD EPA/OP/DPD1

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire **Directions for Use** and the **Conditions of Sale and Warranty** before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The **Directions for Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use** subject to the inherent risks referred to above. **Ciba-Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.** Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing **Conditions of Sale and Warranty**, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.

General Information

D•z•n diazinon AG600 WBC is a water-based concentrate, when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops and other plants. Do not use on food crops grown in greenhouses. Do not use in barns or dwellings. Do not use on livestock.

In the directions for use which follow, best control is obtained when application is made at first sign of pests. Application should be repeated only as directed to maintain control.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not permit children or pets to go onto treated areas until sprays have dried.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Work Safety Rules

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

When handling D•z•n diazinon AG600 WBC, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See **Practical Treatment** section of this label.

Note: The California Department of Agriculture has set a 5-day reentry period for diazinon on grapes, peaches, and nectarines. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of 2 or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

CHEMIGATION

APPLICATION THROUGH AN IRRIGATION SYSTEM (CHEMIGATION) - CRANBERRIES ONLY

D•z•n diazinon AG600 WBC may be applied alone or in combination with other pesticides registered for application through sprinkler irrigation systems to cranberries. To insure compatibility, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for 5 minutes. If the combination remains mixed, or can be remixed readily, the mixture is compatible.

Do not apply this product through any type of irrigation system to other crops. For chemigation, apply this product to cranberries only through sprinkler irrigation systems. Do not apply this product to cranberries through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect the irrigation system used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

Chemigation System Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Sprinkler Chemigation

1. The system must contain a functional check-valve, a vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Operating Instructions

1. Determine the acreage to be treated by the sprinkler system.
2. Measure the appropriate amount of D•z•n diazinon AG600 WBC needed to treat the acreage to be treated. Refer to the specific directions for use for cranberries for the application rate and the amount of water per acre.
3. Add the pre-measured D•z•n diazinon AG600 WBC to the chemigation system supply tank and dilute with water if needed. Proper agitation is needed if D•z•n diazinon AG600 WBC is tank mixed with other pesticides registered for sprinkler chemigation application to cranberries.
4. Start the irrigation system and bring the system up to full pressure, making certain that all sprinkler heads are functioning properly.
5. Engage the chemigation injection or venturi system to add the D•z•n diazinon AG600 WBC to the irrigation lines.
6. Continue to run irrigation system for a few minutes after supply tank is empty to allow material in the irrigation lines to reach the sprinkler heads furthest from the chemigation injection point.

Fruit and Nut Crops

REI - 24 hours

D•z•n diazinon AG600 WBC may be applied using ground or aerial application equipment as specified in the following table. This table indicates the **minimum** amount of water that can be used for the various methods of application. The rate of D•z•n diazinon AG600 WBC for aerial spray or concentrated spray applied to fruit trees by ground equipment is based on a dilute full cover spray applied with conventional ground equipment at a rate of 300-400 gals. of water per acre. The specific amount of product to be used appears under the separate directions for use for each crop. If the **Rate** calls for 13½ fl. oz. of D•z•n diazinon AG600 WBC in 100 gals. of water, this amounts to 40½-54 fl. oz. of product per acre when applied at the usual dilute ground spray volumes of 300-400 gals. of water per acre. In no case should the amount of product used per acre exceed the maximum amount per acre per application that is specified in the **Timing/Special Directions** for individual crops.

Crop	Aerial Application Minimum Gals./Acre	Ground Application Minimum Gals./Acre	
		Dilute	Concentrate
Almonds*	20	100	20
Apricots	20	100	20
Blackberries**	20	100	20
Blueberries	20	100	20
Boysenberries**	20	100	20
Cherries	20	100	20
Cranberries	5	15	—
Dewberries**	20	100	20
Grapes	20	100	20
Loganberries**	20	100	20
Nectarines	20	100	20
Peaches	20	100	20
Plums	20	100	20
Prunes	20	100	20
Raspberries**	20	100	20
Strawberries	20	100	20
Walnuts*	20	100	20

*CA only.

**CA, OR, or WA only.

Observe stated time intervals between last application and harvest, as well as intervals between applications and total amount of product that can be applied per season.

Crop	PHI*
Almonds	**
Apricots	21
Blackberries	7
Blueberries	7
Boysenberries	7
Cherries	21
Cranberries	7
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums, Prunes	21
Raspberries	7
Strawberries	5
Walnuts	45

*PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

** = Dormant spray only.

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds in the orchard are in bloom.

Crop	Pest	Rate	Timing/Special Directions
Almonds (CA only)	San Jose Scale, Parlatoria Scale	13 $\frac{1}{2}$ -20 $\frac{1}{2}$ fl. oz. + 2-3 gals. dormant oil or 1-1 $\frac{1}{2}$ gals. superior type oil in 100 gals. of water	Apply as a dormant spray. Do not apply more than 82 oz. of diazinon or 9 gals. of oil per acre.
Apricots	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers	13 $\frac{1}{2}$ fl. oz. in 100 gals. of water	Apply as infestations occur. Repeat application after 14 days, if necessary.
	Olive Scale Crawlers	6 $\frac{1}{2}$ fl. oz. + 1 $\frac{1}{2}$ gals. of light medium horticultural oil per 100 gals. of water	Apply when scale crawlers are present.
	Apricot Mealybugs	13 $\frac{1}{2}$ fl. oz. oz. per 100 gals. of water	Apply as a cover spray from petal fall to June. Allow 14 days between applications.

Notes: (1) Do not apply more than 54 fl. oz. of product per acre per application and no more than 162 fl. oz. of product per acre per season.
(2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Blue-berries	Cranberry fruitworms, Cherry fruitworms, Blueberry maggots, Aphids, Thrips, Two-spotted Spider Mites	13 1/2 fl. oz. per 100 gals. of water	Apply as infestations occur. Allow 14 days between applications.
	Fire Ants	13 1/2 fl. oz. per 100 gals. of water	To aid in the control of fire ants, slowly apply 1 gal. of dilute mixture over and 6 inches around each fire ant mound. Apply gently to avoid disturbing ants. Use equipment capable of delivering the diluted product as a gentle rain. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness. Note: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours.

Notes: (1) Do not apply more than 27 fl. oz. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Cane-berries CA, OR, and WA only (Black-berries, Boysen-berries, Dew-berries, Logan-berries, Rasp-berries)	Leaf-hoppers, Thrips, Aphids, Two-spotted Spider Mites, Raspberry Fruitworms, Raspberry Sawflies	27 fl. oz./A	<p>Apply as insects occur. Repeat application after 14 days if necessary. Do not make more than 5 applications per season. Do not apply within 7 days of harvest, or illegal residues may occur.</p> <p>Apply in a minimum of 100 gals. of water per acre (200 gals./A maximum).</p> <p>For Raspberry Fruitworms, make one application when blossom buds separate. Make a second application immediately before blossoms open.</p>
	Dryberry Mites	27-54 1/2 fl. oz./A	<p>Apply as a dormant spray in the early Spring when canes are still on the ground and buds are closed. Make a second application when buds are well developed, but still closed.</p> <p>Apply in a minimum of 100 gals. of water per acre. Do not apply within 7 days of harvest, or illegal residues may occur.</p>

Crop	Pest	Rate	Timing/Special Directions
Cane-berries (Contd)	Raspberry Crown Borers (Raspberry Root Borers)	54½ fl. oz./A	Apply in a single application in a minimum of 100 gals. of water per acre as a drench to the crown and lower canes in the Spring before buds break.
Cherries	Eyespotted Bud Moths, Fruit Tree Leafrollers	13½ fl. oz. in 100 gals. of water	Apply in cover sprays as infestations occur beginning at petal fall. Repeat application after 7 days, if necessary.
	Leafhoppers	13½ fl. oz. in 100 gals. of water	Make 3 applications beginning at petal fall and continuing through the growing season as infestations occur. Allow 7 days between applications.
	Cherry Fruitflies	6½-13½ fl. oz. in 100 gals. of water	Make 3-4 applications beginning when adult flies begin to emerge and repeating at 10 day intervals.
	Black Cherry Aphids	13½ fl. oz. in 100 gals. of water	Apply as a cover spray when aphids appear. Repeat application after 10 days if necessary.
	San Jose Scale Crawlers	13½ fl. oz. in 100 gals. of water	Apply during the pre-pink stage or when crawlers first appear.
	Cherry Rust Mites	6½-13½ fl. oz. in 100 gals. of water	Apply to trees as a single supplemental spray after harvest.

Notes: (1) Do not apply more than 54 fl. oz. of product per acre per application and no more than 162 fl. oz. of product per acre per season.
(2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Cranberries	Blackheaded Fireworms (<i>Rhopobota naevana</i>)	54½ fl. oz./A	Apply to larval stage. Pheromone trap captures and sweep monitoring may be used to optimize treatment timing. Repeat application after 14 days, if necessary. A maximum of 6 applications per season is permitted.
	Cranberry Tipworms	54½ fl. oz./A	Presence of damage is not an indicator of active infestation. For best results, target early-stage larvae. Repeat application after 14 days, if necessary. A maximum of 6 applications is permitted.
	Cranberry Fruitworms	54½-82 fl. oz./A	Cranberry growth stage and/or berry inspection for eggs may be used to optimize treatment timing; consult your local extension agent or pest management advisor. Repeat application every 14 days, as needed. A maximum of 4 applications per season is permitted for the 82 fl. oz. rate. A maximum of 6 applications per season is permitted for the 54½ fl. oz. rate.

Notes: (1) This product may be applied to cranberries via ground, aerial, or sprinkler irrigation. For ground application, apply the product in a minimum of 15 gals. of water per acre, for aerial application, a minimum of 5 gals. of water per acre, for chemigation up to 400 gals. of water per acre. Refer to the **Chemigation** section of this label for specific chemigation directions. (2) Do not apply more than 328 fl. oz. of product per acre per season. (3) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Grapes	Pacific Spider Mites, Leafhoppers, Grape Berry Moths, Grape Leaf-folders, Omnivorous Leafrollers, Aphids, Spider Mites	13 1/2-27 fl. oz./A in a minimum of 100 gals. of water	Apply as pest occurs as a thorough cover spray. Repeat application after 7 days, if necessary.
	<i>Drosophila</i> spp.	5 fl. oz./A in a minimum of 100 gals. of water (300 gals. of water maximum)	Apply as pest occurs as a thorough cover spray. Repeat application after 7 days, if necessary.

Notes: (1) Do not apply more than 27 fl. oz. of product per acre per application and no more than 135 fl. oz./A per season. (2) To avoid illegal residues, allow 28 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Nectarines	Oriental Fruit Moths	13½ fl. oz. in 100 gals. of water	Apply at 100% petal fall. Make 2 additional applications at 10 to 15-day intervals. Time the last application to peak adult flight.
	Aphids, Brown Mites (Clover Mites), Two-Spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers, Peach Twig Borers	13½ fl. oz. in 100 gals. of water	Apply as infestations occur. Repeat application after 14 days, if necessary.

Notes: (1) Do not apply more than 54 fl. oz. of product per acre per application and no more than 162 fl. oz./A per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Peaches	Aphids, Brown Mites (Clover Mites), Two-Spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers	13½ fl. oz. in 100 gals. of water	Apply as infestations occur. Repeat application after 14 days, if necessary.
	Oriental Fruit Moths	13½ fl. oz. in 100 gals. of water	Apply at 100% petal fall. Make 2 additional applications at 10 to 15-day intervals. Time the last application to peak adult flight.
	Peach Twig Borers	13½ fl. oz. in 100 gals. of water	Apply as insects occur in May and June. Allow 7 days between applications.
	Leafhoppers	13½ fl. oz. in 100 gals. of water	Make 3 applications beginning at petal fall and continuing through the growing season. Allow 7 days between applications.
	White Peach Scale	20½-27 fl. oz. in 100 gals. of water	Apply post-harvest timed to coincide with peak crawler and immature scale activity. Under conditions of heavy infestation, 2 applications (7 days apart) may be needed. Do not apply before fruit is harvested.

Notes: (1) Do not apply more than 54 fl. oz. of product per acre per application and no more than 162 fl. oz. of product per acre per season.
(2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Plums	Brown Mites (Clover Mites), European Red Mites	13½ fl. oz. in 100 gals. of water	Apply, as needed, every 7 days.
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	6½-13½ fl. oz. in 100 gals. of water	

Notes: (1) Do not apply more than 54 fl. oz. of product per acre per application and no more than 162 fl. oz. of product per acre per season.
 (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Prunes	Brown Mites (Clover Mites), European Red Mites	13½ fl. oz. in 100 gals. of water	Apply, as needed, every 7 days.
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	6½-13½ fl. oz. in 100 gals. of water	

Notes: (1) Do not apply more than 54 fl. oz. of product per acre per application and no more than 162 fl. oz. of product per acre per season.
 (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Straw-berries	Ahpids, Two-Spotted Spider Mites	13½ fl. oz. in 100 gals. of water per acre	Apply as pests occur. Repeat application, if necessary, every 7 days, but do not make more than 3 applications per season.
	Cyclamen Mites	27 fl. oz. in 100 gals. of water per acre	Direct spray to the plant crowns. Plants should be sufficiently agitated to assure thorough coverage of the foliage. Repeat application after 7 days, if necessary. Do not make more than 3 foliar applications per season.
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast in sufficient water to obtain even coverage and immediately incorporate into the top 1-2 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Strawberries (Contd)	Strawberry Leafrollers	10-13½ fl. oz. in 100 gals. of water	Apply 200 gals. of spray per acre when blossoms show color.

Notes: (1) Do not apply more than 27 fl. oz./A per application. (2) Do not apply more than 108 fl. oz. of product per acre per season. (3) To avoid illegal residues, allow a minimum of 5 days between the last application and harvest.

Walnuts (CA only)	Aphids, Spider Mites	27 fl. oz./A	Apply as a foliar spray. Repeat application in 14 days, if necessary. Do not make more than 3 applications per season. Do not apply within 45 days of harvest or after husks open, or illegal residues may occur. Make ground applications in a minimum of 100 gals. of water per acre.
	Codling Moths, Scale Crawlers, Walnut Caterpillars	82 fl. oz./A	

Vegetable Crops

REI - 24 hours

For foliar or soil applications, D•z•n diazinon AG600 WBC should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. D•z•n diazinon AG600 WBC may be applied using ground or aerial application equipment as specified in the following table. This table indicates the **minimum** amount of water that can be used for the various methods of application. The specific amount of product to be used appears under the separate directions for use for each crop.

Crop	Ground Application		Aerial Application Minimum Gals./Acre
	Minimum Gals./ Acre	OR Minimum Gals./1,000 Sq. Ft.	
Beans (Lima, Pole, and Snap) (succulent only)	10	2	5
Beets, Red (Table)	10	2	5
Broccoli	10	2	5
Brussels Sprouts	10	2	5
Cabbage	10	2	5
Cantaloupes	5	1	5
Carrots	10	2	5
Casabas	5	1	5
Cauliflower	10	2	5
Chinese Broccoli	10	2	5
Chinese Cabbage	10	2	5
Chinese Mustard	10	2	5
Chinese Radish (CA, FL only)	10	2	5
Collards	10	2	5
Crenshaws	5	1	5
Cucumbers	5	1	5
Endive (Escarole)	10	2	5
Ginseng	10	2	5
Honeydew Melons	5	1	5
Kale	10	2	5
Lettuce, Head	10	2	5
Lettuce, Leaf	10	2	5
Muskmelons	5	1	5
Mustard	10	2	5
Onions (Bulb and Green)	10	2	5
Parsley	10	2	5
Parsnips	10	2	5
Peas (succulent only)	10	2	5
Peppers	10	2	5
Persian Melons	5	1	5
Potatoes (Irish)*	10	2	5
Radishes	10	2	5
Spinach	10	2	5
Squash, Summer	5	1	5
Squash, Winter	5	1	5
Sweet Corn	20	4	5
Sweet Potatoes	10	2	5

(Contd)

Crop	Ground Application		Aerial Application Minimum Gals./Acre
	Minimum Gals./ Acre	OR Minimum Gals./1,000 Sq. Ft.	
Swiss Chard	10	2	5
Tomatoes	10	2	5
Turnips	10	2	5
Watermelons	5	1	5

*Do not use on commercially grown potatoes which will be hand-harvested.

Observe stated time intervals between last application and harvest, as well as intervals between and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (PHI*) for Foliar Applications

Crop	PHI* (Days)
Beans (Lima, Pole, and Snap)	7
Beets, Red (Table)	14
Broccoli	7
Brussels Sprouts	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casabas	3
Cauliflower	7
Chinese Broccoli	10
Chinese Cabbage	10
Chinese Mustard	10
Chinese Radish	10
Collards	10
Crenshaws	3
Cucumbers	7
Endive (Escarole)	14
Ginseng	30
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Mustard	10
Onions (Bulb and Green)	14
Parsnips	14
Peas	7
Peppers	5
Persian Melons	3
Potatoes (Irish)	35
Radishes	14
Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Swiss Chard	14
Tomatoes	1
Turnips, Roots and Tops	14
Watermelons	3

*PHI - Pre-Harvest Interval or the number of days between the last application and harvest.

Notes: (1) **Soil Incorporation:** Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see **Timing/Special Directions**) using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches, for subterranean cutworms incorporate 3-6 inches. (2) **To Protect Bees:** Foliar applications of this product should not be made to peas or beans if these crops or weeds in the treatment area are in bloom, or to corn during the pollen shed period.

Crop	Pest	Rate	Timing/Special Directions
Beans (Lima, Pole, and Snap) (succu- lent only)	Cutworms	54 ¹ / ₂ -109 ¹ / ₂ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109 ¹ / ₂ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Black Bean Aphids, Mexican Bean Beetles, Dipterous Leafminers	13 ¹ / ₂ -20 ¹ / ₂ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season.
	Cucumber Beetles	10 fl. oz./A	Do not apply within 7 days of harvest, or illegal residues may occur. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay. Make cucumber beetle application in 100 gals. of water per acre.

Crop	Pest	Rate	Timing/Special Directions
Beans (Lima, Pole and Snap) (succulent only) (Cont'd)			<p>Notes: 1) Diazinon will not control organosphosphate-resistant leafminers. 2) To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.</p>
	Leafhoppers, Spider Mites	13 1/2 fl. oz./A	<p>Apply as pests occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season.</p> <p>Do not apply within 7 days of harvest, or illegal residues may occur.</p> <p>Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.</p> <p>Note: To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.</p>

Crop	Pest	Rate	Timing/Special Directions
Beets, Red (Table)	Cutworms	5 1/2 - 10 1/2 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109 1/2 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Dipterous Leafminers	6 1/2 - 13 1/2 fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 14 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>

Crop	Pest	Rate	Timing/Special Directions
Broccoli	Root Maggots	54½-82 fl. oz./A	Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil.
		3-6½ fl. oz. per 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 3-6½ oz. per plant by hand or by tractor mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)

Crop	Pest	Rate	Timing/Special Directions
Broccoli (Contd)	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage- worms, Dipterous Leafminers	6½-13½ fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 7 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>

Crop	Pest	Rate	Timing/Special Directions
Brussels Sprouts	Root Maggots	54½-82 fl. oz./A	Apply to plant beds or fields before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil.
		3-6½ fl. oz. in 50 gals. of water	<p>In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 3-6½ fl. oz. per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant.</p> <p>Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.</p>
	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)

Crop	Pest	Rate	Timing/Special Directions
Brussels Sprouts (Contd)	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage-worms	6½-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest, or illegal residues may occur.

Crop	Pest	Rate	Timing/Special Directions
Cabbage	Root Maggots	54½-82 fl. oz./A	Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil.
		3-6½ fl. oz. in 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 3-6½ fl. oz. per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)

Crop	Pest	Rate	Timing/Special Directions
Cabbage (Contd)	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	6½-13½ fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 21 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>

Crop	Pest	Rate	Timing/Special Directions
Carrots	Cutworms	54 $\frac{1}{2}$ -109 $\frac{1}{2}$ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109 $\frac{1}{2}$ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids	13 $\frac{1}{2}$ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest, or illegal residues may occur.

Crop	Pest	Rate	Timing/Special Directions
Cauliflower	Root Maggots	54½-82 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		3-6½ fl. oz. in 50 gals. of water	<p>In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 3-6½ fl. oz. per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant.</p> <p>Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.</p>
	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)

Crop	Pest	Rate	Timing/Special Directions
Cauliflower (Contd)	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage- worms, Dipterous Leafminers	6½-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications. Do not apply within 7 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Chinese Broccoli	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	6 1/2-13 1/2 fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 10 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>
Chinese Cabbage (Bok Choy and Napa)	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	6 1/2-13 1/2 fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 10 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>

Crop	Pest	Rate	Timing/Special Directions
Chinese Mustard (Gai Choy)	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	6 $\frac{1}{2}$ -13 $\frac{1}{2}$ fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 10 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>
Chinese Radish (Japanese Radish or Daikon) (FL and CA only)	Aphids, Diamondback Moths, Imported Cabbage-worms, Dipterous Leafminers	13 $\frac{1}{2}$ fl. oz./A	<p>Mix in a minimum of 100 gals. of water per acre and apply as a foliar spray as pests occur. Repeat application, as necessary, every 14 days. Do not make more than 3 applications per season.</p> <p>Do not apply within 10 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>

Crop	Pest	Rate	Timing/Special Directions
Collards	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbage-worms, Harlequin Bugs, Corn Earworms	6½-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest, or illegal residues may occur.

Crop	Pest	Rate	Timing/Special Directions
Cucumbers	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms	13½ fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 10 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>

Crop	Pest	Rate	Timing/Special Directions
Endive (Escarole)	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	6½-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Ginseng	Leafhoppers, Aphids, Lygus Bugs, Flea Beetles, Jumping Plant Lice	10-13½ fl. oz./A	Spray when insects first appear. For heavy insect infestations, use 13½ fl. oz. Apply by ground equipment using 10- 100 gals. of water per acre. Do not apply more than one application per growing season. Do not apply during flowering of 3 or 4 year old crops. Do not apply within 30 days of harvest. Do not graze treated areas or feed treated forage to livestock.
Kale	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Kale (Contd)	Aphids, Diamondback Moths, Imported Cabbage- worms, Harlequin Bugs, Corn Earworms	6 1/2-13 1/2 fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest, or illegal residues may occur.
Lettuce	Cutworms	54 1/2-109 1/2 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109 1/2 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	6 1/2-13 1/2 fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate- resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these) and Watermelons	Cutworms	54 ¹ / ₂ -109 ¹ / ₂ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109 ¹ / ₂ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	6 ¹ / ₂ -20 ¹ / ₂ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Mustard	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers, Diamondback Moths, Imported Cabbage- worms	6½-13½ fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 10 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>

Crop	Pest	Rate	Timing/Special Directions
Onions (Bulb and Green)	Onion Maggots	54 ¹ / ₂ -109 ¹ / ₂ fl. oz./A	Broadcast just before planting and mix into the top 3-4 inches of soil. Note: Diazinon will not control organophosphate-resistant onion maggots
	Onion Thrips	13 ¹ / ₂ fl. oz./A	Apply as thrips appear. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest, or illegal residues may occur.
	Wireworms	82-109 ¹ / ₂ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Parsley	Cutworms	54 ¹ / ₂ -109 ¹ / ₂ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109 ¹ / ₂ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Parsnips	Aphids, Flea Beetles, Dipterous Leafminers	6½-13½ fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 14 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>
Peas (succulent only)	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	10-13½ fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season.</p> <p>To avoid illegal residues, allow a minimum of 7 days between the last application and harvest.</p> <p>Vines and hay may be fed to dairy and beef cattle and sheep. To avoid illegal residues, wait 7 days following treatment if forage is to be cut for hay.</p>

Crop	Pest	Rate	Timing/Special Directions
Peas (succulent only) (Contd)	Aphids, Dipterous Leafminers	10-13½ fl. oz./A	Notes: (1) Diazinon will not control organophosphate resistant leafminers. (2) To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.
Peppers	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Serpentine Leafminers	6½-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 5 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Potatoes	Cutworms (Surface and Subterranean)	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Southern Potato Wireworms (South- eastern U.S.)	54-109½ fl. oz./A	For potatoes grown on mineral soils in the Southeast, apply before planting and immediately incorporate into the top 3-6 inches of soil, depending on location of wireworms.
	Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leafminers	6½-10 fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 35 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate- resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Potatoes (Cont'd)	Banded Cucumber Beetles, Leafhoppers, Southern Armyworms	10-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 35 days of harvest, or illegal residues may occur.

Note: Do not use on commercially grown potatoes which will be hand-harvested.

Radishes	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Flea Beetles, Dipterous Leafminers	6½-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate- resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Rutabagas	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Flea Beetles, Dipterous Leafminers	6½-13½ fl. oz./A	<p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 14 days of harvest, or illegal residues may occur.</p> <p>Note: Diazinon will not control organophosphate-resistant leafminers.</p>

Crop	Pest	Rate	Timing/Special Directions
Spinach	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	6½-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
Squash, Summer	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Squash, Summer (Contd)	Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	6 $\frac{1}{2}$ -20 $\frac{1}{2}$ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate- resistant leafminers.
Squash, Winter	Cutworms	54 $\frac{1}{2}$ -109 $\frac{1}{2}$ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109 $\frac{1}{2}$ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	6 $\frac{1}{2}$ -20 $\frac{1}{2}$ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate- resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Sweet Corn	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Seed Corn Maggots	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast before planting and immediately incorporate into the top 4-8 inches of soil.
	Corn Earworms	27-34 fl. oz./A	Apply when silks first appear. Repeat application every 7 days, if necessary. Do not make more than 5 applications per season.
	Corn Rootworm (Adults)	6½-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Corn Leaf Aphids	13½-27 fl. oz./A	Apply in sufficient water to provide complete coverage to the upper portions of the plants. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.

Crop	Pest	Rate	Timing/Special Directions
Sweet Corn (Contd)	Grasshoppers, Spider Mites, Flea Beetles	13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Sap Beetles	27-34 fl. oz./A	Consult local agricultural authorities for proper timing of sprays. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.

Notes: (1) To protect bees, do not apply foliarly to corn during the pollen shed period. (2) Allow 7 days after the last application before grazing or cutting for forage. Forage may be fed to beef and dairy cattle and sheep. (3) Corn may be picked 7 days after the last application.

Sweet Potatoes	Wireworms, Flea Beetles	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Swiss Chard	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.

Crop	Pest	Rate	Timing/Special Directions
Swiss Chard (Contd)	Aphids, Imported Cabbage- worms, Diamondback Moths, Harlequin Bugs	6½-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest, or illegal residues may occur.
Tomatoes	Cutworms	54½-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	6½ fl. oz./A	Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Banded Cucumber Beetles, Fall Army- worms, Southern Armyworms, Beet Armyworms	10-13½ fl. oz./A	Do not apply within one day of harvest, or illegal residues may occur.
	Vinegar Flies (<i>Drosophila</i> spp.)	6½-20½ fl. oz./A	Note: Diazinon will not control organophosphate- resistant leafminers.

Crop	Pest	Rate	Timing/Special Directions
Turnips, Roots and Tops	Cutworms	54 $\frac{1}{2}$ -109 $\frac{1}{2}$ fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	27 fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	82-109 $\frac{1}{2}$ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Flea Beetles, Dipterous Leafminers	6 $\frac{1}{2}$ -13 $\frac{1}{2}$ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate- resistant leafminers.

Field Crops

REI - 24 hours

Soil and foliar applications of D•z•n diazinon AG600 WBC should always be applied as a spray in sufficient water to ensure thorough coverage. For foliar applications, observe stated time intervals between last application and harvest, as well as intervals between applications and total number of applications that can be made per season.

Pre-harvest Interval (PHI*) for Foliar Applications

Crop	PHI* (Days)
Hops	14
Sugar Beets	14

*PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Hops	Aphids, Spider Mites	27 fl. oz./A	Apply as pests occur. Repeat application, as necessary, every 14 days. Do not make more than 4 applications per season. Do not apply within 14 days of harvest.

Crop	Pest	Rate	Timing/Special Directions
Sugar Beets	Wireworms	82-109½ fl. oz./A	Broadcast just before planting. Incorporate into soil 4-6 inches. Do not apply more than 109½ fl. oz. of this product to the soil, per season.
	Aphids, Beet Leafhoppers, Dipterous Leafminers	10-13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
	Grasshoppers	13½ fl. oz./A	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest.

Note: Sugar beet tops may be fed to beef and dairy animals.

Grassland Insects

Ditch Banks, Roadsides, Wasteland, Noncrop Areas, Barrier Strip(s)

Grasshoppers: Spray 10-13½ fl. oz./A in water (minimum of 1 gal. of water per acre) for aerial applications or a minimum of 5 gals. of water per acre for ground applications or in oil (a minimum of 1 gal. of oil per acre) when insects first appear, preferably in the nymphal stage. Thorough coverage of the foliage is essential.

Do not graze or use treated grass for forage or feed.

Insect Control on Ornamentals

REI - 12 hours

To control certain insects on:

Arborvitae	Maple
Azalea	Marigold
Birch	Oak
Boxwood	Pansy
Camellia	Petunia
Carnation	Pine
Chrysanthemum	Ornamental Plum
Crab Apple	Poplar
Dogwood	Rhododendron
Douglas Fir	Rose
Elm	Scotch Pine
Euonymus	Spirea
Gladioli	Spruce
Hawthorn	Sycamore
Holly	Wax Plant (Hoya)
Honey Locust	Willow
Honeysuckle	Yellowwood
Juniper	Yew
Lilac	
Lily	
Locust	

Apply the recommended rates indicated below. Spray when listed pests first appear. Try to spray underside of leaves and penetrate dense foliage. Repeat as necessary.

Insects	Amount of D·z·n diazinon AG600 WBC to Use	
	Rate per 3 Gals. Water	Rate per 100 Gals. Water
Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Dipterous Leafminers, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Obscure Root Weevils, Omnivorous Leafiers, Privet Mites, Scale Crawlers (Cottonycushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Two-spotted Spider Mites, Juniper Webworms, Whiteflies	2 1/2 teaspoons	13 1/2 fl. oz.
Apple-and-Thorn Skeletonizers, Cotoneaster Webworms, European Pine Sawfly, Fall Webworms, Hemlock Chermes, Oak Loopers, Obliquebanded Leafrollers, Pear Slugs, Tent Caterpillars	2 1/2 tablespoons	41 fl. oz.

Precaution: Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, and Gardenia, since injury to the plants may occur.

Mimosa Webworms: Mix 2½ teaspoons in 3 gals. of water (or 13½ fl. oz. in 100 gals. of water) and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

Lawn Pest Control (Excluding Golf Courses and Sod Farms)

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping of sprays. If waterfowl (ducks or geese) can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least ¼ inch of water immediately after application; however, stop watering before puddling occurs.

To control pests listed below, apply recommended rates.

Pest	Rate/1,000 sq. ft.		Remarks
	Diazinon AG600 WBC	Water	
Lawn Chinch Bugs	3 1/2-5 table- spoons	3 gals.	Use higher rate for for longer residual control and in lawns of dense growth, such as St. Augustine lawns.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs), Springtails (Collembola)	3 1/2-5 table- spoons	3 gals.	Spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk when wasps are not active. For frit flies, before treatment, mow grass and water well, but delay application until grass is dry. For billbug control, apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect.
Brown Dog Ticks, Chiggers, Fleas	5 table- spoons	3 gals.	Note: Do not apply to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.

BEST AVAILABLE COPY

Pest	Rate/1,000 sq. ft.		Remarks
	D•z•n diazinon AG600 WBC	Water	
Hyperodes Weevils	5 tablespoons	3 gals.	Treat problem areas in mid-April and again in mid-May.
Millipedes	5 tablespoons	3 gals.	
Rhodesgrass Scales	5 tablespoons	3 gals.	Apply when crawlers first emerge.
White Grubs (such as Japanese Beetle Larvae, European Chafer, Southern Chafer)	5 tablespoons	3 gals.	Apply any time between late July and early October.

Nuisance Pests in Outside Areas

- | | |
|-----------------|-----------------------------------|
| Ants | Fleas |
| Boxelder Bugs | Flies |
| Brown Dog Ticks | Millipedes |
| Chiggers | Sowbugs (Pillbugs) |
| Clover Mites | Spiders |
| Cockroaches | Springtails (<i>Collembola</i>) |
| Crickets | |
| Earwigs | |

Mix 5 tablespoons in 3 gals. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.

To prevent entry of insects into houses, spray a 5-ft. band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 ft.

Repeat application as necessary. **Do not apply to animals. Do not treat plants (other than tree trunks) at this nuisance pest rate.**

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. If freezing occurs, place the product in a warm room (68°F or above) and allow to thaw. Roll or shake container frequently until thawing is complete.

Pesticide

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If pesticide, spray mixture, or rinsate cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container

Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or incinerate, or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

For minor spills, leaks, etc., follow all precautions indicated on this label

and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid contamination of food and feed. Food utensils, such as tablespoons and measuring cups, should not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Do not contaminate ornamental fish ponds.

Statement of Practical Treatment

This product is an organophosphate.

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If inhaled: Remove victim to fresh air. If not breathing, apply artificial respiration, preferably mouth-to-mouth. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Note to Physician: If ingested, induce emesis or lavage stomach. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Personal Protective Equipment

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping of sprays. If waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

D•z•n® trademark of Ciba-Geigy Corporation

©1996 Ciba-Geigy Corporation

Ciba Crop Protection
Ciba-Geigy Corporation
Greensboro, North Carolina 27419

CGA 1L145(DRAFT)

April 18, 1995

Rates/Uses corresponds to CGA 1L128G 034
Dizīn diazinon AG500

December 4, 1995

Revised pending label - cranberries - changed
gallorage to 15 ground, 5 aerial, added rate
range for cranberry fruitworm per Cranberry
Institute

February 9, 1996

Revised rates to be closer equivalent to
AG500; 2/9/96 - changed "Apply to ... in mole
cricket dfu's to broadcast ... added clarification
"Notes" to Strawberry dfu's

April 8, 1996

EPA approved regis. 3/18/96 with Comments.
3/27/96 - revised label accordingly, also delet-
ed PPE and other WPS lang. from container
label - these are allowed to be in booklet.

September 18, 1996

Revised - Added "Low Odor ..." claim

[QUARK/DZN DIAZINON AG600 WBC/DZN DIAZ AG600 WBC-A] - ccg - 9/20/96

RESTRICTED USE PESTICIDE

DUE TO AVIAN AND AQUATIC TOXICITY

FOR RETAIL SALE TO AND USE BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

(Jug Label for 2½ Gal.)

D•z•n® diazinon AG600 WBC

INSECTICIDE
Water-Based Concentrate

For control of certain insects on fruits, nuts, vegetables, field crops, lawns and ornamentals.

This product must not be used on golf courses and sod farms.

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate56.0%
Inert Ingredients:	44.0%
Total:	100.0%

EPA Reg. No. 100-784

EPA Est. 100-AL-1

See directions for use in attached booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

2 1/2 gals.
U.S. Standard Measure

KEEP OUT OF REACH OF CHILDREN.

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid contamination of food and feed. Food utensils, such as tablespoons and measuring cups, should not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets or livestock. Do not contaminate ornamental fish ponds.

Statement of Practical Treatment

This product is an organophosphate.

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If inhaled: Remove victim to fresh air. If not breathing, apply artificial respiration, preferably mouth-to-mouth. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Note to Physician: If ingested, induce emesis or lavage stomach. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates. Rates

above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping of sprays. If waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Chemigation

Refer to supplemental labeling in attached booklet for chemigation use directions for cranberries. Do not apply this product to cranberries through any type of irrigation system, unless the specific directions for chemigation are followed.

Do not apply this product to any other crop or site through any type of irrigation system.

D•z•n® trademark of Ciba-Geigy Corporation

©1996 Ciba-Geigy Corporation


Ciba Crop Protection
 Ciba-Geigy Corporation
 Greensboro, NC 27419

CGA 1L145(DRAFT)

[QUARK/DZN DIAZINON AG600 WBC/DZN DIAZ AG600WBC-A] - CCG - 9/20/96



SEP 2 1988
P/DP/01



96 SEP 24 2:42

REGU LPA: C/P/DPD1