
Fibase read instructions on reve	erse before completing forn	. 100-792	4/13/200	1 Page 17			
≎EPA	United States Environmental Protect Washington, DC 2	•	Registration Amendment Other	OPP Identifier Number NOTIFICATION			
	Applicat	ion for Pesticide - Se	ction I				
Company/Product Number 100-792		2. EPA Produ Mary Waller		. Proposed Classification			
4. Company/Product (Name) Ridomil Gold® PC Granular Fung	icide	PM# 21		x None Restricted			
Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419 Check if this is a new address		•	my product is similar or identical in composition and labeling to: EPA Reg. No.				
			44.45				
Amendment - Explain be	se to Agency letter dated		Final printed labels in res Agency letter dated "Me Too" Application.	Ponse to NOTIFICATION			
X Notification - Explain below. Other - Explain below.			APR 1 3 2001				
to enforcement action and penalties un	rovisions of PR Notice 98-10 and f nt of formula of this product. I undo n is not consistent with the terms of der sections 12 and 14 of FIFRA. iction, Inc. 2) The Conditions of Sa. Protection, Inc. and Zeneca Ag Protection, Inc. and Teneca Ag Protection T	EPA regulations at 40 CFR 152.4 erstand that it is a violation of 18 of PR Notice 98-10 and 40 CFR 1 The following changes are being le and Warranty statement has be oducts, we have chosen to use the change. 3) The copyright date re	6, and no other changes hav U.S.C. Sec. 1001 to willfully r 52.46, this product may be in made via this notification: 1) een changed to reflect the na he former Zeneca warranty st vilects Syngenta. 4) Tradema	make any faise statement to EPA. I violation of FIFRA and I may be subject Company name and address have been ime change. Because Syngenta has been atement as the Syngenta warranty rk statements have been updated to			
		Section - III					
Material This Product Will Child-Resistant Packaging Yes* No *Certification must	Be Packaged In: Unit Packaging Yes No If "Yes" No. pe	Water Soluble Pack Yes No If "Yes"	kaging 2. Type	of Container Metal Plastic Glass Paper			
be submitted	Unit Packaging wgt. Conta	iner Unit Packaging wgt.	container	Other (Specify)			
Location of Net Contents Information Label Container 4. Size(s) F		ze(s) Retail Container	il Container 5. Location of Label Directions On Label On Labeling accompanying product				
6. Manner in Which Label is Aff	fixed to Product	Lithograph Paper glued Stenciled	Other				
		Section - IV					
Contact Point (Complete ite Name	ms directly below for identifica	ation of individual to be conta Title		ocess this application.) elephone No. (Include Area Code)			
•	Certi I have made on this form and vingly faise or misleading state		rue, accurate and comple	6. Date Application Received (Stamped)			
1. Typed Name		Label Group Leader 5. Date		6 ((() 6 (())			
Nan S. Padgett		March 26, 2001		(

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsolete.

Ridomil Gold®



KEEP OUT OF REACH OF CHILDREN. WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

GRANULAR FUNGICIDE

For the control of diseases of beans, cotton, and peanuts caused by Pythium and Rhizoctonia

See additional precautionary statements and directions for use on back of bag.

Ridomil Gold PC is a granular fungicide.

EPA Reg. No. 100-792 EPA Est. 5905-GA-1 **NOTIFICATION**

APR 1 3 2001

PRODUCT ID. **52671**

50 POUNDS NET WEIGHT

syngenta



CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under configurationably foreseeched to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR CROP INJURY.

GENERAL INFORMATION

Ridomil Gold PC is a granular fungicide containing 0.5% Ridomil Gold active ingredient, a systemic fungicide, and 10% PCNB (pentachloronitrobenzene). It provides control of damping-off and seed and seedling rot diseases in beans, cotton, and peanuts caused by Pythium and Rhizoctonia. Ridomil Gold PC will also provide suppression of pod rot in peanuts caused by *Sclerotium rolfsii*.







Ridomil Gold® PC

Note: Ridomil Gold PC is a systemic fungicide having a specific mode of action and could be subject to development of insensitive strains of fungi. Development of insensitivity cannot be predicted. Therefore, Syngenta cannot assume liability for crop damage resulting from insensitive strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance or ways to control any possible Ridomil Gold insensitive strains of fungi which may occur.

Applicator Settings

It is important that the user calibrate each individual granular applicator before its use. The settings in the chart under the **Cotton** section below SHOULD ONLY BE USED AS A STARTING POINT. Even after calibration, it is best to check the actual application rate under your operating conditions. Application rates can vary due to changing moisture conditions, equipment wear, normal variation in product bulk density and particle size distribution.

Calibration: To calibrate a granular applicator, follow the procedure as outlined below:

- 1. Fill the hopper with Ridomil Gold PC.
- 2. Attach a plastic bag to the spout opening.
- Refer to the chart in the Cotton section of this label and adjust the setting on the applicator to match the desired rate per acre.
- 4. Measure off 1,000 linear ft. and drive at the speed selected.
- 5. Weigh or measure the volume of each bag, change settings as needed to obtain the desired rate and repeat steps 2-5 until the setting delivers the correct number of ounces per 1,000 ft.
- 6. Repeat steps 1-5 for each applicator box.

Conversions:

1. Time to cover 1,000 ft. at:

2 mph = 5 min. 40 sec. 3 mph = 3 min. 54 sec. 4 mph = 2 min. 50 sec. 5 mph = 2 min. 16 sec. 6 mph = 1 min. 57 sec.

2. Ounces (wt.) of granules to approximate ounces (volume):

```
6 oz. wt. = 8 fl. oz. (1 level cup)
```

When calibrating granular applicators, follow the recommendations below:

- To reduce granule flow variability, adjust orifice openings by moving the controls in one direction only, such as from the closed position to open.
- Granule flow through an orifice is not always proportional to ground speed, so make calibration checks at the same speed at which you plant and maintain that ground speed as constantly as possible when planting.
- Granular pesticide products vary considerably in size, density, and particle shape, so recalibrate applicators whenever you change brands and/or formulations. The same brand can even change from year to year.
- 4. Changing humidity and temperature during the day can change the flow rate of granules through the applicator's orifice. Recalibrate when conditions change and periodically readjust orifice settings to compensate.
- 5. Know the recommended rate. For band applications, the area actually is the area covered by the band, not total cropland area planted. Some row-crop recommendations are based on treating in-the-row and these rates generally are specified as amounts (ounces) of granules per certain row length (often 1,000 ft.).
- Calibrate frequently. There are several acceptable calibration methods. Learn one. Do the arithmetic first, then complete the material collection phase under field conditions.

Calculate the amount of Ridomil Gold PC needed for band treatment as follows:

Band Width in Inches

Row Spacing in Inches

X Broadcast Rate per Acre

Band Rate per Acre

BEANS

Ridomil Gold PC will control damping-off and seed and seedling rots of dry and green beans (bush and pole types) caused by Pythium and Rhizoctonia when used at planting.

Apply 3/4 lb. (12 oz.) of Ridomil Gold PC per 1,000 linear ft. of row at the time of planting. Adjust the application equipment so the granules are mixed with the soil surrounding the seed.

Note: To avoid possible illegal residues, do not allow the feeding of vines or grazing of foliage by livestock.

Ridomil Gold® PC

5/7

COTTON

Ridomil Gold PC will control damping-off, seed and seedling rot diseases caused by *Pythium* spp. and *Rhizoctonia solani*.

For best results, use high quality seed and plant into a properly prepared seedbed.

Apply 7-10 lbs. of Ridomil Gold PC per 13,000 linear ft. of row (8.6-12.3 oz./1,000 linear ft.) at the time of planting. Adjust the application equipment so the granules are mixed with the soil surrounding the seed. See Chart 1 below for suggested applicator settings and the **General Information** section of this label for calibration instructions.

Note: To avoid possible illegal residues, do not allow the feeding or grazing of cotton foliage by livestock.

Chart 1: Cotton - Suggested Application Settings

		40" Rows In-Furrow Lbs./A				
]	10	9	8	7	
		Equivalent Co./f_con ft.				
Applicator Brand	Speed MPH	12.3	11,1	9.8	8.6	
Gandy	4	28	26.8	25	23.5	
	5	31	30	28	26.3	
	6	34.5	32.5	30.5	29	
Allis Chalmers International Harvester 1980 and Before	4	2.4	2.1	1.9	1.6	
	5	3	2.8	2.4	2.05	
	6	3.6	3.2	2.9	2.5	
Noble 1972	4	19	17	15	13	
and Before	5	23	21	19	17	
	6	27	25	22	20	
Noble (Remcor) 1973-Present	4	14	13	12 or 11	10	
White 5100 Series	5	17 or 16	15	14	13 or 12	
Deutz Allis 1985-Present	6	19	18	16	15 or 14	
John Deere	4	2/12	2/10	2/8	2/5	
Metal Box	5	2/18	2/15	2/12	2/9	
	6	2/22	2/19	2/16	2/13	
John Deere 7000 and 7100	4	28	27	24	22	
Max-Emerge®	5	31	30	28	26	
Meter Scale 01020	6	33	32	31	29	
John Deere 7000 and 7100	4	18	17	16	15	
Max-Emerge	5	21	20	18	17	
Meter Scale 0515	6	23	22	21	19	
International Harvester	4	3.1	2.9	2.6	2.35	
1981-1985 Case 1H	5	3.6	3.4	3.1	2.8	
1985-Present	6	4.5	4.0	3.55	3.2	
Kinze 1980-Present	4	28	27	24	22	
Meter Scale 010 20	5	31	30	28	26	
	6	33	32	31	29	

PEANUTS

Ridomil Gold PC applied at planting will provide control of seed and seedling rots of peanuts caused by Pythium and Rhizoctonia. Applications of Ridomil Gold PC during early pegging will provide control of early pod rot caused by Pythium and Rhizoctonia and will provide suppression of *Sclerotium rolfsii* during early pegging. These treatments will not provide late season control of these diseases.

Seed and Seedling Rots: Apply 12.5-25 lbs. of Ridomil Gold PC per 14,520 linear ft. (36-inch row) in a 4-inch band at the time of planting and mix in the covering soil. Incorporate with mechanical equipment to a depth of 2-4 inches. For double-drilled or double-row plantings, apply 12.5-25 lbs./14,520 linear ft. in a 4-inch band centered on each drill or row. Use the higher rate in fields with a history of seedling disease problems or when weather conditions are favorable for infection and disease development.

Pod Rot: Apply 50 lbs./14,520 linear ft. (36-inch row) at early pegging. For banded applications, adjust the equipment so the granules are distributed uniformly over the row(s) and in the pegging zone. For mid- to late-season control of pod rot, use other EPA-registered fungicides.

Sclerotium rolfsii: When Ridomil Gold PC is used for pod rot control, some suppression of *Sclerotium rolfsii* will be obtained. In areas where *Sclerotium rolfsii* is a problem use other fungicides that are labeled for this disease. Use other professional fall shed for mid- and later seas an control of southern blight/white mold infections.

Notes: To avoid possible illegal residues, (1) Do not allow the feeding or grazing of peanut foliage or harvest residues by livestock. (2) Do not apply more than 75 lbs. Ridomil Gold PC per 14,520 linear ft. of row per season. (3) Do not apply within 75 days of harvest.

ROTATIONAL CROPS

These crops represent those which may be planted (rotated to) according to the interval listed following an application of Ridomil Gold PC to beans, cotton, or peanuts. Do not plant any crop not registered for use with Ridomil Gold active ingredient in soil treated with this active ingredient for a period of 12 months, unless a shorter interval is specified on the following list. Do not plant root crops in fields treated with this product within 12 months of application.

Crop	Rotational Interval	
Alfalfa, Almonds, Apples, Asparagus, Avocados		
Blueberries, Broccoli	0 days	
Cabbage, Cauliflower, Chinese broccoli (gai lon, white flowering broccoli), Chinese cabbage (tight-heading varieties only). Citrus, Cotton, Cucurbit Vegetables		
Deciduous Fruits and Nuts*		
Eggplant		
Grapes		
Hops		
Leafy Vegetables (excluding brassica), Legume Vegetables (beans and peas-succulent and dried)		
Papaya, Peanuts, Peppers, Pineapples		
Raspberries		
Soybeans, Spinach, Stone Fruits, Strawberries		
Tobacco, Tomatoes		
Walnuts		
Brussels Sprouts		
Corn	12 months	
Garlic		
Root Crops		
Wheat		
Crops Not Intended for Food or Feed	0 days	
All Other Crops Intended for Food or Feed	12 months	

^{*}These crops and other perennial crops may be planted immediately following last application of Ridomil Gold PC provided they will not bear harvestable fruit within 12 months.

STORAGE AND DISPOSAL

Pesticide

Store in a dry place. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide is a violation of federal law. Pesticide that cannot be used according to label instructions must be disposed of according to federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Disposal

Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke

For minor spills, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

Causes substantial, but temporary eye injury. Do not get in eyes or on clothing. Harmful if absorbed through skin. Avoid contact with skin.

Statement of Practical Treatment

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

If swallowed: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol.

Note to Physician: If ingested, induce emesis or lavage stomach. Treat symptomatically.

Personal Protective Equipment

Applicators and other handlers must wear:

- . Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- · Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- · Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Cover or incorporate all granules. Do not contaminate water when disposing of equipment wash waters.

Ridomil Gold® trademark of Syngenta

Max-Emerge® trademark of John Deere Company

©2001 Syngenta

Syngenta Crop Protection, Inc. Greensboro, North Carolina 27409 www.syngenta-us.com SCP 792A-M2B 0201