

PM 21

100-797

6/24/96

P2 1/17

Gregory T. Peters, Ph.D.
Ciba Crop Protection
Ciba-Geigy Corporation
P. O. Box 18300
Greensboro, NC 27419-8300

JUN 24 1996

Dear Dr. Peters:

Subject: Mefenoxam 45W (Reg. No. 100-797)
Mefenoxam LS (Reg. No. 100-799)
Label Amendments
Your Correspondence of May 3, 1996

The labeling referred to above, is acceptable provided that you:

1. Submit and/or cite all data required for registration or reregistration of your product under FIFRA sec. 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Stamped copies of the labels are enclosed for your records. Should you have any questions or concerns, please call Kathryn Scanlon on (703) 305-7382.

Sincerely Yours,

Connie B. Welch
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (7505C)

7505C:K.Scanlon:ks:05/17/96:MEFEAPRON.LTR

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	K. Scanlon							
DATE ▶	Jun 19, 1996							

DRAFT

Highlighted

2/17

Booklet Label - 1 lb.
Bag Label - 4 lb.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

JUN 24 1996

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

100-797

*Revised
"Mecorax 450"
(Here and
elsewhere.)*

Apron XL™ WS

Accu-Pak®

FUNGICIDE

A seed treatment chemical for control of *Pythium* and *Phytophthora* causing damping-off, seed rot, and systemic downy mildew diseases of certain crops.

One Pound
Net Weight

Four Pounds
Net Weight

Active Ingredient:

(R)-2-[(2,6-dimethylphenyl)-
methoxyacetylamino]-propionic acid
methyl ester 43.6%

Related Compounds 1.4%

Inert Ingredients: 55.0%

Total: 100.0%

*Deleted
Mecorax*

Apron XL WS is a wettable powder packaged in a water-soluble bag.

EPA Reg. No. 100-797 *Added*

EPA Est. 67545-AZ-1

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements on panel at left and directions for use on panel at right. (For the 4 lb. bag)

See additional precautionary statements and directions for use inside booklet. (For the 1 lb.)

This outer protective container contains Apron XL WS in an inner water-soluble bag. Entire inner bag and contents dissolve in water. After opening outer container, immediately drop entire unopened inner bag into the partially filled sprayer or mix tank. Do not handle the soluble bag or expose it to moisture, since this may cause rupturing.

[QUARK/APRON XL/APRON XL WS] - ccg - 5/1/96

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire **Directions for Use** and the **Conditions of Sale and Warranty** before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The **Directions for Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as *weather conditions, presence of other materials, or the manner of use or application* all of which are beyond the control of Ciba-Geigy or the Seller. All such risks shall be assumed by the Buyer.

Ciba-Geigy warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use** subject to the inherent risks referred to above. **Ciba-Geigy makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Ciba-Geigy or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.** Ciba-Geigy and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing **Conditions of Sale and Warranty**, which may be varied only by agreement in writing signed by a duly authorized representative of Ciba-Geigy.

General Information

Apron XL™ WS is a systemic fungicide seed dressing specifically for control of systemic downy mildews and diseases caused by soil-borne *Pythium* and *Phytophthora* spp. When a rate range is specified, use higher rates of Apron XL WS when the disease pressure is expected to be high.

For control of other soil-borne diseases, such as *Rhizoctonia* spp., Apron XL WS should be applied in combination with other registered seed dressing fungicides.

Note: Apron XL 45W is a systemic fungicide having a specific mode of action. Apron XL 45W could be subject to development of insensitive strains of fungi. Development of insensitivity cannot be predicted. Therefore, Ciba-Geigy cannot assume liability for crop damage resulting from insensitive strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Apron XL 45W insensitive strains of fungi which may occur.

Apron XL WS may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry- or mist-type commercial seed treatment equipment.

Mixing Instructions

Prepare no more mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. **Vigorous agitation is necessary to dissolve the bag and disperse the product. Maintain maximum agitation throughout the spraying operation.** Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use.

Note: 1 lb. of Apron XL WS displaces 0.7 lbs. of water.

Precautions: (1) Water soluble bags of Apron XL WS must be completely dissolved and dispersed in plain water before any other tank mix partner is added to the spray solution. (2) The maximum recommended concentration of Apron XL WS in slurries is 7 oz. of Apron XL WS in 10 ozs. of water per 100 lbs. of seed.

Apron XL WS Alone: Add 1/4 of the required amount of plain water to the spray or mixing tank. With the agitator running, drop the required number of **unopened** soluble bags of Apron XL WS into the tank all at once. Continue agitation while adding the remainder of the water. Begin

Added

application of the spray solution after the bags have dissolved and the material has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Apron XL WS + Tank Mixtures: Add 1/4 of the required amount of plain water to the spray or mixing tank. With the agitator running, drop the required number of **unopened** soluble bags of Apron XL WS, and any other products packaged in water soluble bags, into the tank all at once. After the water soluble bags have dissolved and the products have dispersed uniformly into the mix water, continue to fill the tank with water. Then add the desired amount of other products recommended for tank mixture and allow them to become completely dispersed. Continue agitation to maintain a uniform suspension until all of the spray solution has been applied.

If using Apron XL WS in a tank mixture with other seed treatment chemicals, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate should be exceeded, and the most restrictive label precautions and limitations should be followed. This product should not be mixed with any product which prohibits such mixing.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

Seed Treatment - Domestic Use

Carrots

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

Reduced 50%

Cereal Grains

- Wheat, barley, rye, oats, millet, rice, corn (field corn, sweet corn, and popcorn), buckwheat, and triticale.

For Pythium damping-off control: Apply Apron XL WS as a seed treatment at the rate of 0.27-0.53 ozs. per 100 lbs. of seed.

For Pythium seed rot and damping-off control in field corn when applied in combination with Maxim™ or captan products at labeled rates: *added* Apply Apron XL WS as a seed treatment at 0.035 ozs. per 100 lbs. of seed. *Reduced 54%*

For systemic downy mildew control of sweet corn: Apply Apron XL WS as a seed treatment at the rate of 1.06 ozs. per 100 lbs. of seed. *Reduced 50%*

Helded For Pythium damping-off control in wheat, barley, rye, and oats, when applied in combination with Dividend® or other seed treatment products labeled for disease control in these crops: Apply Apron XL WS as a seed treatment at 0.035 ozs. per 100 lbs. of seed.

For Pythium seed rot and damping-off control in rice when applied in combination with Vitavax® 200, 42-S Thiram, or RTU® Vitavax®-Thiram at labeled rates: Apply Apron XL WS as a seed treatment at 0.035-0.07 ozs. per 100 lbs. of seed. *Reduced 10-50%*

Note: If using Apron XL WS treated rice seed, do not use Ridomil Gold™ for control of Pythium in water seeded rice. *Replaced "Mistakenly"*

Cotton

- For Pythium seed rot and damping-off control:

Crops Not Listed

Apply Apron XL WS as a seed treatment to machine-delinted or acid-delinted cottonseed at the rate of 0.27-0.53 ozs. per 100 lbs. of seed.

Forage Grasses

- Grasses grown for hay grazing or silage, corn (fodder or silage), and small grain grown for hay grazing or silage.

Reduced 50%

For Pythium damping-off control: Apply Apron XL WS as a seed treatment at the rate of 0.27-0.53 ozs. per 100 lbs. of seed.

Forage Legumes

- Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch, and velvet beans (for forage).

For Pythium damping-off and early season Phytophthora control: Apply Apron XL WS as a seed treatment at the rate of 0.53 ozs. per 100 lbs. of seed.

For suppression of early season downy mildew in alfalfa and soybeans: Apply Apron XL WS as a seed treatment at the rate of 0.53 ozs. per 100 lbs. of seed.

Garden/Table Beets

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at the rate of 0.53 ozs. per 100 lbs. of seed.

Legume Vegetables and Okra

- Black-eyed peas, cowpeas, edible soybeans, soybeans, field beans, chick peas (garbanzo beans), field peas, garden peas, green beans, kidney beans, lima beans, navy beans, okra, pinto beans, peas, pole beans, snap beans, string beans, wax beans, lentils, and lupines.

Added

For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

For Pythium damping-off control in field beans, kidney beans, navy beans, pinto beans, and lentils when disease pressure is expected to be low: Apply Apron XL WS as a seed treat-

Added

ment at 0.07-0.27 ozs. per 100 lbs. of seed in combination with other seed treatment products labeled for these crops.

For early season Phytophthora control: Apply Apron XL WS as a seed treatment at 0.53 ozs. per 100 lbs. of seed.

Reduced 55%

For Pythium damping-off and systemic downy mildew control of peas: Apply Apron XL WS as a seed treatment at the rate of 1.06 ozs. per 100 lbs. of seed.

Reduced 62%

Peanuts

- For Pythium seed rot and damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

Reduced 56%

Sorghum

- For Pythium damping-off control on all sorghum cultivars: Use 0.27-0.53 ozs. of Apron XL WS per 100 lbs. of seed applied as a seed treatment.

Reduced 52-75%

For Pythium damping-off control when pressure is expected to be low: Apply Apron XL WS as a seed treatment at 0.07-0.27 ozs. per 100 lbs. of seed in combination with Maxim or captan products at labeled rates.

Handwritten note: 1-23-88

For systemic downy mildew control: Apply 0.53 ozs. of Apron XL WS per 100 lbs. of seed on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew grown in areas with recent history of severe disease, apply 1.06 ozs. of Apron XL WS per 100 lbs. of seed as a seed treatment.

Reduced 50%

Note: Seed treated with the 1.06 ozs. rate should be planted and not carried over to the following year.

Only high quality sorghum seed (90% germ and above) should be treated with Apron XL WS.

Soybeans

- For Pythium damping-off and early season Phytophthora control: Apply Apron XL WS as a seed treatment at the rate of 0.27-0.53 ozs. per 100 lbs. of seed.

Handwritten note: 1-23-88

For Pythium damping-off control when disease pressure is expected to be low: Apply Apron XL WS as a seed treatment at 0.07-0.27 ozs. per 100 lbs. of seed in combination with other seed treatment products labeled for soybeans.

Added

Spinach

Added

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

Note: Do not apply Apron XL WS to spinach which is destined to be grown in the greenhouse. Containers or bags containing Apron-treated spinach seed must be labeled with the restriction. "Do not use seed for spinach destined to be grown in the greenhouse."

Sugar Beets

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at the rate of 0.27-0.53 ozs. per 100 lbs. of seed.

Reduced 50-75%

Sunflower

- For systemic downy mildew control: Apply Apron XL WS as a seed treatment at the rate of 1.06 ozs. per 100 lbs. of seed.

← Reduced

Turfgrasses (All)

- For Pythium damping-off control on golf courses, home lawns, reclamation, and erosion control projects: Apply Apron XL WS as a seed treatment at the rate of 0.53-1.06 ozs. per 100 lbs. of seed.

50%

Note: For long-term control of Pythium in areas when using Apron XL-treated seed, make an application of Subdue® MAXX® 7-10 days after seeding.

↖ Replaced Metconaxa

Seed Treatment-Export Use

FOR TREATMENT OF THE FOLLOWING SEED FOR EXPORT ONLY TO COUNTRIES REQUIRING TREATMENT WITH Apron XL WS BEFORE IMPORTATION. User should verify actual registered rate in country of destination and/or use.

Brassica (Cole) Leafy Vegetables

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

Reduced

Corn

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs.

50%

per 100 lbs. of seed. For systemic downy mildew control: Apply Apron XL WS as a seed treatment at 1.77 ozs. per 100 lbs. of seed.

Cucurbits

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

Reduced 50%

Fruiting Vegetables (e.g. tomatoes, peppers eggplant)

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

Leafy Vegetables

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

Millet

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed. For systemic downy mildew control: Apply Apron XL WS as a seed treatment at 2.0 ozs. per 100 lbs. of seed.

Reduced 57%

Onions

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

Reduced 50%

Peas

- For Pythium damping-off and early season Phytophthora control: Apply Apron XL WS as a seed treatment at the rate of 0.53-1.06 ozs. per 100 lbs. of seed. For downy mildew control, apply Apron XL WS as a seed treatment at 1.06 ozs. per 100 lbs. of seed.

Reduced 57%

Root and Tuber Vegetables

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed.

Note: Do not apply Apron XL WS to potatoes.

Reduced 50%

Sorghum

- For Pythium damping-off control: Apply Apron XL WS as a seed treatment at 0.27-0.53 ozs. per 100 lbs. of seed. For systemic downy mildew control: Apply Apron XL WS as a seed treatment at 1.77 ozs. per 100 lbs. of seed.

Sunflower

- For systemic downy mildew control: Apply Apron XL WS as a seed treatment at 3.1-3.72 ozs. per 100 lbs. of seed.

Reduced
50%

Seed treated in accordance with the export use pattern must be labeled as follows: FOR EXPORT ONLY - NOT FOR DOMESTIC SALE OR USE.

Note: Federal law requires that bags containing treated seeds shall be labeled with the following information:

"This seed has been treated with Apron XL WS fungicide. Do not use for feed, food, or oil purposes. Store away from feeds and foodstuffs." Use with an EPA-approved dye that imparts an unnatural color to the seed.

Storage and Disposal

Storage - Store in a cool, dry place. Do not store this product under wet conditions. Handle outer packaging carefully to avoid breakage of inner soluble bag.

Pesticide Disposal

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Disposal

Do not reuse outer container. Dispose of empty outer container in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if swallowed or absorbed through skin. Avoid contact with skin.

Statement of Practical Treatment

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

Note to Physician: If ingested, induce emesis or lavage stomach. Treat symptomatically.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

<p>User Safety Recommendations</p> <p>Users should:</p> <ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Apply only as specified on this label. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

Accu-Pak®, Dividend®, Maxim™, Ridomil®, Ridomil Gold™ and Subdue® MAXX® trademarks of Ciba-Geigy Corporation

Advised

RTU® trademark of Gustafson, Inc.

Vitavax® trademark of Uniroyal Chemical Company, Inc.

©1996 Ciba-Geigy Corporation

Ciba Crop Protection
Seed Treatment Products
Ciba-Geigy Corporation
Greensboro, North Carolina 27419

CGA 175L1 046 (Booklet)
CGA 175L2 046 (Bag)

[QUARKAPRON XL/APRON XL WS] - ccg - 5/1/96

15/17

Base Label (for Booklet)

Entire Label
100% of Label
Text from Booklet

Apron XL WS

Accu-Pak®

FUNGICIDE

A seed treatment chemical for control of *Pythium* and *Phytophthora* causing damping-off, seed rot, and systemic downy mildew diseases of certain crops.

One Pound
Net Weight

Active Ingredient:	
(R)-2-[(2,6-dimethylphenyl)-	
methoxyacetylamino]-propionic acid	
methyl ester	43.6%
Related Compounds	1.4%
<hr/>	
Inert Ingredients:	55.0%
Total:	100.0%

Apron XL WS is a wettable powder packaged in a water-soluble bag.

EPA Reg. No. 100-797

EPA Est. 67545-AZ-1

See additional precautionary statements and directions for use inside booklet. (For the 1 lb.)

<p>AGRICULTURAL USE REQUIREMENTS</p> <p>Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.</p>
--

KEEP OUT OF REACH OF CHILDREN.

CAUTION

Container Disposal

Do not reuse outer container. Dispose of empty outer container in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if swallowed or absorbed through skin. Avoid contact with skin.

Statement of Practical Treatment

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

Note to Physician: If ingested, induce emesis or lavage stomach. Treat symptomatically.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Apply only as specified on this label. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

Accu-Pak®, Dividend®, Maxim™, Ridomil®, Ridomil Gold™, and
Subdue® MAXX® trademarks of Ciba-Geigy Corporation

Accu-Pak® trademark of Ciba-Geigy Corporation

RTU® trademark of Gustafson, Inc.

Vitavax® trademark of Uniroyal Chemical Company, Inc.

©1996 Ciba-Geigy Corporation

Ciba Crop Protection
Seed Treatment Products
Ciba-Geigy Corporation
Greensboro, North Carolina 27419

CGA 175L1 046 (Booklet)
CGA 175L2 046 (Bag)

[QUARK/APRON XL/APRON XL WS] - ccg - 5/1/96