

100-798

read instructions on reverse before completing form.

12/22/1997



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number  
**NOTIFICATION**

**Application for Pesticide - Section I**

1. Company/Product Number 100-798	2. EPA Product Manager Mary Waller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Ridomil Gold® GR Fungicide <small>(NCP 166L1A 0697)</small>	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Novartis Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419  <input type="checkbox"/> Check if this is a new address	6. <b>Expedited Review.</b> In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____ Product Name _____	

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**NOTIFICATION**  
DEC 22 1997

**Explanation:** Use additional page(s) if necessary. (For Section I and Section II.)

**PR Notice 95-2**  
Company name has been changed from Ciba to Novartis.

**Section - III**


1. <b>Material This Product Will Be Packaged In:</b>			2. <b>Type of Container</b>	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
*Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Unit Packaging wgt. No. per container	<input type="checkbox"/> Glass
				<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____
3. Location of Net Contents information <input type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product		
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____		

**Section - IV**

1. <b>Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)</b>		
Name Lee Hubbard	Title Senior Regulatory Manager	Telephone No. (Include Area Code) 910- 632-7034
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Senior Regulatory Manager	
4. Typed Name Lee Hubbard	5. Date December 10, 1997	

NOTIFICATION  
DEC 22 1997

# Ridomil Gold®



## FUNGICIDE

For control of  
Phytophthora and  
Pythium diseases of  
avocados, citrus,  
cotton, cranberries,  
ginseng, leafy  
vegetables,  
nonbearing deciduous  
fruits and nuts,  
peanuts, raspberries,  
soybeans, spinach,  
sugar beets, and  
tomatoes

<b>Active Ingredients:</b>	
(R)-2-[(2,6-dimethylphenyl)-methoxyacetyl- amino]-propionic acid methyl ester .....	2.43%
<b>Related Compounds</b> .....	0.07%
<b>Inert Ingredients:</b>	97.50%
<b>Total:</b>	100.00%

KEEP OUT OF REACH OF CHILDREN.  
**CAUTION**  
See additional precautionary statements  
on left side of bag and directions for use on  
right side of bag.

Ridomil Gold GR is a granular fungicide.

EPA Reg. No. 100-798  
EPA Est. 5905-GA-01

PRODUCT ID.  
**52581**

**50 POUNDS**  
NET WEIGHT

 **NOVARTIS**

# Ridomil Gold® GR

## DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

**IMPORTANT:** Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

### CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Novartis Crop Protection, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Novartis warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. **Novartis makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Novartis or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.** Novartis and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Novartis.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.** Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.


PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR CROP INJURY.**

### GENERAL INFORMATION

Ridomil Gold GR is a systemic fungicide which provides control of certain diseases caused by members of the Oomycete class of fungi on selected crops.



## Ridomil Gold® GR

4/11

**Note:** Ridomil Gold GR is a systemic fungicide having a specific mode of action and could be subject to development of insensitive strains of fungi. Development of insensitivity cannot be predicted. Therefore, Novartis cannot assume liability for crop damage resulting from insensitive strains of fungi. If treatment is not effective following the use of Ridomil Gold GR as recommended, an insensitive resistant strain of fungi may be present. If the treatment is ineffective due to the presence of a Ridomil Gold GR insensitive strain of fungi, neither Ridomil Gold GR nor any other fungicide with similar action will effectively control that disease. Consideration should then be given to the prompt use of other types of suitable fungicides. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance in your particular crop and disease control situation.

Do not use in nurseries, turf, or landscape plantings, unless specified in this label.

**THIS LABEL IS FOR FIELD USE ONLY AND IS NOT INTENDED FOR USE ON TRANSPLANT TRAYS, GREENHOUSES, LATH HOUSES, FLOAT HOUSES, HYDROPONIC PRODUCTION, OR IN BEDDING PLANT STRUCTURES.**

Calculate the amount of Ridomil Gold GR needed for band treatment by the formula:

$$\frac{\text{band width in inches}}{\text{row spacing in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

### Applicator Settings

It is important the user calibrate each individual granular applicator before its use. The settings in the chart in the **Soybeans** section of the label SHOULD ONLY BE USED AS A STARTING POINT. Even after calibration, it is best to check the actual application rate under your operating conditions. Application rates can vary due to changing moisture conditions, equipment wear, normal variation in product with density and particle size distribution.

**Calibration:** To calibrate a granular applicator, follow the procedure as outlined below:

1. Fill the hopper with Ridomil Gold GR.
2. Attach a plastic bag to the spout opening.
3. Refer to the appropriate chart in the **Soybeans** section of this label and adjust the setting on the applicator to match the desired rate per acre.
4. Measure off 1,000 linear ft. and drive at the speed selected.
5. Weigh or measure the volume of each bag, change settings as needed to obtain the desired rate and repeat steps 2-5 until the setting delivers the correct number of ounces per 1,000 ft.
6. Repeat steps 1-5 for each applicator box.

### Conversions:

1. Time to cover 1,000 ft. at:
  - 4 mph = 2 min. 50 sec.
  - 5 mph = 2 min. 16 sec.
  - 6 mph = 1 min. 57 sec.
2. Ounces (wt.) of granules to approximate ounces (volume):
  - 6 oz. wt. = 8 fl. oz. (1 level cup)

When calibrating granular applicators, follow the recommendations below:

1. To reduce granule flow variability, adjust orifice openings by moving the controls in one direction only, such as from the closed position to open.
2. Granule flow through an orifice is not always proportional to ground speed, so make calibration checks at the same speed you plant at and maintain that ground speed as constant as possible when planting.
3. Granular pesticide products vary considerably in size, density and particle shape, so recalibrate applicators whenever you change brands and/or formulations. The same brand can even change from year to year.
4. Changing humidity and temperature during the day can change the flow rate of granules through the applicator's orifice. Recalibrate when conditions change and periodically readjust orifice settings to compensate.
5. Know the recommended rate. For banded applications, the area actually is the area covered by the band, not total cropland area planted. Some row-crop recommendations are based on treating in-the-row and these rates generally are specified as amounts (ounces) of granules per certain row length (often 1,000 ft.). Others express rates as amount per treated acre which means the total area treated with the pesticide. If rates are expressed as amount per treated acre and banded applications are used, the amount of pesticide used per acre will be proportionately less. The formula above can be used to calculate the amount of Ridomil Gold GR needed per acre of crop when banded applications are made.
6. Calibrate frequently. There are several acceptable calibration methods. Learn one. Do the arithmetic first, then complete the material collection phase under field conditions.

### Preharvest Intervals

Unless otherwise stated in the directions for use for a specific crop, there is no preharvest interval for Ridomil Gold GR.

5/11

## Ridomil Gold® GR

### AVOCADOS

Ridomil Gold GR is a systemic fungicide which provides control of avocado root rot caused by *Phytophthora cinnamomi*. For best results, Ridomil Gold GR should be used before or as soon as early symptoms of disease appear. For new plantings, the use of *Phytophthora* resistant root stocks with Ridomil Gold GR is recommended. Mature trees in advanced stages of decline cannot be cured with Ridomil Gold GR. Applications are not needed during the winter months of November through February.

**Sprinkler Irrigated Orchards:** Apply  $\frac{3}{4}$ -1  $\frac{1}{2}$  oz./sq. yd. evenly to the soil surface under the canopy of the tree. The following table shows the amount of Ridomil Gold GR to use based on the diameter of the tree canopy. Start applications at the beginning of the growing season or at transplanting and make 2 additional applications at 3-month intervals.

**Drip Irrigated Orchards:** Place the recommended amount of Ridomil Gold GR (see following table) on the soil surface directly around the drip emitter at each tree. If there is more than one emitter per tree, distribute the total amount of Ridomil Gold GR among the emitters. Start applications at the beginning of the growing season or at transplanting and make 2 additional applications at 3-month intervals.

Diameter of Tree Canopy in Ft.	Amount of Ridomil Gold GR Per Tree
5	1 $\frac{1}{2}$ -3 oz.
10	6-12 oz.
15 or wider	14-28 oz.

**Notes:** (1) Do not make more than 3 applications per year and do not make the last application within 28 days of harvest, or possible illegal residues may result. (2) Do not apply more than 240 lbs. of Ridomil Gold GR per treated acre per year.

### CITRUS

Includes grapefruit, lemons, limes, oranges, tangelos, tangerines, citrus citron, kumquats, satsuma mandarin, and hybrids of these.

Use Ridomil Gold GR on citrus to control citrus foot rot, root rot, and trunk canker caused by *Phytophthora* spp. Apply to the soil as a broadcast application under the canopy of the trees. If rain is not expected within 3 days of application, water in with  $\frac{1}{2}$ -1 inch of irrigation. For banded applications, use the formula in the **General Information** section for calculating the amount of Ridomil Gold GR needed per acre of trees. If the trees are on a drip irrigation system, distribute the amount of Ridomil Gold GR needed per tree (see table) to the soil surface in the immediate vicinity of the drip emitter(s) at each tree. If there is more than one emitter at each tree, distribute the total amount of Ridomil Gold GR needed among the emitters.

**Notes:** (1) Where nematodes are a problem, best results can be achieved if effective EPA-registered nematicides are used. Nematicides can be used in combination or in sequence with Ridomil Gold GR applications. (2) For all citrus, do not apply more than a total of 240 lbs. of Ridomil Gold GR per treated acre per year.

**Precaution:** For best *Phytophthora* control, a combination of cultural practices and resistant varieties is recommended. The use of Ridomil Gold GR is not recommended in FL for use on the highly susceptible sweet orange rootstock.

#### Citrus in Nurseries (AZ, CA, FL, and PR Only)

Make the first application of Ridomil Gold GR at the time of planting and make 2 additional applications at 3-month intervals during the period when trees are actively growing. Apply 40-80 lbs. per treated acre as a broadcast or banded application to the seedbeds, liners, or bedded stock. For banded applications, use a band wide enough to cover the root systems of the plants.

**Note:** Do not use Ridomil Gold GR in greenhouse nurseries.

#### Citrus Resets or New Plantings (AZ, CA, FL, and PR Only)

Make the first application of Ridomil Gold GR to citrus resets or new plantings at the time of transplanting. Make up to 3 repeat applications at 3-month intervals to coincide with root growth flushes during the growing season. Apply as a broadcast or banded application. For banded applications, use a band wide enough to cover the root systems of the plants.

**Water Ring Treatment:** Uniformly spread 1-1  $\frac{1}{2}$  oz. within the watering ring, then drench with 5 gals. water per tree.

**Soil Surface Treatment:** Apply 40-80 lbs. treated acre (15-30 oz./1,000 sq. ft.) under the canopy of the trees. If natural rainfall is not expected within 3 days, lightly irrigate with  $\frac{1}{2}$ -1 inch of water.

#### Established Plantings

**Soil Surface Applications (FL and PR Only):** Apply 40-80 lbs. per treated acre (15-30 oz./1,000 sq. ft.) as a soil surface treatment under the canopy of the trees. Two additional applications may be made per year at 3-month intervals. Use the following table as a guide for treating individual trees.

6/11

## Ridomil Gold® GR

Diameter of Tree Canopy in Ft.	Amount of Ridomil Gold GR Per Tree
5	1/4-1/2 oz.
10	1 1/4-2 1/2 oz.
15	2 1/2-5 oz.
20	5-10 oz.

**Soil Surface Applications (AZ and CA Only):** Make one application in March-April. One or two additional applications can be made at 3-month intervals to coincide with root flushes. Use the following table to determine the proper rate based on tree size and the number of applications per year. For applications based on broadcast rates, use Ridomil Gold GR at 80 lbs. per treated acre (30 oz./1,000 sq. ft.) when 3 applications are planned and at 120 lbs. per treated acre (45 oz./1,000 sq. ft.) when 2 applications are planned. Immediately after application, irrigate with 1/2-1 inch sprinkler irrigation.

Diameter of Tree Canopy in Ft.	Ounces of Ridomil Gold GR Per Tree	
	2 Applications Per Year	3 Applications Per Year
5	3/4 oz.	1/2 oz.
10	3 1/4 oz.	2 1/2 oz.
15	7 1/2 oz.	5 1/2 oz.
20	12 oz.	10 oz.

### COTTON

Ridomil Gold GR provides control of seed rot and seedling diseases of cotton caused by *Pythium* spp. It does not control seedling diseases caused by other fungi.

#### Seed Rots and Seedling Diseases of Cotton Caused by *Pythium* spp.

Apply 1 1/4-2 1/2 lbs./13,000 linear ft. of row (1 1/2-3 oz./1,000 linear ft.) as an in-furrow application at the time of planting. Mount the application tubes so the granules are mixed with the soil covering the seeds.

For broader spectrum disease control, Ridomil Gold GR can be applied with 10-20 lbs. Terraclor® 10G per 13,000 linear ft. of row as an in-furrow application.

**Note:** Alternatively, Ridomil Gold PC may be used if the predominant pathogens are *Pythium* spp. and *Rhizoctonia* spp.

### CRANBERRIES

Use Ridomil Gold GR as a soil application for control of Phytophthora root rot of cranberries caused by *Phytophthora* spp.

Apply Ridomil Gold GR by ground or air at 20-35 lbs./A as a broadcast soil application for control of Phytophthora root rot of cranberries. Three applications per crop season are recommended. Make the first application in the fall after harvest. Make the second application in the spring, followed by a third application 45 days before harvest.

**Notes:** To avoid possible illegal residues, (1) Do not apply within 45 days before harvest, and (2) Do not apply more than 105 lbs./A per growing season.

### DECIDUOUS FRUITS AND NUTS IN ORCHARDS (NONBEARING)

#### Phytophthora Diseases

Use of Ridomil Gold GR will aid in the control of crown, collar, and root rot of deciduous fruit and nut trees caused by *Phytophthora* spp. when used in conjunction with good cultural practices and rootstocks that are most tolerant to the disease. Ridomil Gold GR applications should be made before symptoms appear, especially in areas favorable for disease development. Ridomil Gold GR will not revitalize trees showing moderate to severe disease symptoms.

Make the first application at the time of planting. (See Notes below.) Up to 2 additional applications should be made at 2 to 3-month intervals during the time when conditions are favorable for disease development. For established plantings, make the first application in the spring before growth starts.

Apply 80 lbs. per treated acre (30 oz./1,000 sq. ft.) uniformly under the canopy of the trees. The treated area is based on the area under the tree canopy or the area of the row to be treated (similar to herbicide strips). Use the formula in the General Information section for calculating the amount of Ridomil Gold GR needed per acre of trees. Soil surface applications of Ridomil Gold GR will not be effective until the fungicide is moved into the root zone by rainfall or irrigation.

**Notes:** (1) Unless labeled for bearing tree crops, do not apply to plantings that will bear harvestable fruit within 12 months of the last application, or possible illegal residues may result. (2) To avoid injury, do not treat the roots directly or concentrate the granules around the tree trunks. (3) Do not apply to trees under stress. (4) For intense plantings (2-3 times the normal planting rate) make applications on a per area basis, i.e., per acre or 1,000 sq. ft. Do not calculate the amount of Ridomil Gold GR on a per tree basis. (5) In CA, do not apply Ridomil Gold GR to newly planted trees within 90 days of planting. (6) Do not apply more than 240 lbs. of Ridomil Gold GR per treated acre per year.



# Ridomil Gold® GR

8/11

**Pythium and Phytophthora:** For full-season control apply 6 oz./1,000 linear ft. of row in a 7-inch band over the row or as a seed furrow application at the time of planting. For early- to mid-season control only, apply 1½-3 oz./1,000 linear ft. of row in the **seed furrow**. When making seed furrow applications, position the applicator tubes so the granules are mixed with the soil covering the seed or are applied in a "T-band." Use the following table to determine the amount of Ridomil Gold GR needed per acre based on the row spacing. Use the high rate in areas with a history of Phytophthora problems.

Suggested Rates Per Acre According to Row Spacing				
Row Spacing	Linear Ft. of Soybean Row Per Acre	Pounds of Ridomil Gold GR Needed Per Acre		
		Rate Desired in Oz.		
		1½ oz.	3 oz.	6 oz.
38 in.	13,756	1½ lbs.	2¾ lbs.	5½ lbs.
36 in.	14,520	1½ lbs.	2¾ lbs.	5½ lbs.
30 in.	17,424	1½ lbs.	3¼ lbs.	6½ lbs.
24 in.	21,780	2 lbs.	4¼ lbs.	8½ lbs.
20 in.	26,136	2½ lbs.	4¾ lbs.	9¾ lbs.

**Note:** Ridomil Gold GR is specific for Pythium and Phytophthora and will not control other diseases that may attack soybeans.

### Suggested Application Settings for a 7-inch Band Width or in the Seed Furrow

Applicator Brand	Speed MPH	Ounces/1,000 ft.				
		1.5	3	4	5	6
Gandy	4	8	13	15	17	19
	5	9	15	17	19	21
	6	10	16	19	21	23
	7	12	18	20	23	25
Allis Chalmers 1979-1985 International Harvester 1980 and Before	4	1	1.3	1.4	1.5	1.8
	5	1.1	1.4	1.5	1.9	2
	6	1.2	1.5	1.8	2	2.3
	7	1.3	1.5	1.9	2.2	2.5
Noble (Remcor) 1973-Present White 5100 Series Deutz Allis 1985-Present	4	2	4	5	6	7
	5	2 or 3	5	6	7	8 or 9
	6	3	5	7	8 or 9	10
	7	4	6	8	10	12
John Deere 7200 and 7300 MaxEmerge 2® White 6100 Series	4	8	12	14	16	18
	5	9	13	16	19	22
	6	10	15	18	22	26
	7	11	16	20	26	30
John Deere 7000 and 7100 Max-Emerge® Meter Scale 0...10...20	4	5	8	10	12	14
	5	6	10	12	14	16
	6	6 or 7	11	14	16	19
	7	7	12	15	19	22
John Deere 1979 and Later Meter Scale 0...5...15	4	4	7	8	9	10
	5	5	7 or 8	9	10	12
	6	5 or 6	8	10	12	14
	7	6	9	11	14	16
International Harvester 1981-1985 Case IH 1985-Present	4	1	1.4	1.55	1.75	2.1
	5	1.1	1.5	1.75	1.85	2.2
	6	1.2	1.65	1.85	2	2.3
	7	1.3	1.8	1.9	2.3	2.5



**SPINACH**

Ridomil Gold GR applied as a soil application will control damping-off in spinach caused by *Pythium* spp., white rust (*Albugo occidentalis*) and downy mildew. Applications may be made preplant incorporated or preemergence.

**Preplant Incorporated Application:** Apply 20-40 lbs. per treated acre as a broadcast soil application and incorporate in the top 2 inches of soil. For banded applications, a 7-inch band is recommended. Use the formula in the **General Information** section of this label to calculate the amount of Ridomil Gold GR needed per acre. When incorporated applications are made at the time of planting, position the drop tubes so the granules are mixed with the soil, covering the seed at a rate of 10-20 oz. of granules per 1,000 linear ft. of row. Avoid placing the granules directly in contact with the seed or injury may occur.

**Surface Application:** Apply 20-40 lbs. per treated acre uniformly to the soil surface at the time of planting. For banded applications, a 7-inch band is recommended. Use the formula in the **General Information** section of this label to calculate the amount of Ridomil Gold GR needed per acre. If natural rainfall is not expected before the seeds start germinating, Ridomil Gold GR should be incorporated mechanically before planting or be moved into the seed zone after planting with 1/2-1 inch sprinkler irrigation.

**White Rust and Downy Mildew Control:** Make the first application at the time of planting according to the above directions. Make up to 2 additional applications of 5 lbs. of Ridomil Gold GR per acre of crop as a side dress application. Make the first side dress application 21 days after planting or after the first cutting. One additional application may be made after the next cutting.

**Notes:** (1) White rust and downy mildew can only be controlled in a preventative disease control program that begins with an application of Ridomil Gold GR or Ridomil Gold EC to the soil at planting. If Ridomil Gold is not used at planting, do not use Ridomil Gold at any other time throughout the season. Do not apply Ridomil Gold GR where rust or downy mildew infections are already established. The use of Ridomil Gold GR in curative applications greatly increases the risk of the fungus developing insensitivity to this active ingredient. The development of insensitivity will destroy the effectiveness of Ridomil Gold in controlling white rust. (2) To avoid possible illegal residues, do not harvest spinach within 21 days of a Ridomil Gold GR application. (3) Do not use Ridomil Gold GR for disease control in greenhouse or field-grown vegetable bedding plants.

**SUGAR BEETS**

Ridomil Gold GR is a systemic fungicide for use in the control of diseases caused by *Pythium* spp. Applications may be made preplant incorporated or as a surface application at planting.

**Preplant Incorporated Application:** Apply 20-40 lbs. per treated acre as a broadcast soil application and incorporate into the top 2 inches of soil. For banded applications, a 7-inch band is recommended. Use the formula in the **General Information** section of this label to calculate the amount of Ridomil Gold GR needed per acre.

**Surface Application:** Apply 20-40 lbs. per treated acre at planting. For banded application, a 7-inch band is recommended. Use the formula in the **General Information** section of this label to calculate the amount of Ridomil Gold GR needed per acre. If natural rainfall is not expected before the seeds begin germinating, Ridomil Gold GR should be incorporated mechanically before planting or be moved into the seed zone after planting with 1/2-1 inch sprinkler irrigation.

**TOMATOES**

Soil applications of Ridomil Gold GR at planting will provide control of damping-off caused by *Pythium* spp. and soil applications applied 4-12 weeks before harvest under the vines will reduce fruit and root rot caused by *Pythium* spp. and *Phytophthora* spp.

**Damping-Off (*Pythium* spp.):** Apply 20 lbs. per treated acre uniformly over the surface of the soil at the time of planting. If natural rainfall is not expected before the seeds begin germinating, Ridomil Gold GR should be incorporated mechanically with 1/2-1 inch sprinkler irrigation. For banded applications, a 7-inch band is recommended. Use the formula in the **General Information** section of this label to calculate the amount of Ridomil Gold GR needed per acre.

**Fruit and Root Rot (*Pythium* spp. and *Phytophthora* spp.):** Additional applications can be made as soil surface applications under the vines to provide control of root and fruit rot. If only one application is planned, apply 20 lbs. per treated acre under the vines 4 weeks after the planting application to 4 weeks before harvest, depending on the expected occurrence of the diseases. If root and fruit rot is a problem throughout the growing season or the occurrence of diseases cannot be predicted, make 2 additional applications at 10 lbs. per treated acre under the vines, one at 4-6 weeks after the planting application and another 4 weeks before harvest. If rainfall is not expected within 3 days of the application, follow as soon as possible with 1/2-1 inch overhead irrigation. Use the formula shown in the **General Information** section to determine the amount of Ridomil Gold GR needed per acre.

Ridomil Gold GR can be shanked into the beds with fertilizer. For calculating the amount of Ridomil Gold GR needed per acre, apply the same amount as would be applied on a 7-inch band at 20 lbs. per treated acre. To obtain uniform disease control, Ridomil Gold GR must be mixed uniformly with the fertilizer.

# Ridomil Gold® GR

10/11

**Notes:** To avoid possible illegal residues, (1) Do not apply more than 40 lbs. per treated acre per season, and (2) Do not apply within 7 days of harvest. (3) Do not use Ridomil Gold GR for disease control in greenhouse or field-grown vegetable bedding plants.

## REPLANTING

If replanting is necessary, additional applications of Ridomil Gold GR can be made, provided that the total amount of active ingredient in Ridomil Gold GR applied does not exceed the maximum specified on the product label for the particular crop.

## ROTATION (PLANTBACK) RESTRICTIONS

Do not plant any crop which is not registered for use with the Ridomil Gold active ingredient in soil treated with this active ingredient for a period of 12 months. See the following list.

Rotation Crop	Planting Time From Last Ridomil Gold GR Application (Days)
Alfalfa (including birdsfoot trefoil), Almonds, Apples, Asparagus, Avocados	0 days
Blueberries, Broccoli	
Cabbage, Cauliflower, Chinese Broccoli (gai lan, white flowering broccoli), Chinese Cabbage (tight-heading varieties only), Citrus, Cotton, Cranberries, Cucurbit Vegetables	
Deciduous Fruits and Nuts*	
Eggplant	
Garlic, Ginseng, Grapes	
Hops	
Leafy Vegetables (excluding Brassica) Legume Vegetables (beans and peas-succulent and dried)	
Onions (dry bulb, green, and seed)	
Papaya, Peanuts, Peppers, Pineapples, Potatoes	
Raspberries, Root and Tuber Vegetables	
Soybeans, Spinach, Stone Fruits, Strawberries, Sugar Beets	
Tobacco, Tomatoes	
Walnuts	
Wheat, barley, and oats	360 days
Corn	360 days
Crops Not Intended for Food or Feed	0 days
All Other Crops Intended for Food or Feed	360 days

\*These crops and other perennial crops may be planted immediately following last application of Ridomil Gold GR provided they will not bear harvestable fruit within 12 months.

## STORAGE AND DISPOSAL

Store in a dry place. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.

### Pesticide Disposal

Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide is a violation of federal law. Pesticide that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

### Container Disposal

Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### CAUTION

Causes eye irritation. Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust.

### Statement of Practical Treatment

**If in eyes:** Flush eyes with plenty of water. Get medical attention if irritation persists.

**If on skin:** Wash with plenty of soap and water. Get medical attention.

**If inhaled:** Move to fresh air.

**Note to Physician:** If ingested, induce emesis or lavage stomach. Treat symptomatically.

### Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### Environmental Hazards

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

Ridomil Gold® trademark of Novartis

MaxEmerge 2® and Max-Emerge® trademarks of John Deere

Terraclor® trademark of Uniroyal Chemical Company, Inc.

©1997 Novartis

Novartis Crop Protection, Inc.  
Greensboro, NC 27419

NCP 166L1A 0897