



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

June 21, 2023

Nestor Algarin  
Regulatory Product Manager  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, NC 27419

Subject: PRIA Label Amendment – **Mefenoxam**: Crop group conversions/revisions for Cottonseed, subgroup 20C; Leaf petiole subgroup 22B; Leafy vegetable, Crop Group 4-16 (except spinach); Vegetable, Brassica, head and stem, group 5-16; Vegetable, fruiting, group 8-10; Vegetable, legume, bean, succulent shelled, subgroup 6-22C; Vegetable, legume, pea, succulent shelled, subgroup 6-22D; Vegetable, stalk and stem, subgroup 22A.

Product Name: Apron XL LS  
EPA Registration Number: 100-799  
Application Dates: 11/23/2021, 11/24/2021  
Petition Number: 1F8970  
Decision Numbers: 580474, 580472

Dear Nestor Algarin:

The application referred to above, submitted under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable under FIFRA section 3(c)(5).

You must submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements


Page 2 of 2  
EPA Reg. No. 100-799  
Decision No. 580474, 580472

EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Stephanie Suarez at 202-566-2918 or at [Suarez.Stephanie@epa.gov](mailto:Suarez.Stephanie@epa.gov).

Sincerely,



Cynthia Giles-Parker, Chief  
Fungicide Branch  
Registration Division (7505T)

Enclosure – stamped “accepted” label

[Master Label]

<b>MEFENOXAM</b>	<b>GROUP</b>	<b>4</b>	<b>FUNGICIDE</b>
------------------	--------------	----------	------------------

## Apron XL® LS

### Fungicide

A seed treatment product for protection against *Pythium* and *Phytophthora* causing damping-off, seed rot, and systemic downy mildew diseases of certain crops

Active Ingredients:

Mefenoxam* .....	33.3%
------------------	-------

Other Ingredients:	66.7%
--------------------	-------

Total:	100.0%
--------	--------

\*CAS Nos. 70630-17-0 and 69516-34-3

Apron XL LS is a flowable concentrate for seed treatment containing 3.0 pounds of mefenoxam per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)


See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-799

EPA Est.

\_\_\_\_\_ gallons

Net Contents



[Batch Code: \_\_\_\_\_ ] (For nonrefillables only.)

## **TABLE OF CONTENTS**

### **1.0 FIRST AID**

### **2.0 PRECAUTIONARY STATEMENTS**

#### **2.1 Hazards to Humans and Domestic Animals**

#### **2.2 Personal Protective Equipment (PPE)**

2.2.1 User Safety Requirements

2.2.2 Engineering Controls

2.2.3 User Safety Recommendations

#### **2.3 Environmental Hazards**

2.3.1 Groundwater Advisory

2.3.2 Surface Water Advisory

## **DIRECTIONS FOR USE**

### **3.0 PRODUCT INFORMATION**

#### **3.1 Resistance Management**

### **4.0 APPLICATION DIRECTIONS**

#### **4.1 Tank Mixtures**

### **5.0 ROTATIONAL CROP RESTRICTIONS**

### **6.0 SEED CONTAINER LABEL REQUIREMENTS**

### **7.0 SEED TREATMENT DIRECTIONS – DOMESTIC USE**

#### **7.1 Artichoke, Globe**

#### **7.2 Brassica Head and Stem Vegetable Crop Group 5-16**

#### **7.3 Bulb Vegetable Crop Group 3-07**

#### **7.4 Cereal Grains**

7.4.1 Barley, Buckwheat, Millet (Pearl and Proso), Oats, Rice, Rye, Triticale, Wheat and Wild Rice

7.4.2 Corn

7.4.3 Sorghum

#### **7.5 Cotton, Crop Subgroup 20C**

#### **7.6 Cucurbit Vegetables Crop Group 9**

#### **7.7 Fruiting Vegetable Crop Group 8-10**

#### **7.8 Grass Forage, Fodder, and Hay Crop Group 17**

#### **7.9 Leaf Petiole Vegetable Crop Subgroup 22B**

#### **7.10 Leafy Vegetable Crop Group 4-16**

#### **7.11 Leaves of Root and Tuber Vegetables (Human Food or Animal Feed) Crop Group 2**

#### **7.12 Legume Vegetable Group – Crop Group 6-22, except Soybean (See Soybean) and Forage and Hay of Legume Vegetable Group – Crop Group 7-22**

#### **7.13 Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay) Crop Group 18**

#### **7.14 Peanut**

#### **7.15 Rapeseed Crop Subgroup 20A**

#### **7.16 Root and Tuber Vegetables Crop Group 1, except Potato**

#### **7.17 Soybean**

#### **7.18 Sunflower**

**7.19 Turfgrass**

**8.0 SEED CONTAINER LABEL REQUIREMENTS – EXPORT USE**

**9.0 SEED TREATMENT DIRECTIONS – EXPORT USE**

**9.1 Brassica Head and Stem Vegetable Crop Group 5-16**

**9.2 Bulb Vegetables Crop Group 3-07**

**9.3 Cereal Grains**

9.3.1 Corn

9.3.2 Millet

9.3.3 Sorghum

**9.4 Cucurbit Vegetables Crop Group 9**

**9.5 Fruiting Vegetable Crop Group 8-10**

**9.6 Leaf Petiole Vegetable Crop Subgroup 22B**

**9.7 Leafy Vegetable Crop Group 4-16**

**9.8 Peas**

**9.9 Root and Tuber Vegetables Crop Group 1, except Potato**

**9.10 Sunflower**

**10.0 STORAGE AND DISPOSAL**

**11.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

## 1.0 FIRST AID

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<p><b>HOTLINE NUMBER</b>            For 24-Hour Medical Emergency Assistance (Human or Animal) or            Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),            Call  <b>1-800-888-8372</b></p>	

## 2.0 PRECAUTIONARY STATEMENTS

### 2.1 Hazards to Humans and Domestic Animals

#### **WARNING/AVISO**

Causes substantial but temporary eye irritation. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

### 2.2 Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥14 mils, neoprene rubber ≥14 mils, nitrile rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, Viton™ ≥14 mils
- Shoes plus socks
- Protective eyewear: faceshield, goggles, or safety glasses

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### 2.2.1 User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### 2.2.2 Engineering Controls

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

### 2.2.3 User Safety Recommendations

#### **User Safety Recommendations**

##### **Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## 2.3 Environmental Hazards

Do not contaminate water when disposing of equipment washwater or rinsate. Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting, including in row ends.

### 2.3.1 Groundwater Advisory

Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### 2.3.2 Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apron XL LS may only be used in accordance with directions on this label or separately published EPA approved supplemental labeling for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Then conduct germination tests with a portion of this treated seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crops when treated with Apron XL LS.

**Restriction: Maximum usage when applying metalaxyl- and mefenoxam- containing products to the same crop within the same season:** Do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.**



### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.** Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, polyvinyl chloride (PVC)  $\geq 14$  mils, or Viton  $\geq 14$  mils
- Shoes plus socks
- Protective eyewear

## 3.0 PRODUCT INFORMATION

Apron XL LS is a systemic fungicide seed treatment specifically for protection against systemic downy mildews and diseases caused by soil-borne *Pythium* and *Phytophthora* spp. When a rate range is specified, use higher specified rates of Apron XL LS when the disease pressure is expected to be high.

For protection against other soil-borne diseases, including those caused by *Fusarium* spp. and *Rhizoctonia* spp., Apron XL LS must be applied in combination with other registered seed treatment fungicides.

Apply Apron XL LS as a water-based slurry through standard liquid-type seed treatment equipment that provides uniform seed coverage. It is advised to use seed treaters with atomizers or spinning discs for better product coverage on the seed.

### 3.1 Resistance Management

Apron XL LS contains mefenoxam, a Group 4 fungicide. Any fungal population may contain individuals naturally resistant to Apron XL LS and other Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Follow appropriate resistance-management strategies.

Mefenoxam belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Apron XL LS or other Group 4 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM guidance for specific crops and pathogens.

- For further information or to report suspected resistance contact Syngenta at 1-866-796-4368. You can also contact your pesticide distributor or university extension specialist to report resistance.

Syngenta encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label.

## 4.0 APPLICATION DIRECTIONS

**Important:** Recirculate Apron XL LS thoroughly before using.

Follow the manufacturer's application instructions for the seed treatment equipment being used.

Apply Apron XL LS as a water-based slurry through standard liquid-type seed treatment equipment that provides uniform seed coverage. It is advised to use seed treaters with atomizers or spinning discs for better product coverage on the seed. The total application volume must be sufficient to provide the desired level of coverage. Dilution is typically done with water or liquid inoculants. Thoroughly mix the specified amount of Apron XL LS into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used. Uneven or incomplete seed coverage may not give the desired level of disease control.

Certain crops require addition of inoculants when the seed is treated or planted. Apron XL LS is compatible with several liquid inoculant products. Consult the maker of the inoculant product and a Syngenta representative for directions before applying Apron XL LS with inoculants.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution. Clean out any unused product from the treater after treating or maintain constant agitation if the leftover slurry will be maintained overnight.

### **Restrictions:**

- Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR Part 153.155(c).
- Allow seed to dry before bagging.

Follow planter manufacturer's specifications for use of talc or other hopper box additives at planting. Seed must be completely dry before adding to planter.

### 4.1 Tank Mixtures

Apron XL LS mixes easily with water and other water-based seed treatments. When mixing with products from other manufacturers, test the compatibility prior to use by conducting a jar test: Mix all intended seed treatments with a proportional amount of water to achieve the desired slurry mixture in a clear glass container. Mix well and allow mixture to sit for one hour. Remix and observe for incompatibility.

**Mixing Apron XL LS with tank-mix partners:** Add ½ of the required water to the mix tank and turn on the agitation. Mechanical agitation is preferred. If using wettable powders, add them first to clean water allowing them to completely disperse prior to adding Apron XL LS or other products. Allow each tank-mix partner to completely disperse before adding the next product. Add the remaining amount of water and agitate. Maintain agitation until the entire slurry mixture has been used.

For protection against other soil-borne diseases, including those caused by *Fusarium* spp. and *Rhizoctonia* spp., apply Apron XL LS in combination with other registered seed treatment fungicides.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**USE RESTRICTION:** When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored with an EPA approved dye including one of the dyes listed in 40 CFR Sections 180.910 and 180.920 to distinguish and prevent subsequent inadvertent use as food for man or feed for animals.

## 5.0 ROTATIONAL CROP RESTRICTIONS

In the event of a crop failure or harvest of a crop grown from seed treated with Apron XL LS, the field may be replanted immediately with any crops.

## 6.0 SEED CONTAINER LABEL REQUIREMENTS

THE FEDERAL SEED ACT REQUIRES THAT CONTAINERS OF SEEDS TREATED WITH APRON XL LS MUST BE LABELED WITH THE FOLLOWING STATEMENTS:

- This seed has been treated with mefenoxam fungicide.
- **DO NOT** use treated seed for feed, food, or oil purposes.

THE US ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON CONTAINERS OF SEEDS TREATED WITH APRON XL LS:

- Store away from feed and foodstuffs.
- **DO NOT** allow children, pets or livestock to have access to treated seeds.
- Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting, including in row ends.
- Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves when handling treated seed.
- Dispose of excess treated seed by burying seed away from bodies of water.
- **DO NOT** contaminate water bodies when disposing of planting equipment washwater.
- Dispose of seed packaging in accordance with local requirements.

- Excess treated seed may be used for ethanol production only if: (1) By-products are not used for livestock feed, and (2) No measurable residues of pesticides remain in ethanol by-products that are used for agronomic practice.
- Treated seed must be incorporated into the soil at the advised depth.
- **Groundwater Advisory:** Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- **For Brassica Head and Stem Vegetables, Bulb Vegetables, Cucurbits Vegetables and Fruiting Vegetables: DO NOT** apply a pre-plant or an at-plant application of mefenoxam or metalaxyl products to seed treated with Apron XL LS.
- **For Bulb Vegetables: DO NOT** apply more than 0.0078 lb ai/A (3.5 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment
- **For Grass Forage, Fodder and Hay: DO NOT** apply more than 0.0081 lb ai/A (3.7 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment
- **For Legume Vegetables (Succulent or Dried) and Foliage of Legume Vegetables, except Soybean: DO NOT** apply more than 0.025 lb ai/A (115 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment
- **For Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay): DO NOT** apply more than 0.0047 lb ai/A (2.1 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment
- **For Peanuts: DO NOT** apply more than 0.071 lb ai/A (32 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment
- **For Root and Tuber Vegetables: DO NOT** apply more than 0.020 lb ai/A (9.2 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment
- **For Soybean: DO NOT** apply more than 0.052 lb ai/A (23 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment
- **For Spinach: DO NOT** use seed for spinach to be grown in the greenhouse or inside environmentally-controlled rooms.

## 7.0 SEED TREATMENT DIRECTIONS – DOMESTIC USE

### 7.1 Artichoke, Globe

Crop			
Artichoke, Globe			
Target Diseases	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off and early-season Phytophthora root rot	0.32 – 2	0.008 - 0.048	0.003 – 0.019
Additional Information			
<ul style="list-style-type: none"> <li>Based on an average seed weight of 11,500 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<b>Maximum Annual Application Rate:</b> DO NOT apply more than 0.0063 lb ai/A/year (0.126 lb ai/100 lb seed) of mefenoxam- or metalaxyl-containing products as a seed treatment.			
<b>Resistance Management:</b> Refer to <b>Section 3.1.</b>			

### 7.2 Brassica Head and Stem Vegetable Crop Group 5-16

Crops			
Broccoli		Cauliflower	
Brussels sprouts		Cultivars, varieties, and hybrids of these commodities	
Cabbage			
Cabbage, Chinese (napa)			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.085 - 0.64	0.002 – 0.015	0.0001 – 0.0007
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 100,000 seeds/lb</li> <li>For seeds sown for transplant production: Use Apron XL LS with a labeled rate of azoxystrobin (e.g., Dynasty®, EPA Reg. No. 100-1159).</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>DO NOT</b> apply a pre-plant or an at-plant application of mefenoxam or metalaxyl products to seed treated with Apron XL LS. Mefenoxam foliar/in-furrow products (e.g., Ridomil Gold® SL, EPA Reg. No. 100-1202) are allowed to be applied beginning 6 weeks after transplanting.</li> <li><b>Maximum Annual Application Rate:</b> DO NOT apply more than 0.00082 lb ai/A/year (0.031 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
<b>Resistance Management:</b> Refer to <b>Section 3.1.</b>			

## 7.3 Bulb Vegetable Crop Group 3-07

Crops			
Chive, fresh leaves		Onion, Beltsville bunching	
Chive, Chinese, fresh leaves		Onion, bulb	
Daylily, bulb		Onion, Chinese, bulb	
Elegans hosta		Onion, fresh	
Fritillaria, bulb		Onion, green	
Fritillaria, leaves		Onion, macrostem	
Garlic, bulb		Onion, pearl	
Garlic, great-headed, bulb		Onion, potato, bulb	
Garlic, Serpent, bulb		Onion, tree, tops	
Kurrat		Onion, Welsh, tops	
Lady's leek		Shallot, bulb	
Leek		Shallot, fresh leaves	
Leek, wild		Cultivars, varieties, and/or hybrids of these	
Lily, bulb			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/ seed*
Pythium damping-off	0.085 - 0.64	0.002 – 0.015	0.0001 – 0.0007
Additional Information			
<ul style="list-style-type: none"> <li>* based on an average seed weight of 100,000 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.0078 lb ai/A (0.031 lb ai/100 lb seed) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment on Bulb Vegetable Crop Group 3-07</li> </ul>			
Resistance Management:			
Refer to <b>Section 3.1</b> .			

## 7.4 Cereal Grains

### 7.4.1 Barley, Buckwheat, Millet (Pearl and Proso), Oats, Rice, Rye, Triticale, Wheat and Wild Rice

<b>Crops</b>			
Barley		Rice	
Buckwheat		Rye	
Millet, Pearl		Triticale	
Millet, Proso		Wheat	
Oats		Wild Rice	
<b>Target Diseases</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed*</b>
<b><u>Barley, Buckwheat, Pearl Millet, Proso Millet, Oats, Rice, Rye, Triticale, Wheat, and Wild Rice</u></b> Pythium damping-off	0.12 - 0.64	0.003 – 0.015	0.00087 – 0.0047
<b><u>Millet</u></b> Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.0023 – 0.0047
Systemic downy mildew	1.28	0.031	0.0093
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 15,000 seeds/lb</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.097 lb ai/A/year (0.062 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
<b>Resistance Management:</b>			
Refer to <b>Section 3.1</b> .			



## 7.4.2 Corn

<b>Crops</b>			
Field Corn Sweet Corn Popcorn			
<b>Target Diseases</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed</b>
<b>Field Corn and Popcorn</b> Pythium seed rot and damping-off	0.0425 - 0.085	0.001 – 0.002	0.0025 – 0.005*
<b>Sweet Corn</b> Pythium seed rot and damping-off	0.32 - 0.64	0.008 – 0.015	0.011 – 0.022**
Systemic downy mildew	1.28	0.031	0.044**
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>• *based on an average seed weight of 1,800 corn seeds/lb</li> <li>• **based on an average seed weight of 3,200 sweet corn seeds/lb</li> <li>• For a greater spectrum of seedling disease protection of corn, combine Apron XL LS with azoxystrobin (e.g., Dynasty, EPA Reg. No. 100-1159), fludioxonil (e.g., Maxim® 4FS, EPA Reg. No. 100-758), picarbutrazox (Vayantis®, EPA Reg. No. 100-1635), or sedaxane (e.g., Vibrance®, EPA Reg. No. 100-1374).</li> <li>• All sweet corn seed treated with Apron XL LS must be planted and not carried over to the following year.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li>• <b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.0012 lb ai/A/year (0.004 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
<b>Resistance Management:</b>			
Refer to <b>Section 3.1</b> .			

### 7.4.3 Sorghum

<b>Crop</b>			
Sorghum			
<b>Target Diseases</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed*</b>
Pythium damping-off on all sorghum cultivars	0.32 - 0.64	0.008 – 0.015	0.0024 - 0.0048
Pythium damping-off when disease pressure is expected to be low	0.085 - 0.32	0.002 – 0.008	0.0006 - 0.0024
Systemic downy mildew on cultivars which possess sorghum downy mildew resistance	0.32 - 0.64	0.008 – 0.015	0.0024 - 0.0048
Systemic downy mildew on cultivars which do not possess resistance to a particular race of sorghum downy mildew grown in areas with recent history of severe disease	1.28	0.031	0.0096
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>• *based on an average seed weight of 14,500 sorghum seeds/lb</li> <li>• Only treat high-quality sorghum seed (90% germination and above) with Apron XL LS.</li> <li>• Seed treated with the 1.28 fl oz rate must be planted and not carried over to the following year.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li>• <b>Maximum Annual Application Rate: DO NOT</b> apply more than 0.0074 lb ai/A/year (0.062 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
<b>Resistance Management:</b>			
Refer to <b>Section 3.1</b> .			

## 7.5 Cotton, Crop Subgroup 20C

Crop			
Cotton			
Target Diseases	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium seed rot and damping-off	0.32 - 1.14	0.008 – 0.027	0.008 – 0.028
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 4.500 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<b>Maximum Annual Application Rate: DO NOT</b> apply more than 0.0104 lb ai/A/year (0.055 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.			
<b>Resistance Management:</b> Refer to <b>Section 3.1.</b>			

## 7.6 Cucurbit Vegetables Crop Group 9

Crops			
Chayote (fruit)	<i>Momordica</i> spp.	Pumpkin	
Chinese waxgourd (Chinese preserving melon)	Balsam apple Balsam pear Bitter melon	Squash, summer ( <i>Cucurbita pepo</i> var. <i>meloepo</i> ) Crookneck squash	
Citron melon	Chinese cucumber	Scallop squash	
Cucumber	Muskmelon	Straightneck squash	
Gherkin	(hybrids and/or cultivars of	Vegetable marrow	
Gourd, edible ( <i>Lagenaria</i> spp.)	<i>Cucumis melo</i> ) Cantaloupe	Zucchini	
Hyotan	Casaba	Squash, winter ( <i>Cucurbita maxima</i> , <i>C. moschata</i> )	
Cucuzza	Crenshaw melon	Butternut squash	
Gourd, edible ( <i>Luffa</i> <i>acutangula</i> , <i>L. cylindrical</i> )	Golden pershaw melon	Calabaza	
Hechima	Honeydew melon	Hubbard squash	
Chinese okra	Honey balls	Squash, winter ( <i>Cucurbita mixta</i> , <i>C. pepo</i> ):	
	Mango melon	Acorn squash	
	Persian melon	Spaghetti squash	
	Pineapple melon	Watermelon (hybrids and/or varieties of <i>Citrullus lanatus</i> )	
	Santa Claus melon		
	Snake melon		
	True cantaloupe		
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.085 - 0.64	0.002 – 0.015	0.0006 – 0.0047
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 15,000 seeds/lb</li> <li>For seeds sown for transplant production: Use Apron XL LS with a labeled rate of azoxystrobin (e.g., Dynasty, EPA Reg. No. 100-1159).</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.0078 lb ai/A/year (0.031 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> <li><b>DO NOT</b> apply a pre-plant or an at-plant application of mefenoxam or metalaxyl products to seed treated with Apron XL LS. Mefenoxam foliar/in-furrow products (e.g., Ridomil Gold SL, EPA Reg. No. 100-1202) are allowed to be applied beginning 6 weeks after transplanting.</li> </ul>			
Resistance Management:			
Refer to <b>Section 3.1</b> .			

## 7.7 Fruiting Vegetable Crop Group 8-10

Crops			
African eggplant	Groundcherry	Pepper, non-bell	
Bush tomato	Martynia	Roselle	
Cocona	Naranjilla	Scarlet eggplant	
Currant tomato	Okra	Sunberry	
Eggplant	Pea eggplant	Tomatillo	
Garden huckleberry	Pepino	Tomato	
Goji berry	Pepper, bell	Tree Tomato	
Cultivars, varieties and/or hybrids of these			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.085 - 0.64	0.002 – 0.015	0.00006 – 0.0005
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 150,000 seeds/lb</li> <li>For seeds sown for transplant production: Use Apron XL LS with a labeled rate of azoxystrobin (e.g., Dynasty, EPA Reg. No. 100-1159).</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.0047 lb ai/A/year (0.031 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> <li><b>DO NOT</b> apply a pre-plant or an at-plant application of mefenoxam or metalaxyl products to seed treated with Apron XL LS. Mefenoxam foliar/in-furrow products (e.g., Ridomil Gold SL, EPA Reg. No. 100-1202) are allowed to be applied beginning 6 weeks after transplanting.</li> </ul>			
Resistance Management:			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1</b>.</li> </ul>			

## 7.8 Grass Forage, Fodder, and Hay Crop Group 17

Crops			
Graminae Family (except Sugarcane and crops included in Forage, Fodder, and Straw of Cereal Grains Crop Group 16)			
<ul style="list-style-type: none"> <li>• Green</li> <li>• Cured</li> </ul>			
Grasses Grown for Hay or Silage			
All Pasture and Range Grasses			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.085 - 0.64	0.002 – 0.015	0.00004 – 0.00033
Additional Information			
<ul style="list-style-type: none"> <li>• *based on average seed weight of 210,000 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>• <b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.0081 lb ai/A (3.7 g ai/A) per calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment on Grass Forage, Fodder, and Hay Crop Group 17.</li> </ul>			
Resistance Management:			
Refer to <b>Section 3.1</b> .			

## 7.9 Leaf Petiole Vegetable Crop Subgroup 22B

Crops			
Cardoon	Fuki		
Celery	Kohlrabi		
Celery, Chinese	Rhubarb		
Celtuce	Udo		
Fennel, Florence (finocchio)	Zuiki		
	Cultivars, varieties, and hybrids of the commodities in crop subgroup 22B		
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.085 - 0.64	0.002 – 0.015	0.000009 – 0.00007
Additional Information			
<ul style="list-style-type: none"> <li>• *based on an average seed weight of 1,000,000 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>• <b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.00002 lb ai/A/year (0.031 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
Resistance Management:			
<ul style="list-style-type: none"> <li>• Refer to <b>Section 3.1</b>.</li> </ul>			

## 7.10 Leafy Vegetable Crop Group 4-16

Crops			
Amaranth, Chinese	Cress, garden	Maca, leaves	
Amaranth, Leafy	Cress, upland ( <i>yellow rocket</i> , <i>winter cress</i> )	Mizuna	
Arugula (Roquette)	Dandelion, leaves	Mustard green	
Aster, Indian	Dock ( <i>sorrel</i> )	Orach	
Blackjack	Dol-nam-mul	Parsley, fresh leaves	
Broccoli, Chinese (gai lon)	Ebolo	Plantain, Buckhorn	
Broccoli Raab (rapini)	Endive ( <i>escarole</i> )	Primrose, English	
Cabbage, Abyssinian	Escarole	Purslane, Garden	
Cabbage, Chinese (bok choy)	Fameflower	Purslane, Winter	
Cabbage, seakale	Feather Cockscomb	Radish, leaves	
Cat's Whiskers	Good King Henry	Rape Greens	
Cham-chwi	Hanover Salad	Rocket, wild	
Cham-na-mul	Huauzontle	Shepherd's Purse	
Chervil, fresh leaves	Jute, leaves	Spinach	
Chrysanthemum, garland	Kale	Spinach, Malabar	
Cilantro, fresh leaves	Lettuce, Bitter	Spinach, New Zealand	
Collards	Lettuce, Head	Spinach, Tanier	
Corn Salad	Lettuce, Leaf	Swiss chard	
Cosmos		Turnip Greens	
		Violet, Chinese, leaves	
		Cultivars, varieties, and hybrids of these commodities	
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.085 - 0.64	0.002 – 0.015	0.00002 – 0.00018
Additional Information			
<ul style="list-style-type: none"> <li>*based on average seed weight of 400,000 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.0194 lb ai/A/year (0.078 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> <li><b>DO NOT APPLY</b> Apron XL LS to spinach that is to be grown in the greenhouse or enclosed environmentally-controlled rooms.</li> <li>Containers or bags of spinach treated with Apron XL LS must be labeled as follows: <b>“DO NOT USE SEED FOR SPINACH TO BE GROWN IN THE GREENHOUSE OR INSIDE ENVIRONMENTALLY-CONTROLLED ROOMS.”</b></li> </ul>			
Resistance Management:			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1.</b></li> </ul>			

## 7.11 Leaves of Root and Tuber Vegetables (Human Food or Animal Feed) Crop Group 2

Crops			
Beet, Garden		Parsnip	
Beet, Sugar		Radish	
Burdock, edible		Radish, Oriental (daikon)	
Carrot		Rutabaga	
Cassava, Bitter and Sweet		Salsify, Black	
Celeriac (celery root)		Sweet Potato	
Chervil, Turnip-Rooted		Tanier(cocoyam)	
Chicory		Turnip	
Dasheen (taro)		Yam, True	
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.085 - 0.64	0.002 – 0.015	0.00003 – 0.00023
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 300,000 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.0203 lb ai/A/year (0.062 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
Resistance Management:			
Refer to <b>Section 3.1</b> .			



## 7.12 Legume Vegetable Group – Crop Group 6-22, except Soybean (See Soybean) and Forage and Hay of Legume Vegetable Group – Crop Group 7-22

Crops			
African yam bean, dry seed American potato bean, dry seed Bean ( <i>Lupinus</i> spp.), succulent shelled and dry seed Andean lupin blue lupin grain lupin sweet lupin white lupin white sweet lupin yellow lupin Bean ( <i>Phaseolus</i> spp.), edible podded French bean garden bean green bean kidney bean navy bean scarlet runner bean snap bean wax bean Bean ( <i>Phaseolus</i> spp.), succulent shelled lima bean scarlet runner bean wax bean Bean ( <i>Phaseolus</i> spp.), dry seed black bean cranberry bean dry bean field bean French bean garden bean great northern bean green bean kidney bean lima bean navy bean pink bean pinto bean red bean scarlet runner bean tepary bean yellow bean	Bean ( <i>Vigna</i> spp.), edible podded asparagus bean catjang bean Chinese longbean cowpea moth bean mung bean rice bean urd bean yardlong bean Bean ( <i>Vigna</i> spp.), succulent shelled blackeyed pea catjang bean cowpea crowder pea moth bean southern pea Bean ( <i>Vigna</i> spp.), dry seed adzuki bean asparagus bean blackeyed pea catjang bean Chinese longbean cowpea crowder pea moth bean mung bean rice bean southern pea urd bean yardlong bean Broad bean (fava bean), succulent shelled and dry seed Chickpea (garbanzo), edible podded, succulent shelled, and dry seed Goa bean (asparagus pea and winged bean), edible podded, succulent shelled, and dry seed Grass pea, edible podded and dry seed Guar bean, edible podded and dry seed	Horse gram, dry seed Jackbean, edible podded, succulent shelled and dry seed Lablab bean (hyacinth bean), edible podded, succulent shelled, and dry seed Lentil, edible podded, succulent shelled and dry seed Morama bean, dry seed Pea ( <i>Pisum</i> spp.), edible podded dwarf pea green pea snap pea snow pea sugar snap pea Pea ( <i>Pisum</i> spp.), succulent shelled English pea garden pea green pea Pea ( <i>Pisum</i> spp.), dry seed dry pea field pea garden pea yellow pea wrinkled pea marrowfat pea green pea Pigeon pea, edible podded, succulent shelled and dry seed Sword bean, edible podded and dry seed Vegetable soybean (edamame), edible podded and succulent shelled Velvetbean, edible podded, succulent shelled and dry seed Winged pea, edible podded and dry seed Cultivars, varieties, and/or hybrids of these commodities Plant parts of any legume vegetable listed in Crop Group 6-22 that will be used as animal feed.	
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.16 - 0.64	0.004 -0.015	0.014 – 0.056

<b>Field Beans, Kidney Beans, Navy Beans, Pinto Beans, and Lentils</b> Pythium damping-off when disease pressure is expected to be low in combination with other seed treatment products labeled for these crops	0.085 - 0.32	0.002 – 0.008	0.0074 – 0.028
Early-season Phytophthora	0.64	0.015	0.056
<b>Peas</b> Systemic downy mildew	1.28	0.031	0.11
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 1,250 seeds/lb</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate: DO NOT</b> apply more than 0.25 lb ai/A/year (0.062 lb ai/100 lb seed/year of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
<b>Resistance Management:</b> Refer to <b>Section 3.1</b> .			

## 7.13 Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay) Crop Group 18

Crops			
Alfalfa		Sainfoin	
Bean, velvet		Trefoil	
Clover		Vetch	
Kudzu		Vetch, crown	
Lespedeza		Vetch, milk	
Lupin			
Target Diseases	Use Rate		
	fl oz/100 lb of seed	lb ai/100 ;b seed	mg ai/seed*
Pythium damping-off			
Early-season Phytophthora	0.64	0.015	0.00033
<b>Alfalfa</b> Suppression of early-season downy mildew			
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 210,000 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> DO NOT apply more than 0.0047 lb ai/A/year (0.031 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
<b>Resistance Management:</b> Refer to <b>Section 3.1</b> .			

## 7.14 Peanut

Crops			
Peanut			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium seed rot and damping-off	0.16 - 0.64	0.004 – 0.015	0.0205 – 0.0822
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 850 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> DO NOT apply more than 0.071 lb ai/A/year (0.031 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
<b>Resistance Management:</b> Refer to <b>Section 3.1</b> .			

## 7.15 Rapeseed Crop Subgroup 20A

Crops			
Borage	Hare's ear mustard	Oil radish	
Crambe	Lesquerella	Poppy seed	
Cuphea	Lunaria	Rapeseed (canola)	
Echium	Meadowfoam	Sesame	
Flax seed	Milkweed	Sweet rocket	
Gold of pleasure	Mustard seed	Cultivars, varieties, and/or hybrids of these	
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.32	0.0008	0.00035
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 100,000 seeds/lb</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.0013 lb ai/A/year (0.015 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
Resistance Management:			
Refer to <b>Section 3.1</b> .			

## 7.16 Root and Tuber Vegetables Crop Group 1, except Potato

<b>Crops</b>			
Arracacha		Horseradish	
Arrowroot		Leren	
Artichoke, Chinese		Parsley, turnip-rooted	
Artichoke, Jerusalem		Parsnip	
Beet, garden		Radish	
Beet, sugar		Radish, oriental (daikon)	
Burdock, Edible		Rutabaga	
Canna, Edible (Queensland arrowroot)		Salsify (oyster plant)	
Carrot		Salsify, black	
Cassava, Bitter & Sweet		Salsify, Spanish	
Celeriac (celery root)		Skirret	
Chayote (root)		Sweet Potato	
Chervil, turnip-rooted		Tanier (cocoyam)	
Chicory		Turmeric	
Chufa		Turnip	
Dasheen (Taro)		Yam Bean (jicama, manioc pea)	
Ginger		Yam, True	
Ginseng			
<b>Target Disease</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed*</b>
Pythium damping-off	0.085 - 0.64	0.002 – 0.015	0.00003 – 0.00023
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 300,000 seeds/lb</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li><b>DO NOT</b> Apply Apron XL LS to potatoes.</li> <li><b>Maximum Annual Application Rate: DO NOT</b> apply more than 0.020 lb ai/A/year (0.062 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
<b>Resistance Management:</b>			
Refer to <b>Section 3.1</b> .			

## 7.17 Soybean

Crops			
Soybean			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off and early-season Phytophthora	0.16 - 0.64	0.004 – 0.015	0.006 – 0.023
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 3,000 soybean seeds/lb</li> <li>For best early-season Phytophthora protection, use the higher specified rate.</li> <li>For a greater spectrum of seedling disease protection of soybean, combine Apron XL LS with azoxystrobin (e.g., Dynasty), fludioxonil (e.g., Maxim 4FS), picarbutrazox (Vayantis), or sedaxane (e.g., Vibrance).</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li><b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.052 lb ai/A/year (0.031 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
Resistance Management:			
Refer to <b>Section 3.1</b> .			

## 7.18 Sunflower

Crops			
Sunflower			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Systemic downy mildew	1.28	0.031	0.031
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 4,500 sunflower seeds/lb</li> </ul>			
USE RESTRICTIONS			
<b>Maximum Annual Application Rate:</b> <b>DO NOT</b> apply more than 0.0025 lb ai/A/year (0.062 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.			
Resistance Management:			
Refer to <b>Section 3.1</b> .			

## 7.19 Turfgrass

<b>Crops</b>			
Turfgrasses (All)			
<b>Target Disease</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed*</b>
Pythium damping-off	0.64 - 1.28	0.015 – 0.031	0.00003 – 0.00006
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>• *based on an average seed weight of 2,200,000 seeds/lb</li> <li>• For use on golf courses, home lawns, reclamation and erosion control projects.</li> <li>• For long-term protection from Pythium in areas when using seed treated with Apron XL LS, make an application of Subdue MAXX® (EPA Reg. No. 100-796) 7-10 days after seeding.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li>• <b>Maximum Annual Application Rate:</b> DO NOT apply more than 0.0161 lb ai/A/year (0.062 lb ai/100 lb seed/year) of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> </ul>			
<b>Resistance Management:</b>			
Refer to <b>Section 3.1</b> .			

## 8.0 SEED CONTAINER LABEL REQUIREMENTS – EXPORT USE

THE FEDERAL SEED ACT REQUIRES THAT CONTAINERS OF SEEDS TREATED WITH APRON XL LS MUST BE LABELED WITH THE FOLLOWING STATEMENTS:

- This seed has been treated with mefenoxam fungicide.
- **DO NOT** use for feed, food, or oil purposes.

THE U.S. ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON CONTAINERS OF SEEDS TREATED FOR EXPORT USE WITH APRON XL LS:

- **TREATED SEED FOR EXPORT ONLY – NOT FOR DOMESTIC SALE OR USE IN THE UNITED STATES.**

**EXPORTER IS RESPONSIBLE FOR VERIFYING SEED CONTAINER LABEL REQUIREMENTS IN COUNTRY OF DESTINATION OR USE. MAXIMUM USE RATES SPECIFIED ON THIS LABEL MUST NOT BE EXCEEDED.**

## 9.0 SEED TREATMENT DIRECTIONS – EXPORT USE

### 9.1 Brassica Head and Stem Vegetable Crop Group 5-16

<b>Crops</b>			
Broccoli	Cauliflower		
Brussels sprouts	Cultivars, varieties, and hybrids of these commodities		
Cabbage			
Cabbage, Chinese (napa)			
<b>Target Disease</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed*</b>
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.00035 – 0.0007
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>• *based on an average seed weight of 100,000 seeds/lb</li> <li>• Verify the actual registered rate in country of destination or use.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li>• Maximum use rates specified on this label must not be exceeded.</li> </ul>			
<b>Resistance Management:</b>			
<ul style="list-style-type: none"> <li>• Refer to <b>Section 3.1</b>.</li> </ul>			



## 9.2 Bulb Vegetables Crop Group 3-07

Crops			
Chive, fresh leaves		Onion, Beltsville bunching	
Chive, Chinese, fresh leaves		Onion, bulb	
Daylily, bulb		Onion, Chinese, bulb	
Elegans hosta		Onion, fresh	
Fritillaria, bulb		Onion, green	
Fritillaria, leaves		Onion, macrostem	
Garlic, bulb		Onion, pearl	
Garlic, great-headed, bulb		Onion, potato, bulb)	
Garlic, Serpent, bulb		Onion, tree, tops	
Kurrat		Onion, Welsh, tops	
Lady's leek		Shallot, bulb	
Leek		Shallot, fresh leaves	
Leek, wild		Cultivars, varieties, and/or hybrids of these	
Lily, bulb			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.00035 – 0.0007
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 100,000 seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
Resistance Management:			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1</b>.</li> </ul>			

## 9.3 Cereal Grains

### 9.3.1 Corn

Crops			
Field Corn			
Sweet Corn			
Popcorn			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.019 – 0.039
Systemic downy mildew	2.2	0.053	0.133
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 1,800 seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
Resistance Management:			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1</b>.</li> </ul>			

### 9.3.2 Millet

<b>Crops</b>			
Millet, Pearl Millet, Proso			
<b>Target Disease</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed*</b>
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.0023 – 0.0047
Systemic downy mildew	2.4	0.058	0.017
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 15,000 seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
<b>Resistance Management:</b>			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1</b>.</li> </ul>			

### 9.3.3 Sorghum

<b>Crops</b>			
Sorghum			
<b>Target Disease</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed*</b>
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.0024 – 0.0048
Systemic downy mildew	2.2	0.053	0.0165
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 14,500 sorghum seeds/lb</li> <li>Only treat high-quality sorghum seed (90% germ and above) with Apron XL LS.</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
<b>Resistance Management:</b>			
Refer to <b>Section 3.1</b> .			

## 9.4 Cucurbit Vegetables Crop Group 9

<b>Crops</b>			
Chayote (fruit)	Muskmelon	Squash, summer ( <i>Cucurbita pepo</i> var. <i>melo</i> )	
Chinese waxgourd (Chinese preserving melon)	(hybrids and/or cultivars of <i>Cucumis melo</i> )	Crookneck squash	
Citron melon	Cantaloupe	Scallop squash	
Cucumber	Casaba	Straightneck squash	
Gherkin	Crenshaw melon	Vegetable marrow	
Gourd, edible ( <i>Lagenaria</i> spp.)	Golden pershaw melon	Zucchini	
Hyotan	Honeydew melon	Squash, winter	
Cucuzza	Honey balls	( <i>Cucurbita maxima</i> , <i>C. moschata</i> )	
Gourd, edible ( <i>Luffa acutangula</i> , <i>L. cylindrical</i> )	Mango melon	Butternut squash	
Hechima	Persian melon	Calabaza	
Chinese okra	Pineapple melon	Hubbard squash	
<i>Momordica</i> spp.	Santa Claus melon	Squash, winter	
Balsam apple	Snake melon	( <i>Cucurbita mixta</i> , <i>C. pepo</i> ):	
Balsam pear	True cantaloupe	Acorn squash	
Bitter melon	Pumpkin	Spaghetti squash	
Chinese cucumber		Watermelon (hybrids and/or varieties of <i>Citrullus lanatus</i> )	
<b>Target Disease</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed*</b>
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.0023 – 0.0047
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 15,000 seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
<b>Resistance Management:</b>			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1</b>.</li> </ul>			

## 9.5 Fruiting Vegetable Crop Group 8-10

Crops			
African eggplant	Groundcherry	Pepper, non-bell	
Bush tomato	Martynia	Roselle	
Cocona	Naranjilla	Scarlet eggplant	
Currant tomato	Okra	Sunberry	
Eggplant	Pea eggplant	Tomatillo	
Garden huckleberry	Pepino	Tomato	
Goji berry	Pepper, bell	Tree Tomato	
Cultivars, varieties and/or hybrids of these			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.00023 – 0.00047
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 150,000 seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
Resistance Management:			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1</b>.</li> </ul>			

## 9.6 Leaf Petiole Vegetable Crop Subgroup 22B

Crops			
Cardoon	Fuki		
Celery	Kohlrabi		
Celery, Chinese	Rhubarb		
Celtuce	Udo		
Fennel, Florence (finocchio)	Zuiki		
Cultivars, varieties, and hybrids of the commodities in crop subgroup 22B			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.000035 – 0.00007
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 1,000,000 seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
Resistance Management:			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1</b>.</li> </ul>			

## 9.7 Leafy Vegetable Crop Group 4-16

Crops			
Amaranth, Chinese	Corn Salad	Mizuna	
Amaranth, Leafy	Cosmos	Mustard greens	
Arugula (Roquette)	Cress, garden	Orach	
Aster, Indian	Cress, upland (yellow rocket, winter cress)	Parsley, fresh leaves	
Blackjack	Dandelion, leaves	Plantain, Buckhorn	
Broccoli, Chinese (gai lon)	Dang-gwi, leaves	Primrose, English	
Broccoli Raab (rapini)	Dillweed	Purslane, Garden	
Cabbage, Abyssinian	Dock (sorrel)	Purslane, Winter	
Cabbage, Chinese (bok choy)	Dol-nam-mul	Radicchio (red chicory)	
Cabbage, seakale	Ebolo	Radish, leaves	
Cat's Whiskers	Endive (escarole)	Rape Greens	
Cham-chwi	Escarole	Rocket, wild	
Cham-na-mul	Fameflower	Shepherd's Purse	
Chervil, fresh leaves	Feather Cockscomb	Spinach	
Chipilin	Good King Henry	Spinach, Malabar	
Chrysanthemum, garland	Hanover Salad	Spinach, New Zealand	
Cilantro, fresh leaves	Huazontle	Spinach, Tanier	
Collards	Jute, leaves	Swiss chard	
	Kale	Turnip Greens	
	Lettuce, Bitter	Violet, Chinese, leaves	
	Lettuce, Head	Cultivars, varieties, and hybrids of these commodities	
	Lettuce, Leaf		
	Maca, leaves		
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.00009 – 0.00018
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 400,000 seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
Resistance Management:			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1</b>.</li> </ul>			

## 9.8 Peas

<b>Crops</b>			
Pea ( <i>Pisum</i> spp.) Dwarf Pea Edible-pod Pea English Pea Field Pea Garden Pea Green Pea Snow Pea Sugar Snap Pea			
<b>Target Diseases</b>	<b>Use Rate</b>		
	<b>fl oz/100 lb of seed</b>	<b>lb ai/100 lb seed</b>	<b>mg ai/seed*</b>
Pythium damping-off and early-season Phytophthora	0.64 - 1.28	0.015 – 0.031	0.056 – 0.112
Downy mildew	1.28	0.031	0.112
<b>Additional Information</b>			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 1,250 seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
<b>USE RESTRICTIONS</b>			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
<b>Resistance Management:</b> Refer to <b>Section 3.1.</b>			

## 9.9 Root and Tuber Vegetables Crop Group 1, except Potato

Crops			
Arracacha		Horseradish	
Arrowroot		Leren	
Artichoke, Chinese		Parsley, turnip-rooted	
Artichoke, Jerusalem		Parsnip	
Beet, garden		Radish	
Beet, sugar		Radish, oriental (daikon)	
Burdock, Edible		Rutabaga	
Canna, Edible (Queensland arrowroot)		Salsify (oyster plant)	
Carrot		Salsify, black	
Cassava, Bitter & Sweet		Salsify, Spanish	
Celeriac (celery root)		Skirret	
Chayote (root)		Sweet Potato	
Chervil, turnip-rooted		Tanier (cocoyam)	
Chicory		Turmeric	
Chufa		Turnip	
Dasheen (Taro)		Yam Bean (jicama, manioc pea)	
Ginger		Yam, True	
Ginseng			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Pythium damping-off	0.32 - 0.64	0.008 – 0.015	0.00012 – 0.00023
Additional Information			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 300,000 seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> <li><b>DO NOT</b> Apply Apron XL LS to potatoes.</li> </ul>			
Resistance Management:			
<ul style="list-style-type: none"> <li>Refer to <b>Section 3.1</b>.</li> </ul>			

## 9.10 Sunflower

Crops			
Sunflower			
Target Disease	Use Rate		
	fl oz/100 lb of seed	lb ai/100 lb seed	mg ai/seed*
Systemic downy mildew	3.75 - 4.45	0.090 – 0.107	0.091 – 0.108
Additional Use Directions			
<ul style="list-style-type: none"> <li>*based on an average seed weight of 4,500 sunflower seeds/lb</li> <li>Verify the actual registered rate in country of destination or use.</li> </ul>			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>Maximum use rates specified on this label must not be exceeded.</li> </ul>			
Resistance Management:			
Refer to <b>Section 3.1</b> .			

## 10.0 STORAGE AND DISPOSAL

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

#### Pesticide Storage

Store in the original container and only in a cool, dry, secure place.

#### Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. Open dumping is prohibited.

#### Container Handling [(less than or equal to 5 gallons)]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

#### Container Handling [(greater than 5 gallons)]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

#### Container Handling [(greater than 5 gallons)]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions and restrictions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**



## 11.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent consistent with applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent consistent with applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

---

Apron XL®, Dynasty®, Maxim®, Ridomil Gold®, Subdue MAXX®, Vayantis®, Vibrance®, the ALLIANCE FRAME, the SYNGENTA Logo, and the PURPOSE ICON are trademarks of a Syngenta Group Company

Viton™ is a trademark of The Chemours Company FC, LLC

©20\_\_ Syngenta

For non-emergency (e.g., current product information) call  
Syngenta Crop Protection at 1-866-796-4368.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

APRON XL LS 799 MAS 0820 AMEND-C 0921-CL - jeb - 05/23/2023  
000100-00799.20210907C.APRON-XL-LS-AMEND-SEPT2021-CL