

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number NOTIFICATION
---	---	--	--

Application for Pesticide - Section I

1. Company/Product Number 100-801	2. EPA Product Manager Mary Waller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Ridomil Gold® Bravo Liquid NCP 801C-M2A 0298	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Novartis Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated February 13, 1998 <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
---	--

NOTIFICATION
 February 13, 1998
MAY 14 1998


Explanation: Use additional page(s) if necessary. (For Section I and Section II.)

PR Notices 95-2 and 97-6: Change in package size; Change of company name to Novartis; and use of "other ingredients" per PRN 97-6; ISK notified Novartis that they changed the name from Bravo 720 to Bravo Weatherstick (alternate brand name); New EC formulation approved 2/13/98 for Ridomil Gold® EC (Mefenoxam EC; 100-801). Submission of Final printed Labeling in response to stamped label sent to Novartis by Ms. Mary Waller, PM 21, on February 13, 1998.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
*Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Unit Packaging wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Lee Hubbard	Title Senior Regulatory Manager	Telephone No. (Include Area Code) 910- 632-7034
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Senior Regulatory Manager	
4. Typed Name Lee Hubbard	5. Date April 28, 1998	

2079

NOTIFICATION

MAY 14 1998

PULL HERE TO OPEN ▶

Ridomil Gold®



Twin-Pak™

FUNGICIDE

For control of certain diseases in potatoes

See additional precautionary statements and directions for use inside booklet.

Following label instructions, apply the entire contents of this container to the target crop once the container is opened.

EPA Est. 34704-MS-2

NCP 801C-M2A 0298

Total **272 fl. oz.**
U.S. Standard Measure

Ridomil Gold EC

Active Ingredients:	
(R)-2-[(2,6-dimethylphenyl)-methoxyacetyl-amino]-propionic acid methyl ester	47.6%
Related Compounds	1.4%
Other Ingredients	51.0%
Total:	100.0%
EPA Reg. No. 100-801	

KEEP OUT OF REACH OF CHILDREN.

CAUTION

32 fl. oz. U.S. Standard Measure

Bravo Weatherstick

Active Ingredient:	
Chlorothalonil (tetrachloro-isophthalonitrile)	54.0%
Inert Ingredients:	46.0%
Total:	100.0%
Contains 6.0 lbs. chlorothalonil per gallon (720 grams per liter)	
EPA Reg. No. 50534-188	

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

240 fl. oz. U.S. Standard Measure

 **NOVARTIS**

Ridomil Gold®/Bravo® Liquid

309

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Novartis Crop Protection, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Novartis warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. **Novartis makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Novartis or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product.** Novartis and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Novartis.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Ridomil Gold/Bravo Liquid

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

GENERAL INFORMATION

Ridomil Gold/Bravo Liquid is a fungicide for the control of early blight (*Alternaria*), late blight (*Phytophthora*), pink rot, and pythium leak in potatoes. Following label instructions, apply the entire contents of this container to the target crop once the container is opened. Each container treats 10 acres of potatoes.

Note: Ridomil Gold active ingredient is a systemic fungicide having a specific mode of action and could be subject to development of insensitive strains of fungi. Development of insensitivity cannot be predicted. Therefore, Novartis cannot assume liability for crop damage resulting from insensitive strains of fungi. If treatment is not effective following the use of Ridomil Gold/Bravo Liquid as recommended, an insensitive strain of fungi may be present. If the treatment is ineffective due to the presence of a Ridomil Gold insensitive strain of fungi, neither Ridomil Gold/Bravo Liquid nor any other fungicide with similar action will effectively control that disease. Consideration should then be given to the prompt use of other types of suitable fungicides. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance on your particular crop and disease control situation.

THIS LABEL IS FOR FIELD USE ONLY AND IS NOT INTENDED FOR USE ON TRANSPLANT TRAYS, GREENHOUSES, LATH HOUSES, FLOAT HOUSES, HYDROPONIC PRODUCTION, OR IN BEDDING PLANT STRUCTURES.

Ridomil Gold®/Bravo® Liquid

4029

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

Mixing Instructions

Note: Slowly invert container several times to assure uniform mixture.

Prepare no more spray mixture than is required for the immediate operation. Use the entire contents of this container once the container is opened. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

Ridomil Gold/Bravo Liquid Alone: Add 1/2-2/3 of the required amount of water to the spray or mixing tank. With the agitator running, add the Ridomil Gold/Bravo Liquid to the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after the Ridomil Gold/Bravo Liquid has completely dispersed into the water. Maintain continuous agitation until all of the mixture has been sprayed.

Ridomil Gold/Bravo Liquid + Tank Mixtures: Consult your dealer or Novartis sales representative for information concerning compatibility with other tank mix partners. **Note:** When using Ridomil Gold/Bravo Liquid in mixture, any product in water-soluble packaging should be added to the tank prior to adding Ridomil Gold/Bravo Liquid. Allow water-soluble packaging to completely dissolve and the mixture product to completely disperse before adding Ridomil Gold/Bravo Liquid to the tank.

Add 1/2-2/3 of the required amount of water to the spray or mixing tank. Again, if tank mixing with any product packaged in water-soluble packets, add this product first with the agitator running. Allow the material to completely dissolve and disperse in the mix water. Continue agitation while adding all of the contents of both chambers of the Ridomil Gold/Bravo Liquid container at once. In general, additional tank mix partners should be added in the following order: wettable powders, dry flowable formulations, liquid flowable formulations, and emulsifiable concentrations. Always allow each tank mix partner to become fully dispersed before adding the next product. Maintain agitation while adding the rest of the water, while adding the other tank mix partners and throughout the mixing and spraying operation.

If using Ridomil Gold/Bravo Liquid in mixture with other products, observe all directions for use crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate should be exceeded and the most restrictive label precautions and limitations should be followed. Ridomil Gold/Bravo Liquid should not be mixed with any product which prohibits such mixing. Tank mixtures or other applications of products are permitted only in those states in which the products are registered.

Important: Ridomil Gold/Bravo Liquid is not recommended for use in combination with Dipel® 4L, Triton AG-98, or Triton B-1956, as phytotoxicity may result.

Application Through Irrigation Systems

Ridomil Gold/Bravo Liquid, alone or in combination with other pesticides which are registered for application through irrigation systems, may be applied through irrigation systems.


Apply this product only through center pivot, solid set, hand move, or moving wheel. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended.

Application Instructions

Ridomil Gold/Bravo Liquid must be applied on the schedule specified in the specific crop use recommendations, not according to the irrigation schedule.



5049

Ridomil Gold®/Bravo® Liquid

The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler irrigation equipment. Users must check with state and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler or drip irrigation equipment.

Note: Do not inject Ridomil Gold/Bravo Liquid at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 15 parts water to 1 part Ridomil Gold/Bravo Liquid in the mix tank. Ridomil Gold/Bravo Liquid is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used, but should be replaced once a year. Do not use Viton, Buna-N, Neoprene, or PVC seals.

Center Pivot Irrigation Equipment

Note: (1) Use only with drive systems which provide uniform distribution. (2) Do not use end guns when chemigating Ridomil Gold/Bravo Liquid through center pivot irrigation systems because of nonuniform application. (3) Plug the first nozzle closest to well head to protect water source.

1. Determine the size of the area to be treated.
2. Determine the time required to apply 1/4 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. Run the system at 80-95% of the manufacturer's rated capacity.
3. Using water, determine the injection pump output when operated at normal line pressure.
4. Determine the amount of Ridomil Gold/Bravo Liquid required to treat the area covered by the irrigation system.
5. Add the required amount of Ridomil Gold/Bravo Liquid and sufficient water to meet the injection time requirements to the solution tank.
6. Make sure the system is fully charged with water before starting injection of the Ridomil Gold/Bravo Liquid solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
7. Maintain constant solution tank agitation during the injection period.
8. Continue to operate the system until the Ridomil Gold/Bravo Liquid solution has cleared the sprinkler head.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment

1. Determine the acreage covered by the sprinklers.
2. Fill injector solution tank with plain water and calibrate the flow rate of the system to deliver the contents of the tank over a 20- to 30-minute time interval.
3. Determine the amount of Ridomil Gold/Bravo Liquid required to treat the area covered by the irrigation system.
4. Add the required amount of Ridomil Gold/Bravo Liquid into the same quantity of water used to calibrate the injection period.
5. Operate the system at the same pressure and time interval established during the calibration.
6. Inject Ridomil Gold/Bravo Liquid at the beginning of the irrigation cycle in 1/2-1 inch of water or as a separate application to maximize the effectiveness of the fungicide. Do not apply in excess of 1 inch of water.
7. Stop injection equipment after treatment is completed. Continue to operate the system until the Ridomil Gold/Bravo Liquid solution has cleared the last sprinkler head.

POTATOES


Ridomil Gold/Bravo Liquid, when used as a foliar application in a preventive disease control program, will provide effective control of early blight, caused by *Alternaria* spp.; late blight, caused by *Phytophthora infestans*; pythium leak caused by *Pythium* spp.; and pink rot caused by *Phytophthora erythroseptica*. Make applications of Ridomil Gold/Bravo Liquid in sufficient water to obtain thorough coverage for ground applications and in a minimum of 10 gals. of water for aerial applications.

It is recommended that this product be used within an Integrated Pest Management Program. Also, vine-kill should occur 14 days before harvest.

Late Blight and Early Blight: Use the entire contents of this container to treat 10 acres of potatoes. Begin preventive applications early in the season when conditions are favorable for disease (before infection), but no later than when the plant foliage meets within the row uniformly across the field. In states other than FL, make up to 3 applications of Ridomil Gold/Bravo Liquid at 14-day intervals. In FL only, make up to 4 applications at 14-day intervals. The labeled rate of a protectant fungicide should be applied between Ridomil Gold/Bravo Liquid applications. Following the 3-4 Ridomil Gold/Bravo Liquid applications, apply the labeled rate of a protectant fungicide on a weekly schedule through the remainder of the season.

Storage Rots: Pythium Leak (caused by *Pythium* spp.) and Pink Rot (caused by *Phytophthora erythroseptica*): For effective control of these storage rots, Ridomil Gold/Bravo Liquid should be used in conjunction with other management practices such as crop rotation. Use the entire contents of this container to treat 10 acres of potatoes. Make the first application at flowering and another one 14 days later. If the field has a history of storage rot problems; make a third application 14 days after the second application.

Note: If applications of Ridomil Gold/Bravo Liquid are being made for late blight control that correspond to the timing of application for storage rot control, additional applications for storage rot are not needed.



6 of 9

Ridomil Gold®/Bravo® Liquid

Notes: (1) To avoid possible illegal residues, do not apply within 14 days of harvest. (2) In states other than FL, you may make up to 3 applications of Ridomil Gold/Bravo Liquid per crop. In FL only, you may make up to 4 applications per crop. (3) If conditions for late blight development are still favorable after making 3 applications of Ridomil Gold/Bravo Liquid, use other fungicides registered for late blight control in potatoes.

Rotation (Plant Back) Restrictions

Do not plant any crop which is not registered for use with the Ridomil Gold active ingredient in soil treated with this active ingredient for a period of 12 months. See the following list.

Rotation Crop	Planting Time From Last Application of Ridomil Gold/Bravo Liquid
Alfalfa (including birdsfoot trefoil), Almonds, Apples*, Asparagus, Avocados	0 days
Blueberries, Broccoli	
Cabbage, Cauliflower, Chinese Broccoli (gai lan, white flowering broccoli), Chinese Cabbage (tight-heading varieties only), Citrus, Cotton, Cranberries, Cucurbit Vegetables	
Deciduous Fruits and Nuts*	
Eggplant	
Garlic, Ginseng, Grapes	
Grass Grown for Seed	
Hops	
Leafy Vegetables (except Brassica vegetables), Legume Vegetables (beans and peas – succulent and dried)	
Onions (Dry Bulb, Green, and Seed)	
Papaya, Peanuts, Peppers, Pineapples, Potatoes	
Raspberries, Root and Tuber Vegetables	
Soybeans, Spinach, Stone Fruits, Strawberries, Sugar Beets	
Tobacco, Tomatoes	
Walnuts	
Cereal Grains (except corn)	40 days
Corn	9 months
Crops Not For Food or Feed	0 days
All Other Crops For Food or Feed	12 months

*These crops may be planted immediately following last application of Ridomil Gold/Bravo Liquid, provided they will not bear harvestable fruit within 12 months.

STORAGE AND DISPOSAL

Storage Instructions

Store in the original container. Using label instructions, apply the entire contents of this container to the target crop once the container is opened. Store in a cool place. Protect from excessive heat.


Pesticide Disposal

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Disposal

Do not reuse empty container. Triple rinse container (or equivalent) and then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by burning. Stay out of smoke from burning containers.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.



7 of 9

PRECAUTIONARY STATEMENTS

Ridomil Gold EC
Hazards to Humans and Domestic Animals
CAUTION
 Causes moderate eye irritation. Do not get in eyes or on clothing. Harmful if swallowed or absorbed through skin. Avoid contact with skin.
Statement of Practical Treatment
If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.
If on skin: Wash with plenty of soap and water. Get medical attention.
If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
Note to Physician: If ingested, induce emesis or lavage stomach. Treat symptomatically.

Bravo Weatherstick
Hazards to Humans and Domestic Animals
WARNING/AVISO
 Causes eye irritation. May cause skin irritation. May be a potential skin sensitizer. Do not get in eyes. Avoid prolonged contact with skin. Avoid breathing spray mist. Do not take internally.
Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation, and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.
Statement of Practical Treatment
If in eyes: Immediately flush eyes with plenty of water and continue for 15 minutes. Seek medical attention for eyes immediately.
If on skin: Wash exposed areas of skin with soap and warm water after handling or using.
If swallowed: Never give anything by mouth to an unconscious person.
Note to Physician: Persons having a temporary allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.


User Safety Recommendations
Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Ridomil Gold EC

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Apply only as specified on this label. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.



8049

Ridomil Gold®/Bravo® Liquid

Bravo Weatherstick

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Do not apply when weather conditions favor drift from treated areas. Apply only to areas specified on label.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat, open flame, or hot surfaces.

Ridomil Gold® and Twin-Pak™ trademarks of Novartis

Bravo® trademark of ISK Biosciences Corporation


Dipel® trademark of Abbott Laboratories

Triton AG-98 and Triton B-1956 trademarks of Rohm and Haas Company

©1998 Novartis

Novartis Crop Protection, Inc.
Greensboro, North Carolina 27419

NCP 801C-M2A 0298



9079

Ridomil Gold



FUNGICIDE

For control of certain diseases in potatoes

Using label instructions, apply the entire contents of this container to the target crop once the container is opened.

EPA Est. 34704-MS-2

©1998 Novartis
Novartis Crop Protection, Inc.
Greensboro, NC 27419
NCP 801C-M2A 0298

Total **272 fl. oz.**
U.S. Standard Measure

Ridomil Gold EC

Active Ingredients:	
(R)-2-[(2,6-dimethylphenyl)-methoxyacetylamino]-propionic acid methyl ester . . .	47.6%
Related Compounds	1.4%
Other Ingredients: 51.0%	
Total:	100.0%

EPA Reg. No. 100-801
Ridomil Gold® and Twin-Pak™ trademarks of Novartis

32 fl. oz.
U.S. Standard Measure

Bravo® Weatherstick

Active Ingredient:	
Chlorothalonil (tetrachloroisophthalonitrile)	54.0%
Inert Ingredients:	46.0%
Total:	100.0%

Contains 6.0 lbs. chlorothalonil per gallon (720 grams per liter)

EPA Reg. No. 50534-188
Bravo® trademark of ISK Biosciences Corporation

240 fl. oz.
U.S. Standard Measure

See directions for use in attached booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

