

100-804

4-10-2002

1/3

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Larry Zang
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

APR 10 2002

Subject: Ridomil Gold Copper Fungicide (Supplemental label)
EPA Reg. No. 100-804
Submission dated April 4, 2002 revising the signal word.

Dear Mr. Zang:

The revised product labeling referred to above, submitted in connection with the registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

A stamped "supplemental label" is enclosed for your records. Submit one copy of your final printed label and incorporate the supplemental labeling into the main label within 18 months. If you have questions about this label review, please contact Dennis McNeilly at (703) 308-6742 or electronically at mneilly.dennis@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Mary L. Waller".

Mary L. Waller
Product Manager 21
Fungicide Branch
Registration Division (7505C)

Enclosure

PRODUCT INFORMATION



Syngenta Crop Protection, Inc.
Greensboro, NC 27409
www.syngenta-us.com


SUPPLEMENTAL LABEL FOR RIDOMIL GOLD® COPPER

FUNGICIDE

EPA Reg. No. 100-804

Contains 65% active ingredient per pound

To aid in control of *Phytophthora* blight caused by *Phytophthora palmivora*.



KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

SCP 804B-S1A 0402

DIRECTIONS FOR APPLICATIONS

Papaya, Black Sapote, Star Apple, Canistel, Mamey Sapote, Mango, Sapodilla

Use Ridomil Gold Copper to aid in the control of Phytophthora blight caused by *Phytophthora palmivora* in nurseries, new plantings in the field and on bearing plants.

Apply 2-3 lbs. (1.3 to 1.95 lb. a.i.) of Ridomil Gold Copper per acre as a trunk or foliar spray. Direct the spray to the trunk of the plant or to the fruit column on the trunk. Apply until the point of run-off. When applying to a fruit column use a minimum of 100 gallons of water per acre. Make up to 4 applications at approximately 21 day intervals. Do not apply within 1 day of harvest.

Do not apply Ridomil Gold Copper if Ridomil Gold[®] EC is used as a soil drench.

Ridomil Gold[®] Copper and the Syngenta logo are trademarks of a Syngenta Group Company

© 2002 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 804B-S1A 0402

Z/SUPPLEMENTAL LABELS/RID GD COP 804B-S1A 0402 – ccg – 4/3/02