	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number NOTIFICATION
	Application for Pesticide - Section I		

1. Company/Product Number 100-806	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Accu-Pak Action Herbicide	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---


NOTIFICATION
APR 13 2001

Explanation: Use additional page(s) if necessary. (For Section I and Section II.). This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. The following changes are being made via this notification: 1) Company name and address have been updated to reflect Syngenta Crop Protection, Inc. 2) The Conditions of Sale and Warranty statement has been changed to reflect the name change. Because Syngenta has been formed by the merger of Novartis Crop Protection, Inc. and Zeneca Ag Products, we have chosen to use the former Zeneca warranty statement as the Syngenta warranty statement. No other changes occur in the statement other than the name change. 3) The copyright date reflects Syngenta. 4) Trademark statements have been updated to reflect Syngenta for those products for which Syngenta holds the trademark. 5) The Internet address has been changed to reflect Syngenta. 6) Other places in the label which referring to the company name have been updated.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No			
*Certification must be submitted		If "Yes" Unit Packaging wgt. No. per Container	If "Yes" Unit Packaging wgt. No. per container	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Nan S. Padgett	Title Label Group Leader	Telephone No. (Include Area Code) 336-632-7567			
Certification				6. Date Application Received (Stamped)	
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.				
2. Signature 		3. Title Label Group Leader			
4. Typed Name Nan S. Padgett		5. Date April 2, 2001			

NOTIFICATION

APR 13 2001

PULL HERE TO OPEN ►

Accu-Pak®

Action®

HERBICIDE

For postemergence control of velvetleaf and certain other broadleaf weeds in soybeans

**KEEP OUT OF REACH OF CHILDREN.
CAUTION**

See additional precautionary statements and directions for use inside booklet.

<p>Active Ingredient: Fluthiacet-methyl (CAS No. 117337-19-6) 4.75%</p> <p>Other Ingredients: 94.25%</p> <p>Total: 100.00%</p>	<p>EPA Reg. No. 100-806 EPA Est. SCP 806A-L1A 0201</p>
--	---

Action is a wettable powder packaged in a water-soluble bag.

This outer protective bag contains Action in 3 oz. water-soluble packets. Entire inner packets and contents dissolve in water. After opening outer bag, immediately dump the required number of unopened packets into a sprayer or mix tank partially filled with water. Do not excessively handle the inner soluble packets or expose them to moisture, since this may cause rupturing.

5 x 3 OUNCE
Water-Soluble Packets

15 OUNCES
Total Net Weight Per Bag

syngenta

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and the restricted-entry interval.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

GENERAL INFORMATION

Action is a selective postemergence herbicide for control of velvetleaf and other broadleaf weeds in soybeans. Action must be applied after emergence of the crop and weeds. There is a wide application window for use in soybeans. This application window ranges from the first trifoliolate to the full flowering stage. Refer to Table 2 for a list of the weeds controlled.

Action will control velvetleaf up to 36 inches tall. Action will also control or partially control certain other annual broadleaf weeds. Partial control means significant activity but not always at a level generally considered acceptable for commercial weed control. When tank mixed, Action enhances the performance of other herbicides. Action does not control grasses. If grasses are present at the time of application, Action should be mixed with an appropriate postemergence herbicide registered for grass control in soybeans.

The amount of Action to apply and the degree of weed control resulting from Action application depends on the weed species present, stage of growth of the weeds, environmental conditions, and growing conditions. Weeds under stress because of lack of moisture, low soil fertility, or mechanical or chemical injury, may not be controlled as well as actively-growing weeds.

To be effective, Action must contact the weed foliage. Weed death or inhibition can be expected quickly – normally within 48 hours after application. Thorough coverage with the spray solution gives the most effective control. A large crop and/or weed canopy as well as a dense crop and weed canopy can prevent the spray from penetrating the canopy and reaching smaller weeds resulting in reduced control.

Soybeans are tolerant of Action when applied according to label directions. Some bronzing, crinkling, or spotting of soybean leaves may occur. Soybeans rapidly outgrow these effects and develop normally with no reduction in yield.

Note: Cotton is very sensitive to Action. To avoid cotton defoliation, do not allow Action to contact this crop.

Grazing/Feeding

Do not graze or feed treated soybean forage or hay to livestock.

MIXING INSTRUCTIONS

Spray Additives

An adjuvant approved for use on growing crops is required with Action for maximum consistent performance.

1. Adjuvants for Action Alone

Use a spray adjuvant from one of these classes for optimum performance:

Nonionic surfactant (NIS) with a minimum of 80% of the constituents effective as spray adjuvant at the rate of 1 qt./100 gals. of spray volume (concentration of 0.25%).

Crop oil concentrate (COC) petroleum- or vegetable-based containing not less than 12% emulsifier at 1-2 pts./A. The concentration should not exceed 2.5% volume/volume. COC is recommended under conditions of dry soil and low relative humidity.

Silicone-based surfactant at a rate of 1 qt./100 gals. or a spray volume concentration of 0.25% or as specified on the adjuvant label.

In addition to an adjuvant, urea ammonium nitrate (UAN) at 1-2 qts./A or spray grade ammonium sulfate (AMS) at 1.5-2.5 lbs./A may also be added to the spray solution. Do not use liquid fertilizer as the total carrier solution.

2. Adjuvants for Action in Tank Mixtures With Other Herbicides

When tank mixing with other herbicides, use the adjuvant recommended for use with the tank mix partner. Follow all restrictions and precautions on the tank mix partner's label.

Mixing Instructions

1. Thoroughly clean the spray equipment before using Action.
2. Maintain maximum agitation throughout the spraying operation.
3. Water-soluble packets must always be the first material put into the spray tank after water. Water-soluble packets must be completely dissolved and dispersed in clean water before any other tank mix partners, including micronutrients or other fertilizers or adjuvants are added to the spray solution. Boron, especially in the form of a micronutrient additive, such as Solubor[®], etc. may prevent water-soluble packets from dissolving.
4. Flush the spray equipment thoroughly after each use and apply rinsate to a previously treated area.

Mixing Action Alone

1. Add 1/4-1/2 of the required amount of clean water to the spray or mixing tank.
2. With the agitator running, drop the required number of **unopened** water-soluble packets of Action into the spray tank. Continue agitation allowing the packets to completely dissolve and the contents of the packets to fully and uniformly disperse.
3. Add the spray adjuvant and continue agitation while adding the rest of the water.
4. Maintain agitation until all of the mixture has been applied.

Action in Tank Mixtures

Action is compatible with most commonly used insecticides, fungicides, and spray adjuvants. **BEFORE MIXING ACTION WITH OTHER REGISTERED PRODUCTS FOR ANY USE ON THIS LABEL, READ THE LABEL OF THE TANK MIX PARTNER TO BE CERTAIN IT IS LABELED FOR THE USE ON THE PARTICULAR CROP AND THAT USE PATTERNS ARE COMPATIBLE WITH THOSE OF ACTION.** When using Action in a tank mixture with other pesticides, observe the most restrictive label limitations and precautions of the products being used.

When Action is mixed with acifluorfen or lactofen-containing products (such as Blazer[®], Galaxy[®], Cobra[®], Storm[®]), velvetleaf control will be improved. However, these tank mixes are not as effective in controlling velvetleaf as the use of Action alone.

1. Add 1/4-1/2 of the required amount of clean water to the spray or mixing tank.
2. Start the agitator running before adding any tank mix partners. With the agitator running drop the required number of **unopened** soluble packets of Action and any of the other products packaged in water-soluble packets into the tank all at once. Continue agitation in the spray tank and allow the packets to completely dissolve and the contents of the packets to fully and uniformly disperse.
3. While maintaining agitation, continue filling the spray tank. When the tank is 3/4 full, add any tank mix partners and allow them to completely and uniformly disperse.
4. Add any other liquid tank mix partners and allow them to completely and uniformly disperse.
5. Add the proper amount of spray adjuvant and continue agitation while adding the remaining water.
6. Complete filling the tank with clean water and maintain sufficient agitation at all times to insure surface action and a uniform mixture.
7. To avoid product degradation, apply Action spray mixture within 48 hours of mixing and do not leave spray in the tank without continuous agitation.
8. After use, thoroughly clean the sprayer according to this label and the tank mix partner's label.

APPLICATION PROCEDURES AND PRECAUTIONS

Chemigation

Do not apply this product through any type of irrigation equipment.

Aerial Application

Do not apply this product aerially.

Ground Application

Mix the product in a minimum of 15 gals. of water and apply at 20-40 psi pressure at the nozzles. If a dense crop and/or weed canopy is present, use up to 40 gals. of water per acre and 70 psi nozzle pressure. **Note:** When using higher nozzle pressures, use extreme caution to avoid spray drift to nearby crops.

Use a pump with enough capacity to maintain a rippling or rolling action in the spray tank. For a uniform spray mixture, agitation during mixing and application is required.

How Much to Use

Table 1. Action Use Rate

	Number of Packets Required			
	Tank Mixed with Roundup® Original or Roundup Ultra	Tank Mixed with Another Broadleaf Herbicide	Alone or With a Grass Control Herbicide	
			Standard Rate	Higher Rate for Difficult to Control Weeds*
	- oz./Acre -			
Acres	1.0	1.2	1.5	2.25
1.3	-	-	-	1
2	-	-	1	-
2.5	-	1	-	-
3	1	-	-	-
4	-	-	2	3
5	-	2	-	-
6	2	-	3	-
8	-	-	4	6
10	-	4	5	-
12	3	-	6	9
20	-	8	10	15
30	10	12	15	-
50	-	20	25	-
60	20	24	30	45
80	-	32	40	60
100	-	40	50	75

*See Table 2 for list of weeds.

WHEN TO APPLY

Weed Stage of Growth

Apply Action after weeds have emerged and are actively growing, but before the weeds have reached the maximum height listed in Table 2. Application after weeds have reached the listed maximum height could result in commercially unacceptable weed control.

Avoid applying Action if weeds cannot be evenly covered with the spray or when spray drift is possible. To reduce spray drift, do not apply if wind speed is 10 mph or greater. Avoid overlapping the spray pattern since this may increase chances for crop injury.

Soybean Stage of Growth

Action may be applied to soybeans from the first trifoliolate through the full flowering stage of development. The latest application date should be no later than 60 days before harvest.

Cultivation

Do not cultivate within 2 days before or 2 days after applying this product.

Application Precautions

1. Do not apply more than 3 oz./A per cropping season.
2. Do not apply if soybeans are under severe stress due to drought, cold weather, hail, flooding, water-logged or compacted soil, disease, insect damage, nutrient deficiency (especially low nitrogen levels), or other causes.
3. Action can be applied in tank mixtures to weeds taller than the maximum heights listed in Table 2 provided the application follows all timing and rate precautions on the tank mix partner's label.
4. Application to weeds that are under severe stress due to drought or to weeds that are taller than the optimum heights listed in Table 2 may result in reduced weed control.
5. Do not irrigate within 4 hours of application of Action. Rainfall or irrigation may wash Action off of the weeds during this period and reduce performance.
6. Observe all precautions and limitations on the label of each product used in tank mixture with Action.

Spray Drift Precautions

Do not allow spray from ground equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including:

1. Do not spray if winds are gusty or if speeds are or become 10 mph or greater.
2. Use extreme caution when conditions favor drift (high temperatures, low relative humidity).
3. Do not apply during a temperature inversion. If an inversion is suspected, consult the local weather service before applying Action.
4. These practices can further reduce drift:
 - a. Using nozzles that provide a uniform droplet size. Use spray nozzles that provide medium-coarse droplets (250-400 microns VMD). Nozzles that produce extremely small droplets are more likely to cause spray drift.
 - b. Apply as close to target plants as practical while maintaining a good spray pattern for spray coverage.

Cleaning Spray Equipment

The sprayer must be cleaned before and after use of this product. Failure to do so may result in unsatisfactory results with Action or injury to other crops sprayed with the equipment. Refer to the label of the product used previously or tank mixed with Action for additional cleaning instructions.

When Action has been used alone, fill the sprayer with clean water, add a commercial sprayer cleaner, circulate the mixture through the system, and spray approximately 1/2 through the hose, boom, and nozzles on an area which has already been treated. Drain the remaining solution, rinse the entire system with clean water, and spray the rinsate on an area which has already been treated.

WEEDS CONTROLLED/APPLICATION RATES**Action Alone**

At the rate and timings listed, Action controls the weeds listed in Table 2 when the product is applied alone after emergence of soybeans and weeds. Action controls emerged weeds listed; it does not provide residual control of weeds that emerge later. Weeds larger than the size in Table 2 may only be partially controlled.

Table 2. Application Rate for Action Applied Alone

Weed Species	1 Packet/2 Acres (1.5 oz./A)		3 Packets/4 Acres (2.25 oz./A)	
	Leaf Stage*	Maximum Height (inches)	Leaf Stage*	Maximum Height (inches)
Anoda, spurred (<i>Anoda cristata</i>)	2	2	3	4
Burcucumber (<i>Sicyos angulatus</i>)	2	2	2	2
Jimsonweed (<i>Datura stramonium</i>)	2**	2**	4	2
Nightshade, black (<i>Solanum nigrum</i>)	2**	2**	2**	2**
Nightshade, eastern black (<i>S. ptycanthum</i>)	2**	2**	2**	2**
Lambsquarters, common (<i>Chenopodium album</i>)	4**	2**	4-6	2
Pigweed, redroot (<i>Amaranthus retroflexus</i>)	4**	2**	4-6	4
Pigweed, smooth (<i>A. hybridus</i>)	2**	2**	2-4	4
Velvetleaf (<i>Abutilon theophrasti</i>)	-	36	-	36
Waterhemp, common (<i>Amaranthus rudis</i>)	-	-	2	2
Waterhemp, tall (<i>A. tuberculatus</i>)	-	-	2	2

*Count individual leaves except the cotyledons.

**Partial control or suppression.

Action in Tank Mixtures With Other Postemergence Herbicides for Improved Broadleaf Weed Control

Action may be applied postemergence at 1 packet/3 acres (1 oz./A) with Roundup Original or Roundup Ultra or at 1 packet/2.5 acres (1.2 oz./A) with other labeled postemergence broadleaf herbicides as a tank mixture. Action enhances the performance of these herbicides by increasing the speed of activity and/or level of control of the weeds listed in Table 2. When mixed with any of the tank mix partners listed in Table 3, Action also controls velvetleaf up to 36 inches tall.

Refer to Table 3 for a list of recommended tank mix partners and follow all recommendations, restrictions, and precautions for each product in the tank mixture. Refer to Table 1 for the amount of Action to use. If a second application is needed, do not apply more than a total of 3 oz. of Action per acre per season.

Table 3. Action – Recommended Tank Mix Partners – Soybeans

Basagran®	Flexstar® HL	Raptor®	Roundup Ultra
Blazer	Galaxy	Reflex®	Scepter® O.T.®
Classic®	Manifest®	Reliance™ STS® SP	Status®
Cobra	Pinnacle®	Result®	Storm
Concert® SP	Pursuit®	Roundup Original	

Action in Tank Mixtures for Grass Weed Control

Action does not provide grass control, but it can be tank mixed at 1 packet/2 acres (1.5 oz./A) to 1 packet/1.3 acres (2.25 oz./A) with any postemergence grass herbicide registered for use in soybeans to provide control of the weeds listed in Table 2 and the grasses listed on the tank mix partner's label.

CROP FAILURE

If soybeans treated with Action are lost due to a natural catastrophe such as hail or frost, soybeans can be replanted immediately, provided this is not restricted on the label of a product used previously or by a product applied in a tank mixture with Action.

For control of weeds in the replanted crop, Action may be applied postemergence a second time but the total amount of Action applied during the cropping season must not exceed 3 oz./A.

ROTATIONAL CROPS

If soybean crop failure occurs, to avoid illegal residues replant soybeans only. All other crops may be planted at normal rotation intervals following application of Action.

Note: For rotational crop restrictions when Action is used in tank mixtures or sequentially with other products, refer to the rotation intervals on the other product label for possible additional restrictions.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in a cool, dry place. Do not store this product under wet conditions. Handle outer bag carefully to avoid breakage of inner water-soluble packets.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Do not reuse outer container. Dispose of empty container in a sanitary landfill, or incinerate, or if allowed by state and local authorities, dispose of by burning. Stay out of smoke from burning container.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

First Aid

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

Note to Physician: If ingested induce emesis and lavage stomach. Treat symptomatically.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

Accu-Pak® and Action® trademarks of Syngenta

Basagran®, Blazer®, Galaxy®, Manifest®, Result®, and Storm® trademarks of BASF Corporation

Classic®, Concert®, Pinnacle®, and Reliance™ STS® trademarks of E.I. DuPont de Nemours and Company, Inc.

Cobra® trademark of Valent Corporation

Flexstar® and Reflex® trademarks of Zeneca Ag Products

Pursuit®, Raptor®, Scepter® O.T.®, and Status® trademarks of American Cyanamid Company

Roundup® trademark of Monsanto Company

Solubor® trademark of United States Borax and Chemical Corporation

©2001 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 806A-L1A 0201



Action®

HERBICIDE

For postemergence control of
velvetleaf and certain other
broadleaf weeds in soybeans

Active Ingredient: Fluthiacet-methyl (CAS No. 117337-19-6) ...	4.75%
Other Ingredients:	54.25%
Total:	100.00%

Action is a wettable powder packaged
in a water-soluble bag.

See directions for use in attached
booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance
with its labeling and with the Worker
Protection Standard, 40 CFR part 170.
Refer to supplemental labeling under
"Agricultural Use Requirements" in
the Directions for Use section for
information about this standard.

EPA Reg. No. 100-806
EPA Est.

5 x 3 OUNCE
Water-Soluble Packets
15 OUNCES
Total Net Weight Per Bag

KEEP OUT OF REACH OF CHILDREN. CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals
Harmful if absorbed through skin. Avoid contact with
skin, eyes, or clothing. Wash thoroughly with soap
and water after handling. Remove contaminated
clothing and wash before reuse.

First Aid

If in eyes: Flush eyes with plenty of water. Get
medical attention if irritation persists.

If swallowed: Call a physician or Poison Control
Center. Drink 1 or 2 glasses of water and induce
vomiting by touching back of throat with finger. Do
not induce vomiting or give anything by mouth to an
unconscious person.

If on skin: Wash with plenty of soap and water. Get
medical attention if irritation persists.

Note to Physician: If ingested, induce emesis and
lavage stomach. Treat symptomatically.

Environmental Hazards

This pesticide is toxic to fish and aquatic inverte-
brates. Do not apply directly to water, to areas
where surface water is present, or to intertidal areas
below the mean high water mark. Drift and runoff
from treated areas may be hazardous to aquatic
organisms in neighboring areas. Do not contaminate
water when disposing of equipment wash water.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open
flame.

Chemigation

Do not apply this product through any type of irriga-
tion system.

Accu-Pak® and Action® trademarks of Syngenta
©2001 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com
SCP 806A-L1A 0201

syngenta

