

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

GFF.COLLF
PREVENTION TITT CIDES AND
TOXIC SUBJECT A ICES

Ruhi Rezaaiyan Regulatory Product Manager Sygenta Crop Protection, Inc. P.O. Box 18300 Greensboro, NC 27419-8300

MAY - 6 2003

Subject:

Dividend™ XL RTA®

EPA Reg. No. 100-826

Amendment dated February 28, 2003

Dear Ms Rezaaiyan:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following changes are made:

PERSONAL PROTECTIVE EQUIPMENT: Change "Water proof gloves" to "Chemical resistant gloves made of any waterproof material"

AGRICULTURAL USE REQUIREMENTS: Change "Water proof gloves" to "Chemical resistant gloves made of any waterproof material"

STORAGE AND DISPOSAL: For clarity, it is suggested that the Storage and Disposal section of both the base label and booklet be changed to read as follows:

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep container tightly closed when not in use. Store out of the reach of children.

SPILL RESPONSE: For minor spills, leaks, etc., follow all precautions indicated on this area and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed on an approved waste facility.

CONTAINER DISPOSAL:

2½ gallon containers - Do not reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning if allowed by state and local authorities. If burned, keep out of smoke.

Syngenta refillable container - If this is a Syngenta refillable container and is to be refilled, do not rinse or introduce any pesticide other than Dividend™ XL RTA®. Reseal and return the container to an authorized Sygenta refilling facility. If the Syngenta container is not to be refilled, triple rinse{or equivalent and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning if allowed by state and local authorities. If burned, keep out of smoke.

A copy of the label stamped "Approved with comments" is included for your records. Please submit one (1) copy of the final printed label before you release the product for shipment.

If you have any questions, please call Robert Westin at (703) 305-6700.

Sincerely,

Dennis McNeilly

Acting Product Manager (22)

Fungicide Branch

Registration Division (7505C)

Enclosure

Booklet Label

Dividend™ XL RTA®

Fungicide

A seed treatment for control of diseases of wheat

Active Ingredients:	
Difenoconazole (CAS No. 119446-68-3)	3.21%
(R)-[(2,6-dimethylphenyl)-methoxyacetylamino]-	
propionic acid methyl ester	0,27%
Other Ingredients:	96.52%
Total:	100.00%

Dividend XL RTA contains 0.31 lbs. difenoconazole and 0.025 lb. (R)-[(2)6-dimethylphenyl)-methoxyacetylamino]-propionic acid methyl ester per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-826

(For 2½ Gallons) EPA Est. 5905-GA-1

SCP 826A-L1 (2½ gals.) SCP 826A-L2 (____ gals.)

2.5 gallons Net Contents

___ gallons
Net Contents

ACCEPTED with COMMENTS In EPA Letter Dated:

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

/00-82 G

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN

If ingested, induce emesis and treat symptomatically.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spiil, Leak, Fire, or Accident), Call

1-800-888-8372

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt and long pants, socks and shoes, and waterproof gloves. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Water proof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash FPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and of Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

7/15

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with the labeling.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restrictedentry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls
- Socks plus shoes
- Waterproof gloves

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.



STORAGE AND DISPOSAL

Pesticide Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal,. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

(For 21/2 Gallons)

Do not reuse empty container. Triple rinse (or equivalent), and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

Container Refilling and Disposal

(For ____ gallons)

Thoroughly clean container before reuse **unless** this is a Syngenta refillable container. If this is a Syngenta refillable container, and is to be refilled, do not rinse or introduce any pesticide other than Dividend TM XL RTA®. Reseal and return the container to an authorized Syngenta refilling facility. If the Syngenta container is not to be refilled, triple rinse (or equivalent), and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.



GENERAL INFORMATION

Dividend XL RTA is a systemic seed dressing that controls or suppresses certain seed-borne, soil-borne, and fall foliar diseases of wheat. It is a mixture of Dividend and Apron XL® LS. Apron XL LS provides Pythium damping-off activity and Dividend provides activity on the remaining diseases claimed on this label. An EPA-approved coloring agent, Pigment Red 48 has been added to the formulation.

Note: Apron XL LS is a systemic fungicide having a specific mode of action. Apron XL LS could be subject to development of insensitive strains of fungi. Development of insensitivity cannot be predicted. Therefore, Syngenta cannot assume liability for crop damage resulting from insensitive strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Apron XL LS insensitive strains of fungi which may occur.

MIXING PROCEDURES

Dividend XL RTA is especially formulated for on-farm treatment, using standard mechanical slurry- or mist-type seed treatment equipment which accurately meters and mixes a flowable seed treatment. The equipment must provide uniform coverage of Dividend XL RTA on the seed. Uneven seed coverage may not give the desired level of disease control. This product does not require the addition of water for application. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment.

WHEAT

Apply Dividend XL RTA to wheat seed at the rates given in the following table:

Winter Wheat

Rate/CWT	Diseases Controlled	Fall Season Foliar Diseases Controlled ¹	Diseases Partiality Controlled ²
10 fl. oz.	Common Bunt ³ Dwarf Bunt Loose Smut Flag Smut Seed-borne Septoria General Seed Rots ⁴ Fusarium Seed Scab Pythium Damping-Off	Fall Season Powdery Mildew Fall Season Leaf Rust Fall Season Septoria Leaf Blotch	Common Roc: Roll (Cochliobolus sp. 1) Fusarium Root Roll Fusarium Crown Flat Take-All Rhizoctonia Roel Rot
5 fl. oz.	Common Bunt ³ Dwarf Bunt Flag Smut Seed-borne Septoria Loose Smut General Seed Rots ⁴ Fusarium Seed Scab Pythium Damping-Off		Common Root Ro: (Cochliobolus spp.) Rhizoctonia Root Rot
2.5 fl. oz. ⁵	Common Bunt Loose Smut		

¹Dividend XL RTA provides control of fall season powdery mildew, fall season leaf rust, and fall season Septoria leaf blotch in winter wheat for the first 6 weeks after planting. For full season control of these foliar diseases, use Tilt® fungicide according to label instructions.

²Partial control can either mean erratic control from good to poor or consistent control at a level below that generally considered acceptable for commercial disease control.

³Dividend XL RTA controls both seed-borne and soil-borne common bunt.

⁴General seed rots controlled include those caused by saprophytic organisms such as Penicillium and Aspergillus.

⁵The 2.5 fl. oz. rate should only be used in the following states: TX, OK, KS, CO, MO, NE, ND, SD, MN, MT.

Spring Wheat

		Diseases Partially
Rate/CWT	Diseases Controlled	Controlled ⁴
10 fl. oz.	Common Bunt'	Take-All
	Loose Smut	Common Root Rot
	Flag Smut	(Cochliobolus spp.)
	Seed-borne Septoria	Rhizoctonia Root Rot
	General Seed Rots ²	
	Fusarium Seed Scab	
	Pythium Damping-Off	
	_	
5 fl. oz.	Common Bunt	Common Root Rot
	Seed-borne Septoria	(Cochliobolus spp.)
	Loose Smut	Rhizoctonia Root Rot
	General Seed Rots ²	
	Fusarium Seed Scab	
	Pythium Damping-Off	
2.5 fl. oz. ³	Common Bunt	
	Loose Smut	
	Fusarium Seed Scab	

¹Dividend XL RTA controls both seed-borne and soil-borne common bunt.

Important: (1) Do not use treated seed for feed, food, or oil. (2) Green forage may not be grazed until 55 days after planting. (3) Do not plant any crop other than wheat within 30 days to fields in which treated seeds were planted.

Note: Federal law requires that bags containing treated seeds shall be labeled with the following information: "This seed has been treated with difenoconazole fungicide. Do not use for feed, food, or oil purposes. Store away from feeds and foodstuffs. Green forage may not be grazed until 55 days after planting. Do not plant any crop other than wheat within 30 days in fields in which treated seeds were planted."

²General seed rots controlled include those caused by saprophytic organisms, such as Penicillium and Aspergillus.

³The 2.5 fl. oz. rate should only be used in the following states: TX, OK, KS, CO, MO, NE, ND, SD, MN, MT,

⁴Partial control can either mean erratic control from good to poor or consistent control at a level below that generally considered acceptable for commercial disease control.

Apron XL®, Dividend™ XL RTA®, Tilt® and the Syngenta logo are trademarks of a Syngenta Group Company U.S. Patent No. 5,266,585

©2003 Syngenta

For non-emergency (e.g. current product information) call Syngenta Crop Protection at 1-800-334-9481.

Syngenta Crop Protection, Inc. Greensboro, North Carolina 27409 www.syngenta-us.com

SCP 826A-L1 (2½ gals.) SCP 826A-L2 (____ gals.)

DIV XL RTA 826A-L1- rr - 02/05/03

Base Label (For 2½ gallons)

Dividend™ XL RTA®

Fungicide

A seed treatment for control of diseases of wheat

Active Ingredients:	
Difenoconazole (CAS No. 119446-68-3)	3.21%
(R)-[(2,6-dimethylphenyl)-methoxyacetylamino]-	
propionic acid methyl ester	<u>0.27%</u>
Other Ingredients:	96.52%
Total:	100.00%

Dividend XL RTA contains 0.31 lb. difenoconazole and 0.025 lb. (R)-[(2-6-dimethylphenyl)-methoxyacetylamino]-propionic acid methyl ester per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See directions for use in attached booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-826

EPA Est. 5905-GA-1

2.5 gallons Net Contents

FIRST AID If on skin or clothing Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN

If ingested, induce emesis and treat symptomatically.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call
1-800-888-8372

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt and long pants, socks and shoes, and waterproof gloves. Wash thoroughly with soap and water after handling.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Container Disposal

Do not reuse empty container. Triple rinse (or equivalent), and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

Dividend™ XL RTA® and the Syngenta logo are trademarks of a Syngenta Group Company

©2003 Syngenta

Syngenta Crop Protection, Inc. Greensboro, North Carolina 27409 www.syngenta-us.com

SCP 826A-L1 (2½ gals.)

DIV XL RTA 826A-L1- rr - 02/05/03