

100-835

7-27-2001

1/7

Please read instructions on reverse before completing form.

	United States	<input type="checkbox"/> Registration	OPP Identifier Number NOTIFICATION
	Environmental Protection Agency	<input type="checkbox"/> Amendment	
	Washington, DC 20460	<input checked="" type="checkbox"/> Other	

Application for Pesticide - Section I

1. Company/Product Number 100-835	2. EPA Product Manager Mary Waller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Thiolux® Dry Flowable Micronized Sulfur	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	JUL 27 2001
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)
 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. The following changes are being made via this notification: 1) Company name and address have been updated to reflect Syngenta Crop Protection, Inc. 2) The Conditions of Sale and Warranty statement has been changed to reflect the name change. Because Syngenta has been formed by the merger of Novartis Crop Protection, Inc. and Zeneca Ag Products, we have chosen to use the former Zeneca warranty statement as the Syngenta warranty statement. No other changes occur in the statement other than the name change. 3) The copyright date reflects Syngenta. 4) Trademark statements have been updated to reflect Syngenta for those products for which Syngenta holds the trademark. 5) The internet address has been changed to reflect Syngenta. 6) Other places in the label which referring to the company name have been updated.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal <input type="checkbox"/>		Plastic <input type="checkbox"/>
*Certification must be submitted			Glass <input type="checkbox"/>		Paper <input type="checkbox"/>
If "Yes" Unit Packaging wgt. No. per Container		If "Yes" Unit Packaging wgt. No. per container		Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section - IV


1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Nan S. Padgett	Title Label Group Leader	Telephone No. (Include Area Code) 336-632-7567
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment of both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Label Group Leader	
4. Typed Name Nan S. Padgett	5. Date July 13, 2001	

2/7

NOTIFICATION

JUL 27 2001

THIOLUX®



Dry Flowable Micronized Sulfur

KEEP OUT OF REACH OF
CHILDREN.

CAUTION

SEE PRECAUTIONARY STATEMENTS
AVOID BREATHING SPRAY MIST

Active Ingredient:	
Sulfur	80.0%
<hr/>	
Other Ingredients:	20.0%
<hr/>	
Total:	100.0%
Guaranteed Analysis:	
Sulfur(s)	80.0%
Derived from elemental sulfur	

EPA Reg. No. 100-835
 EPA Est. 70992-FRA-002
 Made in France
 SCP 835A-L2E 0501

PRODUCT ID. **130**

30 POUNDS
NET WEIGHT

syngenta



Statement of Practical Treatment

If on skin: Wash with soap and plenty of water. Get medical attention.

If in eyes: Flush eyes with plenty of water for at least fifteen minutes and get medical attention.

If inhaled: Remove victim to fresh air.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

FAILURE TO FOLLOW THE DIRECTION FOR USE AND THE PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PEST CONTROL, OR ILLEGAL RESIDUES.

Thiolux is recommended for control of listed plant diseases, insects and spider mites on fruits, vegetables, and ornamentals when used as directed. Additionally, Thiolux is a readily available source of sulfur, a necessary plant nutrient.

GENERAL INFORMATION

Some crops are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use on any crop unless local use has proven sulfur to be safe in that locality. Sulfur may burn foliage or cause fruit injury when temperatures are excessively high. Do not apply at such times. Do not apply to or allow spray drift on sulfur sensitive plants such as apricots, cranberries, spinach. Apply only to dormant Anjou pears. Do not use within two (2) weeks of an oil treatment, except for dormant sprays on pears and apples. For citrus, do not apply within 21 days of an oil application. Follow all other label directions or precautions as indicated. When growing crops for processing, consult processor before applying sulfur.

Nutrient Use

Thiolux applications will provide sulfur for the plant's nutrient requirements and should be considered in the total plant nutrient applications. Thiolux can be used as a micronutrient supplement by either foliar or soil application. Thiolux is recommended but not limited to use on the following crops: alfalfa, almonds, avocados, apples, barley, all beans, broccoli, cabbage, cauliflower, carrots, citrus, corn, grapes, lawns, melons, nectarines, pears, peaches, peppers, plums, prunes, potatoes, peanuts, sorghum, soybeans, sugar beets, strawberries, tomatoes, turnips, walnuts, watermelons, and wheat. Consult your State Agricultural Experimental Station or Extension Specialist for advice in selecting treatments to best fit local conditions.

DIRECTIONS FOR APPLICATION

Fill spray tank half full with water. Pour needed amount into spray tank under agitation. If necessary, add a few drops of oil to avoid foaming. Maintain agitation throughout spray operation.

Use lower recommended rates on following host and pest recommendation table when using concentrate equipment. Use higher rates when using dilute applications.

Dilute Application

Ground Application – Apply specified rate in 20-60 gals. of water per acre.

Orchard Application – Apply specified rate in 100-800 gals. of water per acre.

Concentrate Application

Ground Application – Apply specified rate in 5 gals. or more water per acre.

Orchard Application – Apply specified rate in 20-100 gals. of water per acre. Special concentrate equipment is necessary for these uses.

Aerial Application

Apply specified rate in 3-20 gals. of water per acre. Orchard rates should not be applied in less than 10 gals. of water per acre.

Chemigation

Thiolux may be applied through sprinkler systems including center pivot, lateral move, end tow, side wheel roll, traveler, big gun, solid set, and hand move. Do not apply Thiolux through any other type irrigation system. Do not apply Thiolux through any irrigation system when wind speed favors drift beyond the area intended for treatment.

Crop injury or lack of effectiveness can result from uneven distribution of treated irrigation water. For center pivot systems, apply during one full revolution of the system. For other systems listed above (i.e., portable or solid set systems), apply during the final 15 minutes of the irrigation period. Do not operate irrigation system after application until deposit has thoroughly dried. If you have questions about system operation or calibration, contact State Extension Service specialists or chemigation equipment manufacturers.

THE FOLLOWING PRECAUTIONS MUST BE OBSERVED WHEN APPLYING PESTICIDES AND PLANT NUTRIENTS THROUGH IRRIGATION SYSTEMS:

- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. Public water system means any system for provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection system must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

THIOLUX APPLICATION RATES BY CROP AND PEST**Field Crops, Orchard Crops, Vegetables**

Crop	Pest	lbs./A
alfalfa, cotton	Atlantic mite, red spider mite	3-5
Apply as needed.		
almonds, cherries, nectarines, peaches, plums, prunes	brown rot blossom, blight, cherry leafspot, powdery mildew, rust, scab, leafspot, silver mite, flat mite, red spider mite	10-30
Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days after a period of wet weather. For brown rot of fruit, apply when fruit starts to ripen. For rust of peaches and prunes, apply anytime from late spring to fall in a preventative program.		
apples, pears	powdery mildew, scab, blister mite, red spider mite	10-30
Apply at pre-bloom and petal fall. Apply with cover sprays throughout the season. Dormant applications with oil may be made to apples and pears. Apply to Anjou pears only while dormant.		
asparagus	rust	10-30
Apply after cutting stops. Irrigate and cultivate before applying; repeat at 7 to 10-day intervals throughout the season.		
avocados	brown mite	10-30
Apply as needed.		
barley, corn, oats, rye, sorghum, soybean, wheat	suppression - red spider mite, two-spotted mite, grass banks mite, Pacific mite; control - Atlantic mite (small grain only), powdery mildew (soybean only)	6 7.5-12
Apply when disease first appears and repeat as necessary or with the regularly scheduled insect control program.		
beans, peas, potatoes, rutabagas, turnips	powdery mildew, rust, red spider mite	3-10
Begin applications in early leaf stage. Repeat application every 10-14 days with or without pesticides immediately after rainfall.		
broccoli, brussel sprouts, cabbage, cauliflower, collards, kale, peppers	powdery mildew, rust, red spider mite	3-5
Begin applications in early leaf stage. Repeat application every 10-14 days with or without pesticides immediately after rainfall.		
blackberries, boysenberries, gooseberries, loganberries	powdery mildew	6-15
Begin application when new shoots are 6 inches long and before blossoms open or when disease first appears. Repeat at 7 to 10-day intervals or as necessary.		
blueberries, currants, dewberries, huckleberries	powdery mildew	5-10
Begin application when new shoots are 6 inches long and before blossoms open or when disease first appears. Repeat at 7 to 10-day intervals or as necessary.		
carrots, garlic, onions	Petrobia mite, powdery mildew	3-10
Apply when the disease first appears and repeat as needed.		
citrus	flat mite, aardvark mite, silver mite, rust mite, bud mite, red spider mite, six spotted mite, thrips, yuma spider mite	8-20
Apply in any spray NOT CONTAINING OIL or 21 days following oil treatment.		
cucurbits - cucumbers melons squash	powdery mildew powdery mildew powdery mildew	4-6 6-15 4-10
Begin application when disease first appears. Repeat at 10 to 14-day intervals or as necessary throughout the growing season. Sulfur can injure plants, especially when temperatures reach 95° F. Do not use on sulfur sensitive varieties.		
grapes	powdery mildew, bud mite, blister mite, red spider mite	6
Begin application in late dormant stage or with cutworm spray in early shoot period. Repeat throughout season on 14-day schedule or with pesticides.		
mangoes	powdery mildew	10-30
Begin applications before bloom, repeat in-bloom, after fruit set, and 3 weeks later or as necessary.		
peanuts	rust, red spider mite, leafspot	3-5
Begin application in early leaf stage. Repeat applications as necessary, usually in 10-24 days.		
peppermint, spearmint	powdery mildew	4-6
Apply when mint is 5-6 inches tall or when disease appears. Repeat twice at 30-day intervals. Do not apply within 30 days of harvest.		
pistachio	citrus flat mite	10-30
Begin application when mites are first detected. Repeat applications as necessary.		
raspberries	powdery mildew	6-15
Apply weekly starting at first bloom color. Continue throughout fruit set; repeat as weather requires.		
strawberries	powdery mildew, red spider mite	5-16
Begin application in early leaf stage, repeat application with or without pesticides. For powdery mildew, begin at first signs of disease and repeat as necessary. Do not use on sulfur sensitive varieties.		

continued...

THIOLUX APPLICATION RATES BY CROP AND PEST (continued)**Field Crops, Orchard Crops, Vegetables**

Crop	Pest	lbs./A
sugar beets	powdery mildew, red spider mite	10-20
Apply at 10 to 30-day intervals with or without other pesticides at first symptoms.		
table beets	powdery mildew, red spider mite	3-5
Apply at 10 to 30-day intervals with or without other pesticides at first symptoms.		
tomatoes	russet mite, powdery mildew	6-20
Apply at first symptoms. Thorough coverage is required.		
walnuts	Pacific mite, almond mite, two-spotted mite, red spider mite	10-20

Apply and repeat as necessary throughout season as necessary. Tank mix with chemical mite sprays to increase effectiveness. CAUTION - Certain varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur.

Ornamentals, Grass Seed Production

Ornamentals -- all (see below for host and pest)		(ground) all --> 3-10 (air) all --> 3-20
--	--	---

Apply as aqueous spray for complete plant coverage. Begin when disease first appears and repeat at 5 to 10-day intervals as necessary.

asters, carnations, cosmos, crepe myrtle, dahlia, delphinium, gladiolus, ligustrum, sunflowers, sweet peas, violets, zinnias	powdery mildew	
calendula, cherry laurels, chrysanthemums, hollyhock, smilax	leafspot, powdery mildew	
clematis, columbine, English ivy, laurel, petunia, sage	leafspot	
roses	black spot, powdery mildew	
grass grown for seed	rust	3-5

Apply in tank mix with Tilt® Fungicide for enhanced control of rusts. Begin applications when rust pustules are evident and increasing in number from late spring through early summer. Do not exceed 12 lbs./A of Thiolux per season. Read and follow all application rates and use precautions on companion fungicide label. Do not apply this tank mix through any irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Must be stored under dry conditions.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Thiolux®

7/7

Environmental Hazards

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment and when disposing of equipment wash waters.

Physical and Chemical Hazards

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks, and flame. Do not smoke while applying this product.

Thiolux® and Tilt® trademarks of Syngenta

©2001 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 835A-L2E 0501

