

100-840
PMA-23
8-27-98 1/10

Mr. Thomas J. Parsley
Novartis Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

AUG 27 1998

Dear Mr. Parsley:

Subject: Zoril Rapid 80[™] Herbicide
EPA Registration No. 100-848
Evital 5G
EPA Registration No. 100-840 ✓
Applications and Letters Dated August 13, 1998, Request
To Amend Registration To Revise Alfalfa Directions
To Restrict Use To Certain States and Crop Rotation
Restrictions

The proposed amendments to the subject pesticide product registrations have been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, provided that you:

1. Revise the Precautionary Statements and the Statements of Practical Treatment to reflect the enclosed review of acute toxicity data that were submitted in response to the Sorflurazon RED, 8 Month Response. Please note that the appropriate statements are given for the subject two products.
2. Submit one (1) copy of your final printed label prior to releasing these products for shipment under the subject labeling.

If these conditions are not complied with, the registrations will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of these products constitutes acceptance of these conditions.

A stamped copy of each of these labels is enclosed for your your records.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosures (3)

E. Wilson: Diskette, Metolachlor: 08-27-98

2/10

(Master Label)

Evital® 5G

A GRANULAR HERBICIDE

For the control of certain annual and perennial weeds in
cranberries and alfalfa

Active Ingredient:

Norflurazon: [4-chloro-5-(methylamino)-2-
(α, α, α -trifluoro-*m*-tolyl)-3-(2*H*)-pyridazinone] 5.0%

Inert Ingredients: 95.0%

Total: 100.0%

50 POUNDS
NET WEIGHT

EPA Reg. No. 100-840
EPA Est. 39578-TX-1

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See back panel for additional precautionary statements.

NCP 840A-L1 (DRAFT)

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

AUG 27 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

100-840

APPLICATION RATES PER ACRE

80-120 lbs.	120-160 lbs.	160 lbs.
Barnyardgrass Needlegrass Rice cutgrass (Sicklegrass) Smoke grass	Broomsedge Bog rush Fall panicum Little bluestem (Poverty grass) Nutsedge (<i>Cyperus dentatus</i>) Red root Redtop grass Spike rush Summer grass	Stargrass Switchgrass Woolgrass

The rate of application will depend on many factors. Several are listed below.

Use Lower Rates Where One or More of the Following Exist:	Use Higher Rates Where One or More of the Following Exist:
Open, sanded, well-maintained bogs	A mat of vine and weed stems overlay the soil
Sprinkler irrigation	No sprinkler irrigation
Sandy soils, low in organic matter	Peat soils, high in organic matter
Perennial weed seedlings and young plants	Established perennial weed clumps
Stressed cranberry plants and sensitive varieties (Stevens and McFarlin)	Vigorous cranberry plants

NOTE: The following rates per acre should not be exceeded in any one year:

In a Newly Planted Bog	In an Established Bog
80 lbs.	160 lbs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

),,
),,
),,
),,
),,
),,
),,
),,
),,
),,

SPECIFIC ALFALFA DIRECTIONS

Evital 5G can be used for weed control in alfalfa in the states of AL, AR, AZ, CA, FL, GA, ID, LA, MS, NC, NM, NV, OK, OR, SC, TN, TX, WA, and the portions of CO, KS, and MO south of I-70.

Use Precautions

Evital 5G is only recommended for preemergent weed control in established alfalfa; do not apply to seedling alfalfa until it has emerged and been actively growing for 5 months.

Do not apply Evital 5G within 28 days of harvest.

Soil disturbances, such as rodent burrowing and fire ant mounding, may result in temporary bleaching or reduced growth in alfalfa treated with Evital 5G.

Do not apply to alfalfa planted with small grain or grass companion cover, unless injury to the companion crop is acceptable. Alfalfa seed germinating after Evital 5G is activated in the soil will not become established.

Application Timing and Rates

Evital 5G may be applied to healthy stands of established alfalfa at 20 lbs./A, do not apply to seedling alfalfa until it has emerged and been actively growing for 5 months. An additional 20 lbs./A may be applied later in the first crop year to extend late season weed control.

Application in each of the following crop years may be made at 20-40 lbs./A, depending upon weed pressure, soil texture, and cultural practices:

Use the lower rate on coarse soils (sand, loamy sand, sandy loam) and when low populations of easily controlled annual weeds are expected.

- Use the higher rate on medium- to fine-textured soils (sandy clay through clay) and when heavier weed pressure is expected. The 40 lb. maximum annual rate may be applied in a single application, or in split applications.
- Do not apply more than 40 lbs./A per crop year.

Incorporation by rainfall or irrigation is necessary to activate Evital 5G in soil before weed seeds germinate and emerge. A minimum of one inch of rainfall or irrigation is necessary to incorporate Evital 5G.

Application may be made during dormancy, as long as soil is not frozen, or in actively growing stands. Application should always follow mowing and removal of hay to ensure spray reaches the soil surface. In areas which have less than 15 inches annual moisture (rainfall and snow combined) or less than 4 inches spring rainfall, fall or winter applications are recommended.

Evital 5G will not control emerged weeds. Established weeds or weeds that emerge before activation may be controlled with postemergent herbicides that bear labeling for use in the culture of alfalfa.

Weeds Controlled

Evital 5G applications in alfalfa will control or suppress germinating seed of weed species listed below:

- **Controlled:** Cheat, Annual Bluegrass, Chickweed, Shepherdspurse, Downy Brome, Italian Ryegrass.
- **Suppressed:** Sandbur, Henbit

Rotational Crop Restrictions

Soils with less than 65% sand in AZ and CA and all soils in AL, AR, FL, GA, LA, MS, NC, NM, OK, SC, TN, TX, and the portions of CO, KS, and MO south of I-70 treated with Evital 5G may only be rotated to cotton, soybean, peanut or asparagus within 16 months after application. Soybean and peanut may show pigment loss (bleaching in leaf veins) and/or stand reduction in TX (west of I-35), southwest OK and NM if planted within 16 months following last Evital 5G application.

Soils with 65% sand or more in AZ and CA and all soils in ID, NV, OR, and WA treated with Evital 5G may only be rotated to cotton, soybean, peanut or asparagus within 16 months with a minimum of two summer periods after application. Additionally, intended rotational crops must be bioassayed prior to planting. Rotational crops may be planted when no signs of phytotoxicity (bleaching or pigment loss in leaf veins) are observed in bioassay plantings for 4 months following emergence.

Alfalfa stands treated with Evital 5G should be irrigated as in normal production the year prior to rotation in AZ and CA.

CRANBERRIES

Apply Evital 5G as a single application in the early spring, after removal of winter flood and before weed growth resumes or in the fall after harvest at least 2 weeks before winter flood. Only one application of Evital 5G per year (12

months) can be made.

Apply Evital 5G granules with any suitable calibrated ground or aerial equipment to ensure uniform coverage.

Flooding after application may reduce weed control.

Injury to the cranberry plant may occur where applications overlap or where water stands in low wet areas of a treated bog.

STORAGE AND DISPOSAL

Storage

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Completely empty bag into applicator equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling.

Statement of Practical Treatment

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

Note to Physician: If ingested, induce emesis or lavage stomach. Treat symptomatically.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:


- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate streams, ponds, or watershed areas with unused materials, or with hopper washings when cleaning equipment.

Evital 5G® trademark of Novartis

©1998 Novartis



10/10

Novartis Crop Protection, Inc.
Greensboro, North Carolina 27419
NCP 840A-L1 (DRAFT)

July 25, 1997

Revised: April 27, 1998
Revised: August 11, 1998

EVITAL5G-A [M-LABELS-E] - 8/12/98 - ccg

.....
.....
.....
.....
.....
.....
.....
.....
.....