

100-889

07/29/2005

1/9

Mertect® 340-F
RED Changes 07-13-05
Page 1 of 9

**Mertect® 340-F
Fungicide**

Flowable (Water-Dispersible Suspension)

Active Ingredient:

Thiabendazole: 2-(4-thiazolyl)benzimidazole* 42.3%

Other Ingredients: 57.7%

Total: 100.0%

*4.1 lbs. active ingredient/gallon

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-889

EPA Est.

Product of India
Formulated in the USA

SCP 889XXX
Net Contents

| |
|---|
| A C C E P T E D |
| JUL 29 2005 |
| Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticides registered under EPA Reg. No. 100-889 |

| FIRST AID | |
|--|--|
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p> | |
| <p>HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372</p> | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Causes eye and skin irritation. Avoid contact with eyes, skin, and clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes plus socks

In addition to the above, handlers making application to mushroom houses using hand held sprayers must wear chemical resistant gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

| |
|---|
| <p>User Safety Recommendations</p> <p>Users should:</p> <ul style="list-style-type: none">• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.• Remove clothing/PPE immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.• Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. |
|---|

Environmental Hazards

This pesticide is toxic to fish. Do not apply when weather conditions favor runoff or drift from the target area. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of waste.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

NOT FOR USE IN FORMULATING AGRICULTURAL PESTICIDES.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Manual seed treatments are prohibited

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

STORAGE AND DISPOSAL

Prohibitions

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store Mertect 340-F in a cool, dry place away from food, drink, and animal feeding stuffs. Do not store below 32°F (0°C).

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or alternatives allowed by state and local authorities.

**FOR BULK AND MINIBULK CONTAINERS:
CONTAINER DISPOSAL**

Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS

Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.

REFILL ONLY WITH Mertect 340-F

The contents of this container cannot be completely removed by cleaning. Refilling with materials other than Mertect 340-F will result in contamination and may weaken container. After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

GENERAL INFORMATION

Mertect 340-F is a systemic fungicide which aids in the control of many important plant diseases. If treatment is not effective following use of Mertect 340-F as recommended, a tolerant strain of fungus may be present. Consideration should be given to prompt use of other suitable non-benzimidazole fungicides. Mertect 340-F is not for use in formulating agricultural products.

CROP USE DIRECTIONS

GENERAL – SHAKE OR STIR WELL BEFORE USE.

Clean equipment before using Mertect 340-F. **When Mertect 340-F is dispersed in water, the resulting suspension must be constantly agitated.** Avoid foaming.

Use the higher recommended rates under conditions of heavier disease pressure.

Compatibility information for Mertect 340-F with other chemicals is limited. Consult your supplier or agricultural experiment station for specific recommendations.

Do not apply this product through any type of irrigation system.

Carrots – *Botrytis Cinerea* Gray Mold and *Sclerotinia Sclerotiorum* (Sclerotinia Rot): Dip carrots, before storage, for 5-10 seconds in a suspension containing 41 fl. oz. of Mertect 340-F per 100 gals. of water. Replenish the suspension when the volume is too low or when it becomes dirty.

The runoff and wastes from the dipping operation should not be discarded in a drainage which could contaminate public water systems.

Citrus – Blue mold (*Penicillium digitatum*), Green mold (*Penicillium italicum*), and stem end rot (*Diplodia natalensis*): Dip, flood, or spray harvested fruit with a suspension of Mertect 340-F in water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion. Rate will vary according to application method.

Citrus bin drenches

Use 26 fl.oz. of Mertect 340-F in 100 gal of water to provide 1000 ppm TBZ. Citrus should be drenched for three (3) minutes. Use one (1) gallon of liquid per 1500-2000 lbs citrus or 100 gallons of liquid for about 30 bins. Avoid excessive foaming, use antifoam if needed.

Sprays

Use 52-91 fl.oz. Mertect in 100 gal of water, wax/oil emulsions, or an aqueous dilution of wax/oil emulsion to provide 2000-3500 ppm of TBZ. Keep the mixture under constant agitation. Apply at a rate of one gallon per 8,000-10,000 pounds citrus fruit.

For sporulation control, apply 130 fl. oz of Mertect 340-F in 100 gal of water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion to provide 5000 ppm. Apply at a rate of one gallon per 8,000 – 10,000 lbs citrus fruit.

Mushrooms (*Agaricus* sp.) – Dry Bubble (*Verticillium fungicola*), Wet Bubble (*Mycogone pernicioso*), Cobweb (*Dactylium dendroides*) and Green Mold (*Trichoderma* spp.):

Mertect 340-F should be applied as a drench during normal watering operation. Apply during spawn preparation, any growth stage, and between breaks. Up to 8 fl. oz./1,000 sq. ft. can be applied at casting while a max of 4 fl. oz./1,000 sq. ft. can be applied at the other timings. Do not apply more than 20 fl. oz./1,000 sq. ft. of Mertect 340-F per crop. Do not apply within 12 hours prior to harvest.

Ornamental Bulbs and Corms – *Fusarium* Basal Rot and *Penicillium* Blue Mold: Clean and treat bulbs or corm within 24-48 hours of digging. Prepare suspension of 30 fl. oz. of Mertect 340-F per 100 gals. of water. Submerge stock completely in dipping suspension. Suspension temperature should be 55-75°F (13-24°C). Discard suspension (1) when it becomes dirty, (2) after using five times, or (3) after 24 hours, whichever occurs first.

***Fusarium* (Bulbs and Corms):** Dip bulbs 15-30 minutes, corms 15 minutes.

Blue Mold (Bulbs): Dip 10-15 minutes.

After treatment, dry bulbs or corms in a shaded, well-ventilated area. Curing or retarding may precede or follow treatment.

Potato – *Fusarium* Tuber Rot: Mist unwashed tubers on a conveyor line, with tumbling action, entering storage with 0.42 fl. oz. of Mertect 340-F to each 2,000 lbs. of tubers in sufficient water for complete coverage. If an additional treatment is necessary before shipping, mist the tubers at the same rate or dip the tubers for 20 seconds in a solution containing 0.42 fl. oz. of Mertect 340-F per gal. of water. Do not treat seed potatoes after cutting.

Mertect 340-F for potatoes should be used as part of an Integrated Management Program, which includes prevention of bruising during harvest and maintaining proper storage temperature and relative humidity. Neither cultural practices nor chemical measures will prevent *Fusarium* infection if potatoes are damaged or improperly stored.

Pome Fruit (Apples and Pears) – Blue Mold Rot, Bull's Eye Rot, and Gray Mold (Nest Rot, Cluster Rot), Stem End and Neck Rot: Dip, flood, or spray harvested fruit with a suspension of 16 fl. oz. of Mertect 340-F in 100 gals. of water. Do not treat for over 3 minutes. Treat apples only before and after storage for maximum decay control. Treat pears once.

9/9

Sweet Potato "Seed Roots" – Black Rot, Scurf, and Foot Rot: Dip the seed roots in a suspension containing 8 fl. oz. of Mertect 340-F per 7.5 gals. of water (107 fl. oz. per 100 gals.). Treat the seed roots for 1-2 minutes and plant immediately. Discard the suspension when the volume is too low or when it becomes dirty. Do not use the treated roots for food or feed.

Mertect® and the Syngenta logo are trademarks of a Syngenta Group Company

©2005 Syngenta

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Product of India
Formulated in the USA

SCP 889XXXX

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

Mertect 340F 889 Amd draft 1 nsp & rr 09-23-04
Mertect 340F 889 Amd draft 2 nsp 01-14-05
000100-889.20050713.Mertect 340F.per RED.doc

000100-00889.20050713.Mertect 340F per RED.doc
000100-00889.20050713.Mertect 340F per RED.pdf