

100 - 937

4/13/2001

Page 1 of 8



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number
NOTIFICATION

Application for Pesticide - Section I

1. Company/Product Number 100-937	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Primo MAXX	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

- ☐ Amendment - Explain below. ☐ Final printed labels in response to Agency letter dated _____
- ☐ Resubmission in response to Agency letter dated _____ ☐ "Me Too" Application.
- ☒ Notification - Explain below. ☐ Other - Explain below.

NOTIFICATION

APR 13 2001

Explanation: Use additional page(s) if necessary. (For Section I and Section II.).

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. The following changes are being made via this notification: 1) Company name and address have been updated to reflect Syngenta Crop Protection, Inc. 2) The Conditions of Sale and Warranty statement has been changed to reflect the name change. Because Syngenta has been formed by the merger of Novartis Crop Protection, Inc. and Zeneca Ag Products, we have chosen to use the former Zeneca warranty statement as the Syngenta warranty statement. No other changes occur in the statement other than the name change. 3) The copyright date reflects Syngenta. 4) Trademark statements have been updated to reflect Syngenta for those products for which Syngenta holds the trademark. 5) The Internet address has been changed to reflect Syngenta. 6) Other places in the label which referring to the company name have been updated.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
*Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per Container	If "Yes" Unit Packaging wgt.	No. per container	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Nan S. Padgett	Title Label Group Leader	Telephone No. (Include Area Code) 336-632-7567	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 	3. Title Label Group Leader		
4. Typed Name Nan S. Padgett	5. Date April 2, 2001		

PULL HERE TO OPEN ►

Primo[®]**MAXX**[®]**FOR TURF
GROWTH
MANAGEMENT**

For managing
growth, improving
quality and stress
tolerance, and
edging of warm-
and cool-season
turfgrasses

Active Ingredient:	
Trinexapac-ethyl	
(CAS No. 95266-40-3)	11.3%
Other Ingredients:	88.7%
Total:	100.0%

Primo MAXX is a microemulsion
concentrate.

Product of Switzerland
Formulated in the US¹

EPA Reg. No. 100-937

EPA Est. 1386-OH-1

SCP 937A-L1B 0201

**KEEP OUT OF REACH
OF CHILDREN.**

CAUTION

See additional precautionary
statements and directions for
use inside booklet.

ONE GALLON
U.S. Standard Measure

syngenta**NOTIFICATION****APR 13 2001**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter treated areas without footwear until sprays have dried.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without footwear until sprays have dried.

Observe all precautions and limitations on this label and on the labels of each product used in tank mixture with this product.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN UNEVEN GROWTH REDUCTION OR SEVERELY STUNTED TURF.

GENERAL INFORMATION

Primo MAXX for turf growth management reduces the frequency of mowing and the amount of grass clippings by reducing the growth of warm- and cool-season turfgrasses. Other benefits, such as increased turf density, increased color, and increased turf quality, are frequently observed after Primo MAXX applications.

Primo MAXX can be applied to well-maintained, quality turfgrass areas, such as residential and commercial lawns, golf courses, sod farms, sports fields, cemeteries, and similar areas. Primo MAXX is useful in the management of difficult-to-mow areas and can be used to minimize the need for edging turfgrass along sidewalks, curbs, parking lots, driveways, flower beds, fences and around posts, storage sheds, and trees.

Primo MAXX reaches the growing point by foliar uptake and is rainfast from rainfall or irrigation after one hour. Watering-in is not necessary for activation.

NOTICE TO USER: Plant tolerances to Primo MAXX have been found to be acceptable for the grasses listed on this label. Due to the large number of species and cultivars of grasses, it is impossible to test every one for tolerance to Primo MAXX. Neither the manufacturer nor the seller has determined whether or not Primo MAXX can be used safely on grasses not specified on this label. The professional user should determine if Primo MAXX can be used safely prior to commercial use. Before using Primo MAXX for grasses not listed in the application table, test Primo MAXX on a small scale first. Apply the lower recommended rate for the turf setting (lawn, fairway, etc.) and evaluate for phytotoxicity and growth inhibition to widespread use.

APPLICATION INFORMATION

Environmental conditions, management, and cultural practices that affect turf growth and vigor will influence the response of the turf to Primo MAXX applications. Fertility level, moisture availability, plant vigor, height, and frequency of mowing, etc. have been shown to influence the activity of Primo MAXX.

The rates presented in the **Application Rate Table** provide approximately 50% growth inhibition over a 4-week period with little or no discoloration of turf growing under favorable conditions.

Excessive turf growth, which may occur with high fertilization or during spring flushes, may require higher rates of Primo MAXX. Under these conditions, Primo MAXX rates may need to be increased up to 50% to provide an adequate length of control.

For extended growth suppression up to 8 weeks, when temporary discoloration can be tolerated, a maximum of twice the recommended Primo MAXX rate from the **Application Rate Table** may be applied.

Primo MAXX use rates may need to be reduced up to 50% for turfgrass grown under conditions of low fertility, compaction, or other factors which stress the turf.

Multiple applications of Primo MAXX can be made each growing season, but do not exceed a total of 7.0 fl. oz./1,000 sq. ft. (305 fl. oz./A; 19.0 pts./A) per year.

Mowing

Generally, Primo MAXX provides more suppression when turfgrasses are maintained at lower mowing heights than higher mowing heights. Application rates have been selected for typical mowing heights. The application rate may need to be adjusted depending on actual mowing conditions. To minimize possible turf injury, apply Primo MAXX then wait at least 4 hours before mowing or mow first, wait at least 1 hour, then apply Primo MAXX.

Application Timing

Apply Primo MAXX to actively-growing turf. If turf is going into dormancy because of high or low temperatures or lack of moisture, apply a lower rate of Primo MAXX.

Repeat applications of Primo MAXX may be made as soon as the turf resumes growth or more suppression is desired, but do not apply more than 7.0 fl. oz./1,000 sq. ft. per year.

Equipment

Primo MAXX may be applied with backpack sprayers, hand sprayers, boom sprayers, and with spraygun application devices. Clean spray equipment thoroughly before use. Make sure the sprayer is capable of accurate and uniform application. Calibrate the sprayer before applying Primo MAXX. Rinse sprayer with clean water after use and dispose of rinsate in an approved manner.

Mixing Instructions

Apply Primo MAXX in sufficient carrier (for example 0.5-4.0 gals. of water per 1,000 sq. ft.) to provide uniform and thorough coverage.

Primo MAXX is a microemulsion concentrate (MEC) with minimal odor because the product does not contain petroleum solvents. Primo MAXX mixes completely with water and may be tank mixed with many commonly-used pesticides and liquid fertilizers. Prepare no more mixture than is necessary for the immediate operation. Once Primo MAXX is uniformly mixed, no further agitation is required when the product is used alone, but agitation may be required for tank mixes. If using Primo MAXX in a tank mixture, observe all directions for use, sites, use rate dilution ratios, precautions, and limitations which appear on the tank mix product's label. Do not exceed any label use rate and follow the most restrictive label precautions and limitations. This product should not be mixed with any product which prohibits such mixing.

Backpack and Hand Sprayers (0.5-4.0 gals. capacity)

Primo MAXX Alone: Add all of the required amount of water to the mix tank. Add the appropriate amount of Primo MAXX, close sprayer, and vigorously shake it. Begin application.

Boom and Hand Gun Sprayers

Primo MAXX Alone: Add all of the required amount of water to the spray tank. Then, while agitating, add Primo MAXX.

Primo MAXX + Tank Mixtures: When mixing Primo MAXX with other components (carrier and tank mix pesticide products), add the products to the spray tank using the following procedure:

1. Always check the compatibility of the tank mix using a jar test with proportionate amounts of Primo MAXX, other products to be used, and the water before mixing in the spray tank. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If mixture balls-up, forms flakes, sludges, gels, oily films or layers, or precipitates, it is not compatible and do not use the combination in a tank mix.

2. Fill tank at least $\frac{1}{2}$ full of water.
3. Maintain sufficient agitation during filling to keep the tank mix uniformly suspended.
4. Add all products packaged in water-soluble bags first. Allow the packages to dissolve and the contents to completely disperse into the mix water. Then add water-dispersible granules (WG formulations) while maintaining agitation in the mix tank. Next, add wettable powders (WP formulations). Allow the products to disperse completely before other products are added.
5. Add emulsifiable concentrates (EC).
6. Add flowable liquids (FL) or suspension concentrates (SC).
7. Add Primo MAXX.
8. Add spray adjuvants and spray markers. Use surfactants approved for application to turf. Check surfactant label before use.
9. Add the remainder of the water.
10. Do not leave tank mix combinations in the spray tank for prolonged periods without agitation. Mix and apply all of the spray solution on the same day.

Note: Do not mix Primo MAXX with any product which prohibits such mixing. Do not exceed any label application rate and follow the most restrictive label precautions and limitations. Refer to the tank mix partner label(s) for further information.

Application Near and Around Monuments and Hardscape Materials

Primo MAXX, at normal dilution rates, will not stain brass, bronze, concrete, marble, granite, or other types of stone. Before using Primo MAXX around other materials, test on a small scale basis first.

Turf Pre-Stress Conditioning

Multiple Primo MAXX applications along with cultural practices such as fertilization, irrigation, drainage, mowing height, etc., will delay the onset of stress, improve stress survival, and enhance turf's recovery from stress. Apply Primo MAXX to healthy, actively growing turf before the onset of stress and continue to apply Primo MAXX throughout the growing season as long as the turf remains healthy, but do not apply more than 7.0 fl. oz./1,000 sq. ft. per year. Turfgrass root-mass is often greater in Primo MAXX-treated turf than in similar, untreated turf. As turf top growth slows, energy is redirected to below-ground plant parts, and root and rhizome production increases. Research has demonstrated that Primo MAXX may also enhance the performance of fungicides. Monthly applications of Primo MAXX at the label rate or biweekly applications at $\frac{1}{2}$ the label rate strengthen the turfgrass to help it resist disease. In addition, if mowing is less frequent, and leaf material removal is reduced, contact and systemic products stay at effective concentrations on or in the turf longer. Multiple applications of Primo MAXX will reduce water use and improve drought tolerance. Primo MAXX applications result in smaller, more compact turfgrass and leaf area is reduced for transpiration. Increased turf density reduces moisture loss from soil evaporation, and additional soil moisture is available due to increased root depth and mass. Primo MAXX also may increase carbohydrate levels which have been shown to enhance heat and cold tolerance.

POA ANNUA CONVERSION/RENOVATION

Primo MAXX can be applied to existing turf infested with stands of *Poa annua* as part of an overseeding/renovation program. Such an application allows better germination and seedling growth of more desirable turf. The use of Primo MAXX, with appropriate cultural practices, that help to ensure the vigor and growth of new seedlings, will also result in fewer clippings and thus reduce maintenance traffic on new seedlings. Because Primo MAXX is foliarly absorbed, seed germination is not affected. Apply Primo MAXX 1-5 days before seeding; and before verticutting, scalping, spiking, or other similar operations.

Temporary initial discoloration is possible with aggressive application rates of Primo MAXX to turf with *Poa annua*. The following spring, apply the upper end rate of Primo MAXX for the turf type and setting listed in the Application Rate Table. Actual conversion success will also depend on growing conditions, fertilization, rainfall, and other agronomic and environmental conditions.

BERMUDAGRASS OVERSEEDING

An application of Primo MAXX to bermudagrass will enhance the establishment of cool-season turfgrasses. Application of Primo MAXX, along with cultural practices that help ensure new seedling vigor and growth, will also result in fewer clippings and less maintenance traffic on new seedlings. Primo MAXX is foliarly absorbed, so germination and seedling growth is not affected.

Apply Primo MAXX before verticutting, scalping, spiking, or other similar operations to the bermudagrass. Apply Primo MAXX 1-5 days before seeding.

Temporary initial discoloration of turfgrass is possible with aggressive application rates of Primo MAXX. Use normal seeding rates for your area and turf setting (lawn, fairway, etc.). Actual overseeding success will also depend on growing conditions, fertilization, rainfall, and other agronomic and environmental conditions. For maintenance applications, see the Application Rate Table.

APPLICATION WITH TURF MARKING PAINT

Primo MAXX can extend the duration of marking visibility when applied before or with marking agents. Mix Primo MAXX with water first when combining with latex-based marking agents. Refer to the marking agent's product label for further instructions.

Apply Primo MAXX at 1 oz./gal. of marking paint mix to treat approximately 1,000 sq. ft. of line surface area. Refer to the **Primo MAXX + Tank Mixtures** section of this label for additional instructions.

Notes: (1) Areas treated with Primo MAXX should continue to receive regular maintenance practices, including irrigation; fertilization; and weed, disease, and insect control when necessary, and as recommended for quality turf. Because some herbicides can injure turf, tank mixes with Primo MAXX should be tested on a small scale before widespread use. (2) Primo MAXX may cause temporary yellowing. This usually disappears about one week after application. To minimize yellowing and to enhance the green color of turf, apply readily available nitrogen at 0.2-0.5 lb. of actual nitrogen per 1,000 sq. ft. If desirable, recommended rates of iron per 1,000 sq. ft. can also be used. (3) Full growth regulation by Primo MAXX begins at about 3-5 days after application.

Table 1. Recommended Primo MAXX Application Rates^{1,2}

Turf Type	Residential and Commercial Turf ³		Golf Course Fairways (Cut at 0.5 inch or less)		Golf Course Greens		Edging/Banding ⁴	
	fl. oz./1,000 sq. ft.	fl. oz./A	fl. oz./1,000 sq. ft.	fl. oz./A	fl. oz./1,000 sq. ft.	fl. oz./A	fl. oz./1,000 sq. ft.	fl. oz./A
Cool-Season								
Bentgrass	0.75	33	0.25	11	0.125	6	1.0	44
Fescue, Red	0.75	33					1.0	44
Fescue, Tall (Ky-31)	1.0	44					1.0	44
Fescue, Tall (Turf Types)	0.75	33					1.0	44
Kentucky Bluegrass	0.60	26	0.25	11			0.75	33
Mixture (Bentgrass/ <i>Poa annua</i>)			0.25 ⁵	11	0.125	7		
Mixture (K. Bluegrass/Fescue/Ryegrass)	0.75	33						
Mixture (K. Bluegrass/Ryegrass/ <i>Poa annua</i>)			0.50 ⁵	22				
Ryegrass, Annual	1.0	44					1.0	44
Ryegrass, Perennial	1.0	44	0.50	22			1.0	44
Warm-Season								
Bahiagrass	1.0	44					1-2	44-56
Bermudagrass⁶								
Bermudagrass, Common	0.75	33	0.25	11			1-2	44-56
Bermudagrass, Other Hybrids	0.25	11	0.20	9			C 50-C 75	22-33
Bermudagrass, Tifdwarf	0.20	9	0.20	9	0.062	3	C 50-C 75	22-33
Bermudagrass, Tifgreen (328)	0.25	11	0.20	9	0.125	6	0.75	33
Bermudagrass, Tifway (419)	0.38	16	0.25	11			0.75	33
Buffalograss	1.0	44					1.0	44
Carpetgrass	0.25-C 40	11-18					0.50	22
Centipedegrass	0.50	22					1.0	44
Kikuyugrass	0.30-C 50	13-22	0.30	13			1.0	44
St. Augustinegrass	0.10-0.15	4.50-6.50					C 40-C 60	16-36
St. Augustinegrass, Texas Common	0.10	4.5					0.20	9
Zoysiagrass	0.25	11	0.125	6			0.75	33

¹ These rates should provide 50% suppression of turf growth under good growing conditions for 4 weeks with minimal yellowing.

² Seedheads: At rates equal to or higher than the rates in Table 1, Primo MAXX provides seedhead suppression of hybrid bermudagrass, and partial seedhead suppression of annual bluegrass, bahiagrass, buffalograss, carpetgrass, common bermudagrass, Kentucky bluegrass, and tall fescue. Primo MAXX must be applied prior to seedhead formation. Do not apply more than 7.0 fl. oz./1,000 sq. ft. per year.

³ Including, but not limited to home lawns, parks, recreation areas, golf course roughs, cemeteries, business sites, sports fields, and sod farms.

⁴ Primo MAXX can be applied along the perimeter of lawns, sidewalks, curbs, parking lots, driveways, posts, storage buildings, pet pens, fences, or other areas. Primo MAXX can be used around trees, shrubs, flower beds, and other border plants or similar areas with no injury. Apply Primo MAXX in an 8 to 12-inch band with a single nozzle sprayer. Use the higher concentration to reduce the growth of the turfgrass into adjacent areas.

⁵ Where yellowing of *Poa annua* is a concern, use half this rate and apply more frequently. For *Poa annua* conversion/renovation, where temporary discoloration can be tolerated, twice this rate may be applied.

⁶ Bermudagrass overseeding preparation: 0.5 oz./1,000 sq. ft. for golf fairways and tees; 0.25 oz./1,000 sq. ft. for golf greens.

Precautions: (1) Do not apply Primo MAXX through any type of irrigation system. (2) Do not graze areas or feed clippings to livestock.

STORAGE AND DISPOSAL

Pesticide Storage and Disposal

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Pesticide wastes are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, they must be disposed of according to federal, state, or local procedures. Contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or dispose of by incineration, or by open burning, if allowed by state and local authorities. If container is burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

First Aid

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as nitrile, butyl, neoprene, or barrier laminate)
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Do not apply when weather conditions favor drift from treated areas.

Primo® and MAXX® trademarks of Syngenta
U.S. Patent No. 4,693,745

©2001 Syngenta

Product of Switzerland
Formulated in the USA

Syngenta Crop Protection, Inc.
Turf and Ornamental Products
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 937A-L1B 0201



This booklet manufactured using post-consumer, recycled paper.

Primo® MAXX®

FOR TURF GROWTH MANAGEMENT

For managing growth, improving quality and stress tolerance, and edging of warm- and cool-season turfgrasses

Active Ingredient:

Trinexapac-ethyl (CAS No. 95266-40-3) . . 11.3%

Other Ingredients: 88.7%

Total: 100.0%

Primo MAXX is a microemulsion concentrate.

See directions for use in attached booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-937 EPA Est. 1386-OH-1

Primo® and MAXX® trademarks of Syngenta

U.S. Patent No. 4,693,745

©2001 Syngenta

Product of Switzerland

Formulated in the USA

Syngenta Crop Protection, Inc.

Turf and Ornamental Products

Greensboro, North Carolina 27409

www.syngenta-us.com

SCP 937A-L1B 0201

ONE GALLON

U.S. Standard Measure

KEEP OUT OF REACH
OF CHILDREN.

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

First Aid

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Note to Physician: If a large amount has been ingested, lavage stomach. An aqueous suspension of activated charcoal can be given to absorb remaining toxicant. Treat symptomatically.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Do not apply when weather conditions favor drift from treated areas.

See Primo MAXX label booklet for pesticide disposal instructions.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

syngenta