



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 27, 2016

Matthew C. Flanery
Regulatory Specialist
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419

Subject: Label Amendment – Various label revisions
Product Name: APRON MAXX RTA + MOLY
EPA Registration Number: 100-945
Application Date: 03/11/2016
Decision Number: 515031

Dear Mr. Flanery:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 100-945
Decision No. 515031

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Maryam K. Muhammad by phone at 703-347-0301, or via email at Muhammad.maryam@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be 'HJ', written over a circular scribble.

Hope Johnson, Product Manager 21
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

Apron MAXX® RTA® + Moly

Fungicide

Apron MAXX RTA + Moly is a ready-to-apply seed treatment fungicide for control of certain seed-borne, soil-borne, and seedling diseases of soybeans

Active Ingredients:

Mefenoxam ¹ :	1.02%
Fludioxonil ² :	0.68%
<hr/>	
Other Ingredients:	98.30%
Total:	100.00%

¹CAS Nos. 70630-17-0 and 69516-34-3

²CAS No. 131341-86-1

Apron MAXX RTA + Moly is a flowable concentrate for seed treatment containing 0.099 lb mefenoxam and 0.066 lb fludioxonil per gallon.

Apron MAXX RTA + Moly contains 5.55% molybdenum.

KEEP OUT OF REACH OF CHILDREN.


CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-945

EPA Est.

_____ gallons
Net Contents



FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOTLINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material: barrier laminate, butyl rubber ≥14 mils, natural rubber ≥14 mils, neoprene rubber ≥14 mils, nitrile rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
-

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not contaminate water when disposing of equipment washwater or rinsate.

Physical and Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

Groundwater Advisory

Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use is permitted on-farm and in commercial seed treatment facilities. Do not use for at-plant applications (e.g. hopper box, planter box, etc.). This product is to be applied as a water-based slurry through standard slurry- or mist-type commercial seed treatment equipment.

Where mefenoxam/metalaxyl products used allow the same maximum poundage of active ingredient per acre per calendar year:

If more than one product containing mefenoxam/metalaxyl active ingredient is used on an acre during the same calendar year and the mefenoxam/metalaxyl products used allow the same maximum poundage of active ingredient per acre per calendar year, then the total poundage of all such mefenoxam/metalaxyl products used must not exceed any of the specified individual mefenoxam/metalaxyl product maximum poundage of active ingredient allowed per acre per calendar year.

Where mefenoxam/metalaxyl products used allow different maximum poundage of active ingredient per acre per calendar year:

If more than one product containing mefenoxam/metalaxyl active ingredient is used on an acre during the same calendar year and the mefenoxam/metalaxyl products used allow different maximum poundage of active ingredient per acre per calendar year, then the total poundage of all such mefenoxam/metalaxyl products used must not exceed the lowest specified individual mefenoxam/metalaxyl product maximum poundage of active ingredient allowed per acre per calendar year.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material barrier laminate, butyl rubber ≥14 mils, natural rubber ≥14 mils, neoprene rubber ≥14 mils, nitrile rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils
- Shoes plus socks

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Prior to treatment, conduct germination tests on a portion of seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed when treated with Apron MAXX RTA + Moly.

PRODUCT INFORMATION

Apron MAXX RTA + Moly is a seed treatment fungicide that protects against damping-off and seed rots due to *Pythium*, *Phytophthora*, *Fusarium*, *Rhizoctonia* spp., and early season *Phytophthora* root rot. Apron MAXX RTA + Moly also suppresses seed-borne *Sclerotinia* and *Phomopsis* spp. Apron MAXX RTA + Moly contains the active ingredient mefenoxam found in Apron XL® that provides protection against *Pythium* and early season *Phytophthora* root rot. Apron MAXX RTA + Moly also contains the active ingredient fludioxonil found in Maxim® 4FS that provides protection against the remaining diseases mentioned in this label.

RESISTANCE MANAGEMENT

GROUP	4	12	FUNGICIDES
--------------	----------	-----------	-------------------

Apron MAXX RTA + Moly contains mefenoxam, a Group 4 fungicide, and fludioxonil, a Group 12 fungicide. Mefenoxam belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis. Fludioxonil belongs to the phenylpyrrole class of chemistry which interferes with osmotic signal transduction.

Fungal populations may contain individuals naturally resistant to Group 4 or 12 fungicides and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies such as alternation with fungicides with a different mode of action and/or tank mixes established for the crop and use area.

Use should be based on an IPM program that includes field sanitation, scouting, historical information related to pesticide use, and crop rotation. The IPM program should also consider cultural, biological, and other chemical control practices.

Syngenta encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

For additional information on Fungicide Resistance Management:

- Contact Syngenta representatives at 1-800-334-9481.
- Contact your local extension specialist or certified crop advisor.
- Visit the Fungicide Resistance Action Committee (FRAC) on the web at: <http://www.frac.info>.

MIXING INSTRUCTIONS

Apron MAXX RTA + Moly is especially formulated for on-farm treatment, using standard mechanical slurry or mist-type seed treatment equipment which accurately measures and mixes a flowable seed treatment. The equipment must provide uniform coverage of Apron MAXX RTA + Moly on the seed. Uneven seed coverage may not give the desired level of disease control. The product does not require the addition of water for application. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment. Do not mix Apron MAXX RTA + Moly fungicide with other pesticides unless sufficient compatibility, loading rates, and crop safety factors are determined.

Apron MAXX RTA + Moly has been found to be compatible with some liquid inoculant products. Apron MAXX RTA + Moly may be mixed or applied sequentially with approved liquid inoculants. Consult the manufacturer of the liquid inoculants and a Syngenta Crop Protection seed treatment representative for directions before applying Apron MAXX RTA + Moly with liquid inoculants.

Apron MAXX RTA + Moly contains an EPA-approved colorant that imparts an unnatural Blue color to the seed as required by the Federal Seed Act.

SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that containers of treated seeds shall be labeled with the following statements:

- This seed has been treated with mefenoxam and fludioxonil fungicides.
- Do not use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on containers of seeds treated with Apron MAXX RTA:

- Ground Water Advisory: Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- Excess treated seed may be used for ethanol production only if: (1) By-products are not used for livestock feed, and (2) No measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.
- Do not allow children, pets, or livestock to have access to treated seed.
- Store away from feeds and foodstuffs.

- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of all excess treated seed by burying away from bodies of water. .
- Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.

CROP USE DIRECTIONS

Soybean Seed Treatment

Apron MAXX RTA + Moly provides protection against damping-off and seed rots due to *Pythium*, *Phytophthora*, *Fusarium*, *Rhizoctonia* spp., and early season *Phytophthora* root rot. Apron MAXX RTA + Moly also suppresses seed-borne *Sclerotinia* and *Phomopsis* spp. For these soybean diseases, Apron MAXX RTA + Moly is applied at the rate of 5.0 fl. oz. per 100 lbs. of seed. The specified Apron MAXX RTA + Moly rate delivers 3.75 grams active ingredient of mefenoxam and 2.5 grams active ingredient of fludioxonil per 100 kilograms of seed. If the target fields have a history of high *Phytophthora* pressure, then use 5.0 fl. oz. of Apron MAXX RTA + Moly with a tank mix of 0.16-0.48 fl. oz. of Apron XL per 100 lbs. of seed. Read and follow all label directions and restrictions for Apron XL use.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in the original container and only in a cool, dry, secure place.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining

contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Apron MAXX® RTA®, Apron XL®, Maxim®, the ALLIANCE FRAME, the Syngenta Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

©20XX Syngenta

Manufactured For:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

APR MAXX RTA+Moly 945 MAS 0614 AMEND-D MAR2016 CLEAN – rk – 7/25/16
000100-00945.20160311D.APRONMAXXRTAMOLY-AMEND-D-MAR2016-CL.PDF