



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

100-952

Date of Issuance:

6/22/2015

NOTICE OF PESTICIDE:

Registration

Reregistration

(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

PPZ 1.55% Multi-Purpose  
Fungicide

Name and Address of Registrant (include ZIP Code):

Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, NC 27419-8300

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on her/his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

EPA received a label amendment request submitted by email on 6/22/15. EPA grants this request under the authority of section 3(c)(5) of FIFRA, as amended. With this accepted labeling, all requirements set forth in the Reregistration Eligibility Decision for propiconazole have been satisfied. Therefore, EPA reregisters the product listed above. This action is taken under the authority of section 4(g)(2)(c) of FIFRA, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Submit one (1) copy of final printed labeling. Amended labeling will supersede all previously accepted labels. A copy of your label stamped "Accepted" is enclosed for your records. Products shipped after 12 months from the date of this Notice or the next printing of your label, whichever occurs first, must bear the new revised label.

Signature of Approving Official:

Shaja Joyner, Product Manager 20  
Fungicide and Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Date:

6/22/2015

Page 2 of 2  
EPA Reg. No. 100-952  
Decision No. 454601

EPA Form 8570-6

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 8/27/12
- Alternate CSF 1 dated 8/27/12

If you have any questions, please contact Erik Kraft by phone at 703-308-9358, or via email at [kraft.erik@epa.gov](mailto:kraft.erik@epa.gov).

Enclosure

**Master Label**

**PPZ 1.55% Multi-Purpose Fungicide**

**[ALT Package 1:** Ortho Dial'N Spray® Hose-End Sprayer, Other Brand of Hose-End Sprayer

**ALT Package 2:** Lock'N Spray® Cartridge Ready-to-Use Sprayer Label Spray - closed system, user just attaches hose

**ALT Package 3:** READY-SPRAY® - closed system, user just attaches hose

**ALT Package 4:** Pump-up Pressure Sprayer (user pours concentrate)]

Controls brown patch, powdery mildew, rust, blights, scab and [&] more  
[Contains Banner®]

Active Ingredient:

Propiconazole (CAS No. 60207-90-1) ..... 1.55%

Other Ingredients: ..... 98.45%

Total: ..... 100.00%


**KEEP OUT OF REACH OF CHILDREN.**


**WARNING**

See back panel booklet for additional precautionary statements.

EPA Reg. No. 100-952

EPA Est.



Stops & Prevents Diseases	Dollar spot, rust, brown patch, powdery mildew, blights, scab, leaf spot, and more!
Where to Use	On lawns. Can also be used on roses, flowers, trees, shrubs, and other ornamental plants.
Amount	38.4 oz treats up to 3,840 sq ft. For lawns, see inside label for Area to Cover
	Questions or Comments: Call [insert phone number] <b>Medical Information:</b> <b>Call [insert phone number]</b>

16 fl oz(1 pt) 473 ml

32 fl oz(1 qt) 946 ml

38.4 fl oz (1.13 L)

\_\_\_ ounce(s) pint(s), quart(s)

Net Contents

Optional Marketing Statements Below

Just mix with water and [&] spray  
For use in tank or hose-end sprayer

*[icon]* Water-Based

Systemic fungicide for turf and ornamentals

Multi-Purpose

Multi-Purpose Disease Control

Controls most common lawn and plant diseases

Prevents and [&] stops major diseases on roses, lawns, trees, shrubs and [&] flowers

Stops major diseases on roses, lawns, trees, shrubs and [&] flowers

Prevents a large variety of diseases

Effective on 30 listed diseases

Stops [a large] variety of [common] garden/lawn/rose/flower/tree/shrub/ornamental  
[plant] diseases

Stops and [&] prevents diseases in your yard and [&] garden

Stops and [&] prevents diseases on lawns, roses, flowers, ornamental(s) [plants], trees  
and [&] shrubs

Prevents [lawn] diseases before they happen

Broad spectrum [and [&] systemic] disease control

Promotes beautiful healthy lawns by controlling major diseases

[Use] on roses, flowers, lawns, trees and [&] shrubs

[Use] on lawns and [&] ornamental plants

Systemic Action

Systemic disease control

Absorbs into the plant

Can't be washed off by rain or sprinklers [watering] after spray has dried

Rain-Proof Once Spray is Dry

Rain-Proof When Spray is Dry

Won't wash off by rain or sprinklers [watering] after spray has dried.

Lasting protection and [&] control

Long lasting disease control

Works up to 4 weeks [28 days, 1 month] for certain diseases

Up to 4-week [28-day, 1-month] protection for certain diseases

Stimulates growth of newly planted cool-season lawns and one quart (entire contents)  
treats up to 3,200 sq ft.

Treats [Covers] up to 1,600 sq ft [*pint*], up to 3,200 sq ft [*quart*], up to 3,840 sq ft.

*[ounce]*

Larger Size, 20% More!!

<b>FIRST AID</b>	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move the person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>HOT LINE NUMBER</b> For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call <b>1-800-888-8372</b>	

**PRECAUTIONARY STATEMENTS** *[optional illustration]*

**Hazards to Humans and Animals**

**WARNING**

Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if inhaled. Harmful if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse. Wear protective eyewear (goggles, face shield, or safety glasses), long-sleeved shirt and long pants, socks, shoes, and gloves.

## Environmental Hazards

This pesticide is toxic to fish and shrimp. Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential, or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

**Or**

**NOTICE:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

**OPEN**  
**Resealable Label**  
**for Directions & Precautions**  
**PRESS TO RESEAL**

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Shake well before using.

#### **Use Restrictions**

Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.

Do not allow adults, children, or pets to enter the treated area until sprays have dried.

#### **STORAGE AND DISPOSAL** *[Optional Illustration]*

Do not contaminate water, food, or feed by storage or disposal.

#### **Pesticide Storage**

Store in a cool, dry place, away from heat or open flame.

#### **Pesticide Disposal**

**If empty:** Do not reuse this container. Place in trash or offer for recycling if available.

**If partly filled:** Call your local solid waste agency or call 1-800-334-9481 for disposal instructions. Never place unused product down any indoor or outdoor drain.

[End of Master Label text]

**[ALT Package 1: Ortho Dial'N Spray Hose-End Sprayer, Other Brand of Hose-End Sprayer]**

**PRODUCT INFORMATION**

PPZ 1.55% Multi-Purpose Fungicide [alternate brand name] stops and [&] prevents diseases without harming lawns or delicate blooms and buds. [Continues protecting up to 28 days for certain diseases.]

Plus, PPZ 1.55% Multi-Purpose Fungicide [alternate brand name] provides lasting protection [up to 28 days (for certain diseases)] because [since] lawn and plant foliage absorb it, and it won't wash off with rain or watering after spray has dried.

PPZ 1.55% Multi-Purpose Fungicide [alternate brand name] promotes beautiful, healthy and vigorous lawns, roses, flowers, shrubs and shade trees by controlling their major [common] diseases.

PPZ 1.55% Multi-Purpose Fungicide [alternate brand name] stops and prevents lawn and plant diseases. Lawn and plant foliage absorb PPZ 1.55% Multi-Purpose Fungicide resulting in protection that lasts up to 4 weeks (for certain diseases).

Thank you for choosing [company/brand name]. You'll like the results and so will your lawn and garden.

<b>PRODUCT FACTS</b>	
[Treats] [Covers] up to 1,600 sq ft. [pint]	
[Treats] [Covers] up to 3,200 sq ft. [quart]	
<b>STOPS &amp; PREVENTS DISEASES</b>	Dollar spot, rust, brown patch, powdery mildew, blights, scab, leaf spot and [&] more!
[optional illustration(s)]	
<i>[option to list other diseases here that are listed elsewhere on this label]</i>	
<b>WHERE TO USE</b>	[On] lawns. Can also be used on roses, flowers, trees, shrubs and [&] other ornamental plants
[optional illustration(s)]	
<i>[option to list other sites here that are listed elsewhere on this label]</i>	
<b>AMOUNT TO USE</b>	<b>4 tsp per gallon of water</b>
	For lawns, see inside label for "AREA TO COVER."
[DNS illustration]	Easy to apply with [an Ortho Dial 'n Spray] [a hose-end sprayer] [a tank sprayer].
[phone icon]	<b>Questions, Comments or Medical Information?</b> call [telephone number] e-mail [e-mail address]



---

## USE PRECAUTIONS

---

Specially formulated for residential use  
Do not use fruit from treated crab apple trees for food purposes.  
Do not apply to “Stanley” type plums.

---

## APPLICATION INSTRUCTIONS

---

### ESTABLISHED LAWNS *[optional illustration]*

#### When to Mow

Apply after mowing **or** allow sprayed lawn to completely dry before mowing.

#### How to Apply

*[walking man icon]*

- Measure lawn. Calculate square feet by multiplying length times width.
- Spray evenly over measured area. Treat entire lawn.

<b>MIXING INSTRUCTIONS</b> for Established Lawns
--

<b>Amount to Use:</b> 4 tsp per gallon of water for each 33 1/3 sq ft.
--

<i>[DNS illustration]</i> When using <b>Ortho Dial 'n Spray</b> :
---

1. Set dial to 4 tsp.
-----------------------

<i>[pint]</i>
---------------

2. Add 16 fl oz (entire contents) to sprayer jar. <b>DO NOT</b> add water.
--

3. Spray evenly over 800 sq ft.
---------------------------------

<i>[quart]</i>
----------------

4. Add 32 fl oz (entire contents) to sprayer jar. <b>DO NOT</b> add water.
--

5. Spray evenly over 1,600 sq ft.
-----------------------------------

1 Tablespoon (Tbsp) = 3 teaspoons (tsp)

1 fl oz = 2 Tbsp

*Alternative “MIXING INSTRUCTIONS” to be used with the optional column “AREA TO COVER.”*

<b>MIXING INSTRUCTIONS</b> for Established Lawns
--

<b>Amount to Use:</b> 4 tsp per gallon of water
---

<i>[DNS illustration]</i> When using <b>Ortho Dial 'n Spray</b> :
---

1. Set dial to 4 tsp.
-----------------------

<i>[pint]</i>
---------------

2. Add 16 fl oz (entire contents) to sprayer jar. <b>DO NOT</b> add water.
--

3. See table for “AREA TO COVER.”
-----------------------------------

<i>[quart]</i>
----------------

4. Add 32 fl oz (entire contents) to sprayer jar. <b>DO NOT</b> add water.
--

5. See table for “AREA TO COVER.”
-----------------------------------

1 Tablespoon (Tbsp) = 3 teaspoons (tsp.)

1 fl oz = 2 Tbsp

### When to Apply

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified, treat at the first sign of disease. Reapply as directed in “COMMENTS” until lawn shows signs of recovery. Resume applications if disease reoccurs.

<b>DISEASES CONTROLLED</b>	<b>AREA TO COVER</b> [Entire contents treats] [Using one [quart] [pint] treats]	<b>COMMENTS</b>
Anthracnose	[pint] 1,600 sq ft [quart] 3,200 sq ft	Apply every 14 days.
Brown patch	[pint] 800 sq ft [quart] 1,600 sq ft	Start applications in May or June before disease occurs. Apply every 14 days. For southern lawns, start applications at first sign of disease in spring or fall.
Dichondra rust on dichondra lawns	[pint] 800 sq ft [quart] 1,600 sq ft	Apply every 21 days.
Dollar spot	[pint] 1,600 sq ft [quart] 3,200 sq ft	Apply every 28 days.
Fusarium patch	[pint] 800 sq ft [quart] 1,600 sq ft	Apply every 28 days from fall to early spring.
Gray leaf spot	[pint] 1,600 sq ft [quart] 3,200 sq ft	Apply every 14 days in the summer.
Leaf spot (melting out)	[pint] 1,600 sq ft [quart] 3,200 sq ft	Apply every 14 days.
Pink patch	[pint] 800 sq ft [quart] 1,600 sq ft	Apply every 21 days
Powdery mildew	[pint] 1,600 sq ft [quart] 3,200 sq ft	Apply every 21-28 days.
Red thread	[pint] 800 sq ft [quart] 1,600 sq ft	Apply every 21 days.
Rust	[pint] 1,600 sq ft [quart] 3,200 sq ft	Apply every 21-28 days.
Snow mold	[pint] 800 sq ft [quart] 1,600 sq ft	Apply twice in the fall prior to snow cover waiting 14 days between each application.
Stripe smut	[pint] 1,600 sq ft [quart] 3,200 sq ft	Apply in fall after grass is dormant or in early spring before grass starts to grow.
Summer patch	[pint] 800 sq ft [quart] 1,600 sq ft	Start applications in late May or early June and apply every 14 days up to 6 times.
Take-all patch	[pint] 800 sq ft [quart] 1,600 sq ft	Apply twice in the fall (September and October) and twice in the spring (April and May).  Southern lawns also need to be treated twice in the summer (June-August).

## ADDITIONAL USE - GROWTH STIMULATOR *[optional illustration]*

In addition to its fungicide uses, *PPZ 1.55% Multi-Purpose Fungicide* can be used as a growth stimulator for faster establishment of newly planted **cool-season** lawns.

*[alternative text]*

In addition to preventing and [&] stopping major diseases on roses, lawns, trees, shrubs and [&] flowers, *PPZ 1.55% Multi-Purpose Fungicide* promotes faster establishment of newly planted **cool-season** lawns.

### MIXING INSTRUCTIONS for Newly Planted *Cool Season* Lawns

**Amount to Use:** 4 tsp per gallon of water for each 66 2/3 sq ft

*[DNS illustration]* When using **Ortho Dial 'n Spray**:

1. Set dial to 4 tsp.

*[pint]*

2. Add 16 fl oz (entire contents) to sprayer jar. **DO NOT** add water.

3. Spray evenly over 1,600 sq ft.

*[quart]*

4. Add 32 fl oz (entire contents) to sprayer jar. **DO NOT** add water.

5. Spray evenly over 3,200 sq ft.

1 Tablespoon (Tbsp) = 3 teaspoons (tsp)

1 fl oz = 2 Tbsp

### When to Apply

Lawn Seedlings: Apply at the 2- to 3-leaf stage of growth for faster root development and top growth.

## ORNAMENTAL PLANTS [ROSES, FLOWERS, SHRUBS AND [&] SHADE TREES]

*[optional illustration]*

When used as directed, this product is safe to plants and controls the diseases listed. This product may injure African violets, begonias, Boston fern, geraniums or other unlisted plants, and it may not control unlisted diseases.

### MIXING INSTRUCTIONS for Ornamental Plants [Roses, Flowers, Shrubs and [&] Shade Trees]

**Amount to Use:** 4 tsp per gallon of water

*[DNS illustration]* When using **Ortho Dial 'n Spray**:

1. Set dial to 4 tsp.

2. Pour product into sprayer to fill jar one-quarter to one-half full. **DO NOT** add water.

3. After spraying, unused product can be poured back into its original container.

1 Tablespoon (Tbsp) = 3 teaspoons (tsp)

1 fl oz = 2 Tbsp

### How to Apply

- Apply product to the point just when it begins to run off the leaves.

*[alternative text]*

- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces, flowers, stems and branches) including new growth.
- Allow spray to completely dry before watering.

### When to Apply

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified (in COMMENTS), treat at the first sign of disease and reapply every 21 days until plant shows signs of recovery. Resume applications if disease reoccurs.

PLANTS	DISEASES CONTROLLED	COMMENTS
Amelanchier	Powdery mildew	
Apple (non-bearing)	Scab	Start applications when leaves start growing in the spring.
Apple (non-bearing)	Rust	Start applications when buds just begin to break open. Make 3 applications.
Apple (non-bearing)	Powdery mildew	Start applications in early spring.
Ash	Powdery mildew	
Azalea	Monilinia flower blight	Apply when there is 5% bloom and again at 70% bloom.
Calendula	Powdery mildew	
Carnation	Rust	
Cherry (ornamental)	Powdery mildew, leaf spot	Start applications in early spring.
Chrysanthemum	Ray Blight	
Crab apple	Scab	Start applications when buds just begin to break open.
	Rust	Start applications when buds just begin to break open. Make 3 applications.
	Powdery mildew	
Crape myrtle	Powdery mildew	
Delphinium	Powdery mildew	
Dogwood	Anthrachnose	
Douglas Fir, Shasta Fir	Needle Rust	For Shasta Fir, apply in May-June before disease occurs.
Gomphrena	Alternaria leaf spot	
Hawthorns, Poplars	Rust	
Iris	Rust	
Juniper	Phomopsis blight	
Lilac	Powdery mildew	
Marigold	Alternaria leaf spot	
Monarda	Powdery mildew	
Oak	Leaf Spot	
Phlox	Powdery mildew	
Red tip photinia	Leaf spot	
Pine	Tip blight	Start applications in mid-late winter (November-January) and apply every 2 months up to 3 times.

PLANTS	DISEASES CONTROLLED	COMMENTS
Pyracantha	Blight	
Rhododendron	Flower blight	Apply when there is 5% bloom and again at 70% bloom.
Roses	Powdery mildew	[Thorough coverage is needed for best results.] Start applications when buds just begin to break open.
Roses	Black spot and [&] rust	<i>[optional text]</i> For severe black spot on roses alternate every 7 to 10 days between this product and Rose Pride Funginex® Rose & Shrub Disease Control.  <i>[optional text]</i> For severe black spot on roses use Rose Pride Funginex Rose & Shrub Disease Control.
Snapdragons	Rust	
Sweet William	Leaf spot	
Zinnia	Powdery mildew	

### BEARING FRUIT AND [&] NUT TREES]

[optional illustration]

When used as directed, this product is safe to plants and controls the diseases listed.

#### MIXING INSTRUCTIONS for Bearing Fruit and [&] Nut Trees]

**Amount to Use:** 4 tsp per gallon of water

[DNS illustration] When using **Ortho Dial 'n Spray:**

1. Set dial to 4 tsp.
2. Pour product into sprayer to fill jar one-quarter to one-half full. **DO NOT** add water.
3. After spraying, unused product can be poured back into its original container.

1 Tablespoon (Tbsp) = 3 teaspoons (tsp)

1 fl oz = 2 Tbsp

#### How to Apply

- Apply 4 tsp (1 1/3 ounce) of product to 250 to 500 square feet of bearing fruit and nut trees. Apply spray mixture to the point just when it begins to run off the leaves.

*[alternative text]*

- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces, flowers, stems and branches) including new growth.
- Allow spray to completely dry before watering.

#### When to Apply

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified (in COMMENTS), treat at the first sign of disease and reapply every 21 days until plant shows signs of recovery. Resume applications if disease reoccurs.

<b>BEARING FRUIT AND NUT TREES</b>	<b>DISEASES CONTROLLED</b>	<b>COMMENTS</b>
Apricots, Cherries, Nectarines, Peaches, Plums and Prunes	Brown Rot Blossom blight, Powdery Mildew, Cherry Leaf Spot and Fruit Brown Rot	Start applications in early Spring and spray every 21 days, making no more than 5 applications for the entire season. Up to 3 applications may be made from early bloom through petal fall. Up to 2 applications may be made from the end of petal fall through harvest.

## **SWEET CORN**

*[optional illustration]*

When used as directed, this product is safe to plants and controls the diseases listed.

<b>MIXING INSTRUCTIONS</b> for Sweet Corn
<p><b>Amount to Use:</b> 4 tsp per gallon of water  <i>[DNS illustration]</i> When using <b>Ortho Dial 'n Spray</b>:</p> <ol style="list-style-type: none"> <li>1. Set dial to 4 tsp.</li> <li>2. Pour product into sprayer to fill jar one-quarter to one-half full. <b>DO NOT</b> add water.</li> <li>3. After spraying, unused product can be poured back into its original container.</li> </ol>

1 Tablespoon (Tbsp) = 3 teaspoons (tsp)

1 fl oz = 2 Tbsp

### **How to Apply**

- Apply 4 tsp (1 1/3 ounce) of product to 250 to 500 square feet of sweet corn.

*[alternative text]*

- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces).
- Allow spray to completely dry before watering.

### **When to Apply**

Apply when disease first appears and continue every 14 days, making no more than 4 applications. Do not apply to sweet corn within 14 days of harvest.

<b>SWEET CORN</b>	<b>DISEASES CONTROLLED</b>	<b>COMMENTS</b>
Sweet Corn	Leaf blights, Rust, Gray Leaf Spot and Eye Spot	Apply when disease first appears and continue every 14 days, making no more than 4 applications. Do not apply to sweet corn within 14 days of harvest.

---

**IMPORTANT** *[optional illustration]*

---

*PPZ 1.55% Multi-Purpose Fungicide* is a preventative fungicide. Applications should be made before diseases occur for best results.

*[alternative text]*

This product does not repair [No fungicide repairs] plants or lawns damaged by diseases. It *PPZ 1.55% Multi-Purpose Fungicide [alternate brand name]* stops [the] disease(s) and prevents reinfestation of treated areas. As treated plants and lawns grow they recover from disease.

If you need more information about lawn and ornamental diseases, contact your local County Extension Service.

*[Reentry icon]* People and pets may enter treated area after spray has dried.

**[ALT Package 2: Lock 'N Spray Cartridge Ready-to-Use Sprayer Label]**

---

**PRODUCT INFORMATION**

---

**FOR USE WITH THE Lock 'N Spray [logo] [SYSTEM]**

[Product name] stops and [&] prevents diseases without harming lawns or delicate blooms and buds. [Continues protecting up to 28 days for certain diseases.]

Plus, [product name] provides lasting protection [up to 28 days (for certain diseases)] because [since] lawn and plant foliage absorb it, and it won't wash off with rain or watering after spray has dried.

[Product name] promotes beautiful, healthy and vigorous lawns, roses, flowers, shrubs and shade trees by controlling their major [common] diseases.

[Product name] in the unique Lock 'N Spray System makes it fast and easy to stop and prevent lawn and plant diseases. Lawn and plant foliage absorb [product name] resulting in protection that lasts up to 4 weeks (for certain diseases). Simply attach the Lock 'N Spray sprayer to this cartridge and the correct amount of PPZ 1.55% Multi Purpose Fungicide] and water mix automatically. No measuring. No mixing. No messy cleanup. And Lock 'N Spray cartridges are spill-proof!

The unique Lock 'N Spray System makes PPZ 1.55% Multi Purpose Fungicide easy to apply -- simply attach the Lock 'N Spray sprayer to this cartridge and the correct amount of [product name] and water mix automatically. No measuring. No mixing. No messy cleanup. And Lock 'N Spray cartridges are spill-proof!

Thank you for choosing [brand or company name]. You'll like the results and so will your lawn and garden.

<b>PRODUCT FACTS</b>	
[Treats] [Covers] up to 1,600 sq ft. [pint]	
[Treats] [Covers] up to 3,200 sq ft. [quart]	
<b>STOPS &amp; PREVENTS DISEASES</b>	Dollar spot, rust, brown patch, powdery mildew, blights, scab, leaf spot and [&] more!
[optional illustration(s)]	
[option to list other diseases here that are listed elsewhere on this label]	
<b>WHERE TO USE</b>	[On] lawns. Can also be used on, roses, flowers, trees, shrubs and [&] other ornamental plants.
[optional illustration(s)]	
[option to list other sites here that are listed elsewhere on this label]	
[phone icon]	<b>Questions, Comments or Medical Information? call [telephone number]</b>



---

## USE PRECAUTIONS

---

Specially formulated for residential use

Do not use fruit from treated crab apple trees for food purposes.  
Do not apply to “Stanley” type plums.

<b>HOW TO USE THE Lock ‘N Spray [logo] SYSTEM</b>	
<b>CONNECT</b> <i>[illustration #1]</i>	Connect sprayer to hose.
<b>ATTACH</b> <i>[illustration #2]</i>	Attach cartridge to sprayer. Turn cartridge until it clicks into place.
<b>SPRAY</b> <i>[illustration #3]</i>	To begin spraying, rotate sprayer nozzle tip to direct spray up or down, or remove for a 20 to 25 foot (ft) jet spray. Turn on water and squeeze trigger.
<b>FINISH</b> <i>[Reverse illustration #2]</i>	<ul style="list-style-type: none"> <li>To stop spraying, release trigger, unlock cartridge and remove from sprayer. (No need to turn off water.)</li> <li>To clean sprayer, flush with water, squeezing and releasing trigger in 2 short bursts. Turn off water.</li> <li>Squeeze trigger to drain water from sprayer, then disconnect sprayer from hose.</li> </ul>
Cartridge reseals itself automatically - No spills, no leaks!	

---

## APPLICATION INSTRUCTIONS

---

### ESTABLISHED LAWNS *[optional illustration]*

#### When to Mow

Apply after mowing **or** allow sprayed lawn to completely dry before mowing.

#### How to Apply

*[walking man icon]*

- Measure lawn. Calculate square feet by multiplying length times width.
- Spray evenly over measured area. Treat entire lawn.

#### Area to Cover

One pint (entire contents) treats [covers] 800 sq ft.

One quart (entire contents) treats [covers] 1,600 sq ft.

Alternative text - add “AREA TO COVER” column to the table below.

### When to Apply

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified, treat at the first sign of disease. Reapply as directed in “COMMENTS” until lawn shows signs of recovery. Resume applications if disease reoccurs.

<b>DISEASES CONTROLLED</b>	<b>AREA TO COVER</b> [Entire contents treats] [Using [one quart (32 oz)] [one pint (16 oz)] treats] <i>[optional column]</i>	<b>COMMENTS</b>
Anthracnose	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft	Apply every 14 days.
Brown patch	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft	Start applications in May or June before disease occurs. Apply every 14 days.  For southern lawns, start applications at first sign of disease in spring or fall.
Dichondra rust on dichondra lawns	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft	Apply every 21 days.
Dollar spot	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft	Apply every 28 days.
Fusarium patch	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft	Apply every 28 days from fall to early spring.
Gray leaf spot	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft	Apply every 14 days in the summer.
Leaf spot (melting out)	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft	Apply every 14 days.
Pink patch	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft	Apply every 21 days
Powdery mildew	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft	Apply every 21-28 days.
Red thread	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft	Apply every 21 days.
Rust	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft	Apply every 21-28 days.
Snow mold	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft	Apply twice in the fall prior to snow cover waiting 14 days between each application.
Stripe smut	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft	Apply in fall after grass is dormant or in early spring before grass starts to grow.
Summer patch	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft	Start applications in late May or early June and apply every 14 days up to 6 times.

DISEASES CONTROLLED	AREA TO COVER [Entire contents treats] [Using [one quart (32 oz)] [one pint (16 oz)] treats] <i>[optional column]</i>	COMMENTS
Take-all patch	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft	Apply twice in the fall (September and October) and twice in the spring (April and May).  Southern lawns also need to be treated twice in the summer (June-August).

**ADDITIONAL USE - GROWTH STIMULATORS** *[optional illustration]*

In addition to its fungicide uses, *[product name]* can be used as a growth stimulator for faster establishment of newly planted **cool season** lawns.

*[alternative text]*

In addition to preventing and [&] stopping major diseases on roses, lawns, trees, shrubs and [&] flowers, *[product name]* promotes faster establishment of newly planted **cool season** lawns.

**Area to Cover**

One pint (entire contents) treats [covers] 1,600 sq ft.  
One quart (entire contents) treats [covers] 3,200 sq ft.

**When to Apply**

Lawn Seedlings: Apply at the 2- to 3-leaf stage of growth for faster root development and top growth.

**ORNAMENTAL PLANTS [ROSES, FLOWERS, SHRUBS, AND [&] SHADE TREES**  
*[optional illustration]*

When used as directed, this product is safe to plants and controls the diseases listed. This product may injure African violets, begonias, Boston fern, geraniums or other unlisted plants, and it may not control unlisted diseases.

**How to Apply**

- Apply product to the point just when it begins to run off the leaves. *[alternative text]*
- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces, flowers, stems and branches) including new growth.
- Allow spray to completely dry before watering.

### When to Apply

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified (in COMMENTS), treat at the first sign of disease and reapply every 21 days until plant shows signs of recovery. Resume applications if disease reoccurs.

PLANTS	DISEASES CONTROLLED	COMMENTS
Amelanchier	Powdery mildew	
Apple (non-bearing)	Scab	Start applications when leaves start growing in the spring.
Apple (non-bearing)	Rust	Start applications when buds just begin to break open. Make 3 applications.
Apple (non-bearing)	Powdery mildew	Start applications in early spring.
Ash	Powdery mildew	
Azalea	Monilinia flower blight	Apply when there is 5% bloom and again at 70% bloom.
Calendula	Powdery mildew	
Carnation	Rust	
Cherry (ornamental)	Powdery mildew, leaf spot and [&]	Start applications in early spring.
Crab apple	Scab	Start applications when buds just begin to break open.
	Rust	Start applications when buds just begin to break open. Make 3 applications.
	Powdery mildew	
Chrysanthemum	Ray Blight	
Crape myrtle	Powdery mildew	
Delphinium	Powdery mildew	
Dogwood	Anthraco nose	
Douglas Fir, Shasta Fir		For Shasta Fir, apply in May-June before disease occurs.
Gomphrena	Alternaria leaf spot	
Iris	Rust	
Hawthorn	Rust	
Juniper	Phomopsis blight	
Lilac	Powdery mildew	
Marigold	Alternaria leaf spot	
Monarda	Powdery mildew	
Oak	Leaf Spot	
Phlox	Powdery mildew	
Red tip photinia	Leaf spot	
Pine	Tip blight	Start applications in mid-late winter (November-January) and apply every 2 months up to 3 times.
Poplars	Rust	
Pyracantha	Blight	
Rhododendron	Flower blight	Apply when there is 5% bloom and again at 70% bloom.

<b>PLANTS</b>	<b>DISEASES CONTROLLED</b>	<b>COMMENTS</b>
Roses	Powdery mildew	[Thorough coverage is needed for best results.] Start applications when buds just begin to break open.
Roses	Black spot and [&] rust	<i>[optional text]</i> For severe black spot on roses alternate every 7 to 10 days between this product and Rose Pride Funginex Rose & Shrub Disease Control.  <i>[optional text]</i> For severe black spot on roses use Rose Pride Funginex Rose & Shrub Disease Control.
Snapdragons	Rust	
Sweet William	Leaf spot	
Zinnia	Powdery mildew	

### **BEARING FRUIT AND [&] NUT TREES]**

[optional illustration]

When used as directed, this product is safe to plants and controls the diseases listed.

#### **How to Apply**

- Apply 4 tsp (1 1/3 ounce) of product to 250 to 500 square feet of bearing fruit and nut trees. Apply spray mixture to the point just when it begins to run off the leaves.

*[alternative text]*

- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces, flowers, stems and branches) including new growth.
- Allow spray to completely dry before watering.

#### **When to Apply**

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified (in COMMENTS), treat at the first sign of disease and reapply every 21 days until plant shows signs of recovery. Resume applications if disease reoccurs.

<b>BEARING FRUIT AND NUT TREES</b>	<b>DISEASES CONTROLLED</b>	<b>COMMENTS</b>
Apricots, Cherries, Nectarines, Peaches, Plums and Prunes	Brown Rot Blossom blight, Powdery Mildew, Cherry Leaf Spot and Fruit Brown Rot	Start applications in early Spring and spray every 21 days, making no more than 5 applications for the entire season. Up to 3 applications may be made from early bloom through petal fall. Up to 2 applications may be made from the end of petal fall through harvest.

## SWEET CORN

*[optional illustration]*

When used as directed, this product is safe to plants and controls the diseases listed.

### How to Apply

- Apply 4 tsp (1 1/3 ounce) of product to 250 to 500 square feet of sweet corn.

*[alternative text]*

- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces).
- Allow spray to completely dry before watering.

### When to Apply

Apply when disease first appears and continue every 14 days, making no more than 4 applications. Do not apply to sweet corn within 14 days of harvest.

SWEET CORN	DISEASES CONTROLLED	COMMENTS
Sweet Corn	Leaf blights, Rust, Gray Leaf Spot and Eye Spot	Apply when disease first appears and continue every 14 days, making no more than 4 applications. Do not apply to sweet corn within 14 days of harvest.

---

## IMPORTANT *[optional illustration]*

---

PPZ 1.55% Multi-Purpose Fungicide is preventative fungicide. Applications should be made before diseases occur for best results.

*[alternative text]*

This product does not repair [No fungicide repairs] plants or lawns damaged by diseases. It *[product name]* stops [the] disease(s) and prevents reinfestation of treated areas. As treated plants and lawns grow they recover from disease.

If you need more information about lawn and ornamental diseases, contact your local County Extension Service.

*[reentry icon]* People and pets may enter treated area after spray has dried.

**[ALT Package 3: READY-SPRAY - closed system [or other brand Hose-end Sprayer], user just attaches hose]**

---

**PRODUCT INFORMATION**

---

Treats [Covers] up to 1,600 sq ft [*pint*]; up to 3,200 sq ft [*quart*], up to 3,480 sq ft [*ounce*].

<b>Product Facts</b>	
<b>STOPS &amp; PREVENTS DISEASES</b>	Dollar spot, rust, brown patch, powdery mildew, blights, scab, leaf spot and more!
<b>WHERE TO USE</b>	On lawns. Can also be used on roses, flowers, trees, shrubs, and other ornamental plants
<b>AMOUNT TO USE</b>	[16 oz treats up to 1,600 sq ft.] [32 oz treats up to 3,200 sq ft.] [38.4 oz treats up to 3,480 sq ft.] For lawns, see inside label for Area to Cover
[image of telephone]	<b>Questions, Comments, or Medical Information call 1-800-888-8372.</b>

---

**USE PRECAUTIONS**

---

Specially formulated for residential use

Do not use fruit from treated crab apple trees for food purposes.  
Do not apply to “Stanley” type plums.

---

**APPLICATION INSTRUCTIONS**

---

**ESTABLISHED LAWNS** [*optional illustration*]

**When to Mow**

Apply after mowing **or** allow sprayed lawn to completely dry before mowing.

**How to Apply**

[*walking man icon*]

- Measure lawn. Calculate square feet by multiplying length times width.
- Spray evenly over measured area. Treat entire lawn.

## How to Use the Ready-Spray System

- Connect**
1. Shake well before using
  2. Connect sprayer to hose
  3. Turn on water
- Spray**
1. To begin spraying, point nozzle in the direction you want to spray.
  2. [Bend small plastic tab back and] Turn knob [clockwise] to ON position.
  3. Spray evenly over measured area. See table page \_ for “AREA to COVER.”
- Finish**
1. To stop spraying, [turn knob counter clockwise] to OFF position.
  2. Turn off water.
  3. Relieve water pressure by [bending plastic tab back and] turning knob to ON position until water slows to a drip. Then turn knob back to OFF position.
  4. Disconnect sprayer from hose.

## Area to Cover

[One pint (entire contents) treats [covers] [up to] 800 sq ft. [1,600 sq ft.]  
[One quart (entire contents) treats [covers] [up to] 1,600 sq ft. [3,200 sq ft.]  
[38.4 ounces treats [covers] up to 3,840 sq ft.]  
*[Alternative text - add “AREA TO COVER” column to the table below.]*

## When to Apply

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified, treat at the first sign of disease. Reapply as directed in “COMMENTS” until lawn shows signs of recovery. Resume applications if disease reoccurs.



<b>DISEASES CONTROLLED</b>	<b>AREA TO COVER</b> [Using one quart (32 oz) ] [Using one pint (16 oz)] [38.4 ounces] treats	<b>COMMENTS</b>
Anthracnose	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft [38.4 ounce] 3,840 sq ft	Apply every 14 days.
Brown patch	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft [38.4 ounce] 3,840 sq ft	Start applications in May or June before disease occurs. Apply every 14 days.  For southern lawns, start applications at first sign of disease in spring or fall.
Dichondra rust on dichondra lawns	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft [38.4 ounce] 3,840 sq ft	Apply every 21 days.
Dollar spot	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft [38.4 ounce] 3,840 sq ft	Apply every 28 days.
Fusarium patch	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft [38.4 ounce] 3,840 sq ft	Apply every 28 days from fall to early spring.
Gray leaf spot	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft [38.4 ounce] 3,840 sq ft	Apply every 14 days in the summer.
Leaf spot (melting out)	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft [38.4 ounce] 3,840 sq ft	Apply every 14 days.
Pink patch	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft [38.4 ounce] 3,840 sq ft	Apply every 21 days
Powdery mildew	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft [38.4 ounce] 3,840 sq ft	Apply every 21-28 days.
Red thread	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft [38.4 ounce] 3,840 sq ft	Apply every 21 days.
Rust	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft [38.4 ounce] 3,840 sq ft	Apply every 21-28 days.
Snow mold	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft [38.4 ounce] 3,840 sq ft	Apply twice in the fall prior to snow cover waiting 14 days between each application.
Stripe smut	[ <i>pint</i> ] 1,600 sq ft [ <i>quart</i> ] 3,200 sq ft [38.4 ounce] 3,840 sq ft.	Apply in fall after grass is dormant or in early spring before grass starts to grow.
Summer patch	[ <i>pint</i> ] 800 sq ft [ <i>quart</i> ] 1,600 sq ft [38.4 ounce] 3,840 sq ft	Start applications in late May or early June and apply every 14 days up to 6 times.

<b>DISEASES CONTROLLED</b>	<b>AREA TO COVER</b> [Using one quart (32 oz) ] [Using one pint (16 oz)] [38.4 ounces] treats	<b>COMMENTS</b>
Take-all patch	[pint] 800 sq ft [quart] 1,600 sq ft [38.4 ounce] 3,840 sq ft	Apply twice in the fall (September and October) and twice in the spring (April and May).  Southern lawns also need to be treated twice in the summer (June-August).

### **ADDITIONAL USE - GROWTH STIMULATOR** *[optional illustration]*

In addition to its fungicide uses, *[product name]* can be used as a growth stimulator for faster establishment of newly planted **cool season** lawns.

#### *[alternative text]*

In addition to preventing and [&] stopping major diseases on roses, lawns, trees, shrubs and [&] flowers, *[product name]* promotes faster establishment of newly planted **cool-season** lawns.

### **Area to Spray for Newly Planted Cool Season Lawns**

Amount to Use:            One pint (entire contents) treats [covers] 1,600 sq ft.  
                                     One quart (entire contents) treats [covers] 3,200 sq ft.  
                                     38.4 oz. treats up to 3,840 sq ft.

1. Follow “How to use the Ready-Spray System instructions from page [\_\_].
2. Spray evenly over [1,600] [3,200] [3,840] sq ft.

### **When to Apply**

Lawn Seedlings: Apply at the 2- to 3-leaf stage of growth for faster root development and top growth.

### **ORNAMENTAL PLANTS [ROSES, FLOWERS, SHRUBS AND [&] SHADE TREES**

*[optional illustration]*

When used as directed, this product is safe to plants and controls the diseases listed. This product may injure African violets, begonias, Boston fern, geraniums or other unlisted plants, and it may not control unlisted diseases.

### **How to Apply**

- **Follow “How to use Ready-Spray System instructions from page [\_\_].**
- Apply product to the point just when it begins to run off the leaves.

#### *[alternative text]*

- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces, flowers, stems and branches) including new growth.

- Allow spray to completely dry before watering.

### When to Apply

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified (in COMMENTS), treat at the first sign of disease and reapply every 21 days until plant shows signs of recovery. Resume applications if disease reoccurs.

PLANTS	DISEASES CONTROLLED	COMMENTS
Amelanchier	Powdery mildew	
Apple (non-bearing)	Scab	Start applications when leaves start growing in the spring.
Apple (non-bearing)	Rust	Start applications when buds just begin to break open. Make 3 applications.
Apple (non-bearing)	Powdery mildew	Start applications in early spring.
Ash	Powdery mildew	
Azalea	Monilinia flower blight	Apply when there is 5% bloom and again at 70% bloom.
Calendula	Powdery mildew	
Carnation	Rust	
Cherry (ornamental)	Powdery mildew, leaf spot and [&]	Start applications in early spring.
Chrysanthemum	Ray Blight	
Crab apple	Scab	Start applications when buds just begin to break open.
	Rust	Start applications when buds just begin to break open. Make 3 applications.
	Powdery mildew	
Crape myrtle	Powdery mildew	
Delphinium	Powdery mildew	
Douglas Fir, Shasta Fir	Needle Rust	For Shasta Fir, apply in May-June before disease occurs.
Dogwood	Anthraxnose	
Gomphrena	Alternaria leaf spot	
Hawthorn	Rust	
Iris	Rust	
Juniper	Phomopsis blight	
Lilac	Powdery mildew	
Marigold	Alternaria leaf spot	
Monarda	Powdery mildew	
Oak	Leaf Spot	
Phlox	Powdery mildew	
Poplar	Rust	
Red tip photinia	Leaf spot	
Pine	Tip blight	Start applications in mid-late winter (November-January) and apply every 2 months up to 3 times.
Pyracantha	Blight	

PLANTS	DISEASES CONTROLLED	COMMENTS
Rhododendron	Flower blight	Apply when there is 5% bloom and again at 70% bloom.
Roses	Powdery mildew	[Thorough coverage is needed for best results.] Start applications when buds just begin to break open.
Roses	Black spot and [&} rust	<i>[optional text]</i> For severe black spot on roses alternate every 7 to 10 days between this product and Rose Pride Funginex Rose & Shrub Disease Control.  <i>[optional text]</i> For severe black spot on roses use Rose Pride Funginex Rose & Shrub Disease Control.
Snapdragons	Rust	
Sweet William	Leaf spot	
Zinnia	Powdery mildew	

## BEARING FRUIT AND [&} NUT TREES]

[optional illustration]

When used as directed, this product is safe to plants and controls the diseases listed.

### How to Apply

- Apply 4 tsp (1 1/3 ounces) of product to 250 to 500 square feet of bearing fruit and nut trees. Apply spray mixture to the point just when it begins to run off the leaves.

*[alternative text]*

- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces, flowers, stems and branches) including new growth.
- Allow spray to completely dry before watering.

### When to Apply

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified (in COMMENTS), treat at the first sign of disease and reapply every 21 days until plant shows signs of recovery. Resume applications if disease reoccurs.

<b>BEARING FRUIT AND NUT TREES</b>	<b>DISEASES CONTROLLED</b>	<b>COMMENTS</b>
Apricots, Cherries, Nectarines, Peaches, Plums and Prunes	Brown Rot Blossom Blight, Powdery Mildew, Cherry Leaf Spot and Fruit Brown Rot	Start applications in early spring and spray every 21 days, making no more than 5 applications for the entire season. Up to 3 applications may be made from early bloom through petal fall. Up to 2 applications may be made from the end of petal fall through harvest.

## **SWEET CORN**

*[optional illustration]*

When used as directed, this product is safe to plants and controls the diseases listed.

### **How to Apply**

- Apply 4 tsp (1 1/3 ounces) of product to 250 to 500 square feet of sweet corn.

*[alternative text]*

- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces).
- Allow spray to completely dry before watering.

### **When to Apply**

Apply when disease first appears and continue every 14 days, making no more than 4 applications. Do not apply to sweet corn within 14 days of harvest.

<b>SWEET CORN</b>	<b>DISEASES CONTROLLED</b>	<b>COMMENTS</b>
Sweet Corn	Leaf Blights, Rust, Gray Leaf Spot and Eye Spot	Apply when disease first appears and continue every 14 days, making no more than 4 applications. Do not apply to sweet corn within 14 days of harvest.

## [ALT Package 4: Pump-Up Pressure Sprayer]

---

### PRODUCT INFORMATION

---

[Treats] [Covers] up to [1,600 sq ft] [*pint*] [3,200 sq ft]. [*quart*].

Product Facts	
<b>STOPS &amp; PREVENTS DISEASES</b>	Dollar spot, rust, brown patch, powdery mildew, blights, scab, leaf spot and more!
<b>WHERE TO USE</b>	On lawns. Can also be used on roses, flowers, trees, shrubs, and other ornamental plants
<b>AMOUNT TO USE</b>	[16 fl oz treats up to 1,600 sq ft.] [32 fl oz treats up to 3,200 sq ft.]
[image of telephone]	<b>Questions, Comments, or Medical Information call 1-800-888-8372.</b>

Specially formulated for residential use

---

### MIXING PROCEDURES

---

Shake well before using. Apply only with a pump-up pressure sprayer, concentrate plus water-style, or a hose-end sprayer.

**NOTE:**      ½ fl oz= 1 Tablespoon  
                 1 fl oz= 2 Tablespoons

Food utensils, such as measuring cups and spoons, must not be used for food purposes after use in measuring pesticides.

### Pump-Up Pressure Sprayer

1. Fill with required amount of water.
2. Add the amount of concentrate specified in the charts that follow.
3. Shake well before application.

---

## APPLICATION INSTRUCTIONS

---

**STOP. DO NOT ALLOW PEOPLE OR PETS TO CONTACT TREATED AREAS UNTIL SPRAY HAS DRIED.**

### ROSES AND FLOWERS

#### Use Directions

1. Apply product to the point just when it begins to run off of the leaves.
2. Allow spray to completely dry before watering.

Plant	Disease	Amount of Product	Comments
Roses	Powdery Mildew	1 Tablespoon (1/2 fl oz) per gallon of water	Thorough coverage needed for best results.  Begin application at bud break.  Treat every 10 days.
	Black Spot Rust	2 Tablespoons (1 fl oz) per gallon of water	Treat every 10 days.  <b>Note:</b> May not provide 100% control of black spot under severe conditions.
Calendula	Powdery Mildew	1 Tablespoon (1/2 fl oz) per gallon of water	Apply every 30 days.
Carnation, Iris	Rust	1 Tablespoon (1/2 fl oz) per gallon of water	Apply every 10-14 days.
Chrysanthemums	Ray Blight	2 Tablespoons (1 fl oz) per gallon of water	Apply every 7 days.
Delphinium	Powdery Mildew	1 Tablespoon (1/2 fl oz) per gallon of water	Apply every 14 days.
Marigold	Leaf blight, Leaf spots	2 Tablespoons (1 fl oz) per gallon of water	Apply every 14 days.
Phlox, Zinnia	Powdery Mildew	1 Tablespoon (1/2 fl oz) per gallon of water	Apply every 10-14 days.

Plant	Disease	Amount of Product	Comments
		oz) per gallon of water	
Snapdragons	Rust	2 Tablespoons (1 fl oz) per gallon of water	Apply every 10 days.
Sweet William	Leaf Spot	1 Tablespoon (1/2 fl oz) per gallon of water	Apply every 14 days.

### LAWNS *[optional illustration]*

#### Use Directions

1. **When to Mow:** Apply after mowing **or** allow sprayed area to completely dry before mowing.
2. Measure the length and width of area to be treated before application. Calculate square feet by multiplying length times width. For example, an area 25 ft long and 20 ft wide is 500 sq ft.
3. **How to Apply:** Spray evenly over measured area. Treat entire lawn.

#### Established Lawns

Disease	Amount of Product	Comments
Brown Patch	5 fl oz in 1 gallon of water per 250 sq. ft. of lawn	If needed, repeat treatment every 10 days.  First application should be in May or June before disease is present.
Dollar Spot, Powdery Mildew, Rust, Anthracnose	3 fl oz in 1 gallon of water per 250 sq ft of lawn	Apply every 14 days.
Fusarium Patch	5 fl oz in 1 gallon of water per 250 sq ft of lawn	If needed, repeat every 21 days when conditions are favorable for disease to occur.
Necrotic Ring Spot	9 fl oz in 1 gallon of water per 250 sq ft of lawn	Apply in the fall and/or the early spring.
Red Thread, Pink Patch, Dichondra Rust on Dichondra Lawns	5 fl oz in 1 gallon of water per 250 sq ft of lawn	If needed, repeat every 21 days when conditions are favorable for disease to occur.
Snowmold (Gray or Pink)	5 fl oz in 1 gallon of water per 250 sq ft of lawn	Apply twice in the fall prior to snow cover waiting 14 days between each application.
Spring Dead Spot	9 fl oz in 1 gallon of water per 250 sq ft of lawn	Apply when conditions are favorable for disease development.
Stripe Smut	3 fl oz in 1 gallon of water per 250 sq ft of lawn	Apply in Fall after grass is dormant or in early Spring before



Disease	Amount of Product	Comments
		grass starts to grow.
Take-All Patch	5 fl oz in 1 gallon of water per 250 sq ft of lawn	Apply twice in the Fall (Sept. and Oct.) and twice in the spring (April and May).  Southern lawns also need to be treated twice in the summer (June - August).
Zoysia Patch	5 fl oz in 1 gallon of water per 250 sq ft of lawn	Make one application in the early fall prior to development of disease symptoms.

### New Seedings

Amount of Product	Comments
5 fl oz in 2 gallons of water per 500 sq ft of lawn	Apply at the 2 to 3-leaf stage of growth for faster root development and top growth.

### TREES AND SHRUBS

#### Use Directions

1. Spray just until product runs off leaves.
2. Allow the spray to dry before watering plants.

Plant	Disease	Amount of Product	Comments
Ash	Powdery Mildew	3 Tablespoons (1 1/2 fl oz) per gallon of water	Apply every 21 days.
Azalea	Flower Blight	1/2 Tablespoon (1/4 fl oz) per gallon of water	Apply when there is a 5% bloom and again at 70% bloom.
Crab Apple	Scab Rust	1/2 Tablespoon (1/4 fl oz) per gallon of water	Apply when buds just begin to break open. For scab, apply every 14 days. For rust, make 3 applications, beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1 1/2 Tablespoons (3/4 fl oz) per gallon of water	Apply every 14 days.
Crape Myrtle	Powdery Mildew	3 Tablespoons (1 1/2 fl oz) per gallon of water	Apply every 21 days.

<b>Plant</b>	<b>Disease</b>	<b>Amount of Product</b>	<b>Comments</b>
Dogwood	Anthraco nose	1 Tablespoon (1/2 fl oz) per gallon of water	Apply every 14-28 days.
Douglas Fir Shasta Fir	Needle Rust	3 Tablespoons (1 1/2 fl oz) per gallon of water	For Douglas Fir, apply every 21 days. For Shasta Fir, apply in May - June before disease occurs.
Hawthorn, Poplars	Rust	1/2 Tablespoon (1/4 fl oz) per gallon of water	Apply every 14 to 21 days.
Jeffrey Pine	Tip Blight	3 Tablespoons (1 1/2 fl oz) per gallon of water	Make the first application in mid-late winter (Nov.- Jan).  Apply every 2 months up to 3 times.
Juniper	Blight	1 Tablespoon (1/2 fl oz) per gallon of water	Apply every 14 to 21 days.
Lilac	Powdery Mildew	3 Tablespoons (1 1/2 fl oz) per gallon of water	Apply every 21 days.
Oak Red Tip Photinia	Leaf Spot	3 Tablespoons (1 1/2 fl oz) per gallon of water	Apply every 14 days.
Pines	Conifer Blights	3 Tablespoons (1 1/2 fl oz) per gallon of water	Apply every 14 to 21 days.
Pyracantha	Leaf Spot	1/2 Tablespoon (1/4 fl oz) per gallon of water	Apply every 14 to 21 days.
Rhododendron	Flower Blight	1/2 Tablespoon (1/4 fl oz) per gallon of water	Apply when there is a 5% bloom and again at 70% bloom.

## NON-BEARING FRUIT AND NUT TREES

### Use Directions

1. Non-bearing fruit and nut trees are those trees that will not produce fruit for at least one year after application of this product.
2. Apply product to the point just when it begins to run off of the leaves.

Plant	Disease	Amount of Product	Comments
Apple	Scab	1/2 Tablespoon (1/4 fl oz) per gallon of water	Start applications when leaves start growing in the Spring and spray every 14 days.
	Rust	1/2 Tablespoon (1/4 fl oz) per gallon of water	Make 3 applications beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1 Tablespoon (1/2 fl oz) per gallon of water	Start applications in early Spring and spray every 14 days.
Citrus	Greasy Spot	4 Tablespoons (2 fl oz) per gallon of water	Apply June-August every 21 days.
Pecan	Scab Powdery Mildew Downy Spot Zonate Leafspot Fungal Leaf Scorch Brown Spot	2 Tablespoons (1 fl oz) per gallon of water	Start applications at bud break and make 3 applications at 14-day intervals.
Walnut	Anthraxnose	2 Tablespoons (1 fl oz) per gallon of water	Apply every 14 days.

## BEARING FRUIT AND [&] NUT TREES]

[optional illustration]

When used as directed, this product is safe to plants and controls the diseases listed.

### How to Apply

- Apply 4 tsp (1 1/3 ounce) of product to 250 to 500 square feet of bearing fruit and nut trees. Apply spray mixture to the point just when it begins to run off the leaves.

*[alternative text]*

- Spray thoroughly to cover all plant surfaces (upper and lower leaf surfaces, flowers, stems and branches) including new growth.
- Allow spray to completely dry before watering.

### When to Apply

This product is highly effective when it is used at the first sign of disease. Unless otherwise specified (in COMMENTS), treat at the first sign of disease and reapply every 21 days until plant shows signs of recovery. Resume applications if disease reoccurs.

<b>BEARING FRUIT AND NUT TREES</b>	<b>DISEASES CONTROLLED</b>	<b>AMOUNT OF PRODUCT</b>	<b>COMMENTS</b>
Apricots, Cherries, Nectarines, Peaches, Plums and Prunes	Brown Rot Blossom blight, Powdery Mildew, Cherry Leaf Spot and Fruit Brown Rot	4 Teaspoons (2/3 fl oz) per gallon of water applied to 250 to 500 square feet	Start applications in early Spring and spray every 21 days, making no more than 5 applications for the entire season. Up to 3 applications may be made from early bloom through petal fall. Up to 2 applications may be made from the end of petal fall through harvest.

<b>SWEET CORN</b>	<b>DISEASES CONTROLLED</b>	<b>AMOUNT OF PRODUCT</b>	<b>COMMENTS</b>
Sweet Corn	Leaf blights, Rust, Gray Leaf Spot and Eye Spot	4 Teaspoons (2/3 fl oz) per gallon of water applied to 250 to 500 square feet	Apply when disease first appears and continue every 14 days, making no more than 4 applications. Do not apply to sweet corn within 14 days of harvest.

---

### **IMPORTANT** *[optional illustration]*

---

*PPZ 1.55% Multi-Purpose Fungicide* is a preventative fungicide. Applications should be made before diseases occur for best results.

#### *[alternative text]*

This product does not repair [No fungicide repairs] plants or lawns damaged by diseases. It *[product name]* stops [the] disease(s) and prevents reinfestation of treated areas. As treated plants and lawns grow they recover from disease.

If you need more information about lawn and ornamental diseases, contact your local County Extension Service.

[reentry icon] People and pets may enter treated area after spray has dried.

[Master label text continued]

[The Syngenta [alternate brand] Guarantee

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.]

[phone icon] Medical Information call 1-800-888-8372

Product Information call \_\_\_\_\_

For non-emergency (e.g., current product information) call  
Syngenta Crop Protection at 1-800-334-9481.

©200X Syngenta

Banner® and the Syngenta logo are trademarks of a Syngenta Group Company

[Ready-Spray® [and Ortho Rose Pride Funginex®] [is a trademark] [are trademarks] of  
The Scotts Company]

Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, NC 27419-8300  
www.syngenta-us.com

PPZ 1.55 MULTI-PUR FUNG 952 MAS 0210 DCI AMEND MAY2012-HI – sw – 5/18/12  
000100-00952.20120521.PPZ\_1.55%\_MULTI-PUR\_FUNG-DCI\_AMEND-MAY2012-HI.PDF

PPZ 1.55 MULTI-PUR FUNG 952 MAS 0210 DCI AMEND MAY2012-Version B-Hilite – pl – 6/18/15  
000100-00952.20120521B.PPZ\_1.55%\_MULTI-PUR\_FUNG-DCI\_AMEND-MAY2012-HI.pdf

PPZ 1.55 MULTI-PUR FUNG 952 MAS 0210 DCI AMEND MAY2012-Version B-Clean – pl – 6/24/15  
000100-00952.20120521B.PPZ\_1.55%\_MULTI-PUR\_FUNG-DCI\_AMEND-MAY2012.pdf