100 - 973

1/15/2003

pe 129

HELIX™

Insecticide with Fungicides

A seed treatment product for control of certain insects and diseases of canola

For use in commercial seed treatment facilities with closed transfer systems only. No open transfer.

Active Ingredients:	
Thiamethoxam (CAS No. 153719-23-4)	10.30%
Difenoconazole (CAS No. 119446-68-3)	1.24%
(R)-[(2,6-dimethylphenyl)-methoxyacetylamino]-	
propionic acid methyl ester (CAS No. 70630-17-0)	0.39%
Related Compounds	
Fludioxonil (CAS No. 131341-86-1)	
Other Ingredients:	87.93
Total:	100.00%

Helix contains the following amounts of active ingredient per gallon: 1.12 lbs. thiamethoxam; 0.13 lb. difenoconazole; 0.04 lb. (R)-[(2,6-dimethylphenyl)methoxyacetylamino]-propionic acid methyl ester; 0.01 lb. fludioxonil.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See storage, disposal, and additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-973

EPA Est. 71478-CAN-001

Product of Canada

SCP 973MAS (Draft)

U.S. Standard Measure

ACCI	÷,	P	ΤE	D
JAN	I	5	2003	
Under the Fe Fungicide, an as amonded, Registered un EPA Reg. No.	ot.	r tb	e peeuci	oe (

. . . .

.

. . . . - -

• • • •

.

. . .

12 - 10 ⁻¹	FIRST AID
lf inhaled	 Move the person to fresh air. If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center, doctor, or 1-800-888-8372 immediately for treatment advice.
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center, doctor, or 1-800-888-8372 immediately for treatment advice
lf on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center, doctor, or 1-800-888-8372 immediately for treatment advice.
If swallowed	 Call a poison control center, doctor, or 1-800-888-8372 immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
-	Note to Physician ific antidote if Helix is ingested. Induce emesis or lavage symptomatically.
Have the produc or doctor, or goir	t container or label with you when calling a poison control center ng for treatment.
For 24 H	HOT LINE NUMBER our Medical Emergency Assistance (Human or Animal) or al Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372

•

.

239

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if inhaled or absorbed through the skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment

Workers involved in treating, cleanup, or maintenance of seed treatment equipment must wear:

- Chemical-resistant coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material Category A (e.g., barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton).
- Head gear
- Respirator protection (i.e., half-mask respirator with suitable dust filter or fresh air hood).

Baggers, sewers, and stackers must wear:

- Regular coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material Category A (e.g., barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton).
- Head gear
- Respirator protection (i.e., single use dust masks, quarter or half-mask respirator, fresh air hood).

Forklift operators must wear regular coveralls over long-sleeved shirt and long pants and gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

This product is toxic to wildlife and highly toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

• Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff (See manual at the following internet address:

http://www.wcc.nrcs.usda.gov/water/quality/common/pestmgt/core4.html). Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

• Spray Drift Advisory

Do not allow this product to drift.

Physical and Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

4 7 9

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on agricultural establishments in hopper-box, planter-box, slurry-box, or other seed-treatment applications at or immediately before planting.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

Pesticide Storage and Disposal

Store in a cool, dry place. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Container Disposal

Triple rinse (or equivalent), then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

GENERAL INFORMATION

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngentz,

579

no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

Helix is a seed treatment product that contains both insecticide and fungicide components with activity against flea beetles and certain seed-borne and damping-off diseases of canola. Thiamethoxam, the insecticide component, provides activity against flea beetles, and the remaining components are fungicides with activity against seed-borne blackleg (*Leptosphaeria maculans*), seed-borne Alternaria, and the seedling disease complex (damping-off, seedling blight, seed rot, and root rot) caused by *Pythium* spp., *Fusarium* spp., and *Rhizoctonia* spp. Thiamethoxam, difenoconazole, (R)-[(2,6-dimethylphenyl)-methoxyacetylamino]-propionic acid methyl ester, and fludioxonil are the active ingredients in Cruiser™ 5FS, Dividend®, Apron XL® LS, and Maxim® 4FS, respectively.

Note: (R)-[(2,6-dimethylphenyl)-methoxyacetylamino]-propionic acid methyl ester is a systemic fungicide having a specific mode of action. (R)-[(2,6-dimethylphenyl)-methoxyacetylamino]-propionic acid methyl ester could be subject to development of insensitive strains of fungi. Development of insensitivity cannot be predicted. Therefore, Syngenta cannot assume liability for crop damage resulting from insensitive strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible insensitive strains of fungi which may occur.

MIXING PROCEDURES

Helix is especially formulated as a ready-to-apply seed treatment. An EPAapproved coloring agent has been added to the formulation. No additional colorant, dyes, binders, or water are needed. Apply utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect or disease control. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment. Allow seed to dry before bagging.

Federal law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with thiamethoxam insecticide and fludioxonil, difenoconazole, and (R)-[(2,6-dimethylphenyl)methoxyacetylamino]-propionic acid methyl ester fungicides. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs."

CANOLA

To provide early season protection against flea beetles, seed-borne blackleg (*Leptosphaeria maculans*), seed-borne Alternaria, and the seedling disease complex (damping-off, seedling blight, seed rot, and root rot) caused by *Pythium* spp., *Fusarium* spp., and *Rhizoctonia* spp., apply Helix at 23 fl. ozs. of product per 100 lbs. of seed.

Rotational Crops: Treated areas may be replanted immediately following harvest or as soon as practical following the last application with canola. All other crops may be planted 30 days after the last application of Helix.

Apron XL®, Cruiser™, Dividend®, Helix™, Maxim® and the Syngenta logo are trademarks of a Syngenta Group Company U.S. Patent Nos. 5,852,012 and 6,022,871

©2002 Syngenta

For non-emergency (e.g. current product information) call Syngenta Crop Protection at 1-800-334-9481.

Product of Canada

Syngenta Crop Protection, Inc. Seed Treatment Products Greensboro, North Carolina 27409 www.syngenta-us.com

SCP 973A-L1 (Draft)

i

HEL 973MAS-DRAFT 1 CLEAN - Ig- 9-24-02

87

939

<u>е</u> ссіл

r c

CHRON

...

1

9/02 – Revised Environmental Hazards, PPE glove statement; revised container disposal; added Physical and Chemical Hazards; made minor formatting changes.

*