

100-974

05/17/2001

1/20



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number: 100-974
Date of Issuance: MAY 17, 2001

Term of Issuance: Conditional

Name of Pesticide Product: Platinum™ Ridomil Gold®

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300
Attn: Frederick J. Pearson

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(C), provided that you submit the following data and label revisions:

1. Submit and/or cite all data required for registration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Before you release the product for shipment revise the EPA Registration number to read "EPA Reg. No. 100-974".
3. Submit the following information to satisfy the data gaps in the data base for Thiamethoxam and its metabolite.

Residue Chemistry:

- 860.1340: Modification of Method AG-675 to add microwave extraction of liver or a new enforcement method for liver; additional radiovalidation data for beef liver;
- 860.1380: Storage stability data continued to cover the conditions and intervals that plant and animal commodity samples were stored (up to 30 months);

Signature of Approving Official:
151
Tina E. Levine, Ph.D., Chief
Insecticide-Rodenticide Branch
Registration Division 7505C

Date:
MAY 17 2001

EPA form 8570-6 (EPA reg. No. 100-974) continued from page 1...

- 860.1480: Reanalysis of the cow liver **samples** from the feeding study using a microwave extraction step;
- 860.1900: Reanalysis of field rotational **crop samples** for metabolite CGA-265307 and description of storage conditions for future uses;
- 860.1300: Plant metabolism study on a **leafy vegetable** (Crop Group 4 or 5)
- 860.1300: Poultry metabolism study.

Toxicology Data:

870.6300: Developmental neurotoxicity study.

Ecological Effects Data:

72-4b: Daphnia chronic study repeated to fulfill guideline requirements.

Water Monitoring Studies

Two Small-Scale Prospective Ground Water (PGW) studies done according to Guideline 166.1, Draft Small-Scale Prospect Groundwater Guidance Document. This includes the current PGW study in Michigan and one other PGW study in a second State. The registrant will propose several potential sites, located at the 95 percentile of vulnerability and warm climate, for this second study. EPA will approve the final selection. The second site must be operational by May, 2002. The registrant should shall consult with EPA on the degradates to be included in the study. The PGW studies will continue until terminated with agreement by the USEPA.

Monitoring of community water systems using ground water as their source water and private wells. During the summer, 2001, EPA and the registrant will discuss the design of this monitoring program with interested states. Preliminary discussions have begun with Washington, California, Texas, Idaho, Michigan, Florida, North Carolina, and New York. It is anticipated that other states will want to join these discussion. The registrant will submit a protocol for the monitoring studies by December 31. 2001.

- 4. Submit two (2) copies of your **final printed label** for before your release the product for shipment. Please refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

3/20

3

A stamped copy of the label is enclosed for your records. If you have any questions regarding this action, please contact Dani Daniel at 703 305-5409.

Platinum™ Ridomil Gold®

INSECTICIDE/FUNGICIDE

For control of certain insect pests and diseases of potatoes

FOR USE IN POTATOES ONLY

Active Ingredient:

Thiamethoxam (CAS No. 153719-23-4):	4.5%
(R)-[(2,6-dimethylphenyl)- methoxyacetylamino]- propionic acid methyl ester* (CAS No: 70630-17-0)	9.0%
Other Ingredients	86.5%
Total	100.0%

*Also known as mefenoxam.

Platinum™ Ridomil Gold® is a soluble liquid containing approximately 0.406 pounds of thiamethoxam active ingredient per gallon and 0.812 pounds of mefenoxam active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING

See additional precautionary statements and directions for use on container label.

SCP

EPA Reg. 100-OTU

EPA Est.

U.S. Standard Measure

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

MAY 17 2001

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

100-974

FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have a person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
<p style="text-align: center;">HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call 1-800-888-8372</p>	
<p style="text-align: center;">NOTE TO PHYSICIAN There is no specific antidote if Platinum Ridomil Gold is ingested. Induce emesis or lavage stomach. Treat symptomatically.</p>	

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling. Remove clothing and wash before reuse.

Personal Protection Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to inter-tidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff (See manual at the following internet address: <http://www.nrcs.usda.gov/water/quality/common/pestmgmt/files/core4.html>).

Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

Spray Drift Advisory

Do not allow this product to drift.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, POOR DISEASE CONTROL AND/OR ILLEGAL RESIDUES.

Storage and Disposal

Storage

Store in a cool dry place.

Pesticide Disposal

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

Container Disposal

Do not reuse empty container. Triple rinse container and offer for recycling or reconditioning, or puncture and dispose of using one of the following methods: 1) depositing in a sanitary landfill, 2) incineration, or, 3) burning, if allowed by state and local ordinances. Stay out of smoke from burning containers.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup, procedures and disposal of wastes. In the event of a major spill, fire, or other emergency call 1-800-888-8372, day or night.

General Information

Platinum™ Ridomil Gold® controls certain sucking and chewing insects of potatoes, including Colorado potato beetle, potato leafhopper, potato psyllid, flea beetle, and several aphid species. It also controls storage and tuber rots of potatoes, including Pythium leak caused by *Pythium* spp., pink rot caused by *Phytophthora erythroseptica*, and Pythium seedling disease caused by *Pythium* spp.

Platinum Ridomil Gold is a systemic product that is rapidly absorbed by plant roots when properly applied using one of the methods described in the "Recommended Applications of Platinum Ridomil Gold" section of this label. It moves readily throughout the growing plant to protect potato foliage from insect damage. It also moves systemically into developing tubers to protect potatoes against certain soil borne diseases.

Platinum Ridomil Gold is a selective pesticide which should not adversely impact beneficial arthropods and is compatible with integrated pest management programs.

Platinum Ridomil Gold has a wide margin of plant safety and will not adversely impact potato yields when used in accordance with this label.

NOTE: Do not use a foliar application of thiamethoxam (Actara) following an in-furrow or soil application of thiamethoxam (Platinum Ridomil Gold).

Resistance Management

Platinum Ridomil Gold is a systemic insecticide/fungicide containing a specific insecticide mode of action and a specific fungicide mode of action and could be subject to development of resistant strains of either insects or fungi. Development of resistance cannot be predicted. Syngenta cannot assume liability for crop damage resulting from resistant strains of insects or fungi. Insect pests and disease pathogens are known to develop resistance to products that are used repeatedly. The use of this product should conform to sound resistance management strategies established for the crop and use area. Syngenta encourages responsible product stewardship to insure effective long-term control of the insects and diseases on this label.

If treatment is not effective following the use of Platinum Ridomil Gold as recommended, a resistant strain of insect or fungus may be present. In this case, use of other products with different modes of action should be considered. Consult with your state extension service for guidance on your particular insect and disease control situation.

Mixing and Application Procedures

DO NOT APPLY PLATINUM RIDOMIL GOLD AS A BROADCAST FOLIAR SPRAY.

DO NOT APPLY PLATINUM RIDOMIL GOLD THROUGH ANY TYPE OF IRRIGATION SYSTEM.

Mixing Procedures

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area. Keep product container tightly closed when not in use.

Platinum Ridomil Gold Alone: Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the Platinum Ridomil Gold to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the Platinum Ridomil Gold has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Platinum Ridomil Gold + Tank Mixtures: If using Platinum Ridomil Gold in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix product label. No label dosage rate should be exceeded, and the most restrictive label precautions and limitations should be followed. This product should not be mixed with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. In general tank mix partners should be added in this order: products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables, such as Platinum Ridomil Gold, liquids and emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide

sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

Note: When using Platinum Ridomil Gold in tank mixtures, all products in water-soluble packaging should be added to the tank before any other tank mix partner, including Platinum Ridomil Gold. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

Platinum Ridomil Gold + Liquid Fertilizer: Platinum Ridomil Gold may be mixed with water or liquid fertilizer. Spray solutions of Platinum Ridomil Gold should be applied to the field immediately. Do not allow the spray mixture to remain in the spray tank overnight.

Compatibility: Platinum Ridomil Gold is compatible with most commonly soil-applied pesticides. However, since it is not possible to test all possible mixtures, the user should pre-test to insure the physical compatibility and lack of phytotoxic effect of any proposed mixtures with Platinum Ridomil Gold. To determine the physical compatibility of Platinum Ridomil Gold with other products, use a jar test, as described below.

Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

The crop safety of all potential tank mixes on all crops has not been tested. Before applying any tank mixture not specifically recommended on this label, the safety to the target crop should be confirmed.

Application Procedures

Soil Application

Spray nozzles or metering orifices should provide accurate and uniform spray deposition. Platinum Ridomil Gold may be applied as an in-furrow banded spray at-planting, or as a banded spray during a row-mark (mark-out) procedure prior to planting. To help insure accuracy, calibrate sprayer before each use. Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective pest control. For information on spray equipment and calibration, consult sprayer manufacturers and / or State Extension Service specialists.

Impregnated Fertilizer Application

Platinum Ridomil Gold may be impregnated on fertilizer and applied in-furrow at planting or as a banded application during a row mark (mark-out) procedure prior to planting.

Dry Bulk Granular Fertilizer: Many dry bulk fertilizers may be impregnated or coated with Platinum Ridomil Gold and used to control insects and disease in potatoes. When applying Platinum Ridomil Gold with dry bulk granular fertilizer, follow all directions for use and precautions regarding rates per acre, and pests controlled found in the "Recommended Applications of Platinum Ridomil Gold" section of this label.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the individual and or/company selling the pesticide/fertilizer mixture.

Prepare the pesticide/fertilizer mixture by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray Platinum Ridomil Gold onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray onto the fertilizer only, avoiding the walls of the blender. Do not pour or dribble Platinum Ridomil Gold directly from the product container onto the fertilizer.

Blender Mixing Directions: Fill the blender with the required amount of dry bulk granular fertilizer to be used. Start the blender. Spray the Platinum Ridomil Gold directly onto the moving fertilizer. Allow the mixture to blend for at least 5 minutes or until uniform. Spread the pesticide/fertilizer mixture as soon as possible.

Calculate the amounts of Platinum Ridomil Gold by the following formula:

$$\frac{\text{oz./acre Platinum Ridomil Gold}}{\text{lbs./acre Fertilizer}} \times 2,000 \text{ lbs./ton.} = \text{oz./ton Platinum Ridomil Gold}$$

Important: When more than 1000 lbs./A of dry bulk granular fertilizer is to be applied, mix Platinum Ridomil Gold with water to improve coverage on the dry fertilizer. Mix one part Platinum Ridomil Gold with up to 2 parts water [1:2] in a mix tank before application to fertilizer. Use a maximum of 2 qts. liquid / ton of fertilizer. To achieve best results, apply a minimum of 150 lbs./A of dry impregnated fertilizer per acre.

Precautions: Do not impregnate Platinum Ridomil Gold on raw agricultural limestone, since adsorption will not be achieved. Limestone prills which contain a binding agent, and fertilizer blends containing limestone can be impregnated.

Application: For best results, apply the mixture uniformly to the soil with properly calibrated equipment as soon as practical after blending. Non-uniform application of the pesticide/fertilizer mixture may result in unsatisfactory pest control.

Recommended Applications of Platinum Ridomil Gold

CROP	PEST	RATE PER 1000 FT. ROW	REMARKS
POTATO	<u>Insects¹:</u> Colorado potato beetle Aphids Potato leafhopper Potato psyllid Flea beetles <u>Pythium seedling disease:</u> (<i>Pythium</i> spp.) <u>Storage and tuber diseases:</u> Pink Rot (<i>Phytophthora erythroseptica</i> , Pythium Leak (<i>Pythium</i> spp.)	<p align="center">2.2 fl. oz/1000 ft row</p> <p align="center">(See rate conversion chart below for rates per acre)</p>	<p align="center">Apply to the soil using one of the methods described below</p>

Application methods:

- 1) Apply specified rate as a 6-8" banded spray directly over or under the seed pieces in the furrow during the planting operation, or as a 6-8" banded spray during the bedding or row-mark (mark-out) procedure prior to planting. Use a minimum of 3 gallons/A of water carrier per acre for banded sprays.
- 2) Impregnate on dry fertilizer and incorporate within the planted hill during the planting operation, or apply impregnated fertilizer during the bedding or row-mark procedure (mark-out) prior to planting. See the directions for fertilizer impregnation contained in the "Mixing and Application Procedures" section of this label.

Restrictions:

- (1) Make only one soil application per season and do not exceed a total of 38 fluid oz Platinum Ridomil Gold (0.121 lbs. thiamethoxam) per acre during each growing season.
- (2) Do not apply more than one foliar application of a Ridomil Gold prepack (Ridomil Gold MZ, Ridomil Gold Bravo, Ridomil Gold Copper) following a Platinum Ridomil Gold application at planting. A foliar application of a Ridomil Gold prepack at tuber initiation may be needed in addition to the soil application of Platinum Ridomil Gold: a) when conditions are conducive for disease development, b) when potato variety is susceptible or moderately susceptible (pink rot/Pythium leak), or, c) in areas with a long growing season.

¹ If additional insect control measures are needed during the season, it is recommended that products with a different mode of action be used.

Platinum Ridomil Gold Conversion Chart

Fluid ounces per 1000 row feet	Row Spacing	Fluid ounces per acre	Lbs. active thiamethoxam per acre	Lbs. active mefenoxam per acre
2.2 oz per 1000 row ft	30	38	0.121	0.242
	32	36	0.114	0.228
	34	34	0.108	0.216
	36	32	0.102	0.204
	38	30	0.095	0.190

Rotational (Plantback) Restrictions

Areas treated with Platinum Ridomil Gold may be replanted to potatoes or any other crop immediately following harvest or crop loss as long as the plantback intervals in the following table are observed. The plantback interval is defined as the period from the time of application (at planting in this case) to the planting date of the subsequent rotational crop. Any cover crop planted for erosion control or soil improvement may be planted at any time following application if the cover crop is not grazed or harvested for food or feed.

Rotation Crop	Plantback Interval
Cover Crops Not Intended for food or feed	0 days
Cole Crops	
Cotton	
Cucurbit Vegetables	
Tuberous and Corm Vegetables	
Leafy Vegetables	
Fruiting Vegetables	
Tobacco	
Wheat, Barley	
Oats, Rye, Peanuts, Asparagus	120 days
Cranberries, Raspberries	
Root and Tuber vegetables (other than tuberous and corm)	
Alfalfa, Clover, Hops	
Legume Vegetables (beans and peas-succulent and dried), Soybeans	
Grasses*	270 days
Corn	
All other crops Intended For food or feed	365 days

* Any grass in the Gramineae family (either green or cured), except for cereal grains, sugarcane, or any enclosed pasture grasses or grasses grown for hay or silage such as bermudagrass, bluegrass, brome grass, or fescue.

16/20

Platinum™ Ridomil Gold® trademark of Syngenta
U.S. Patent No. 6,022,871

©2001 Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP

Back Page of Booklet

Platinum™ Ridomil Gold®

INSECTICIDE/FUNGICIDE

For control of certain insect pests and diseases of potatoes

FOR USE IN POTATOES ONLY

Active Ingredient:

Thiamethoxam (CAS No. 153719-23-4):	4.5%
(R)-[(2,6-dimethylphenyl)- methoxyacetylamino]- propionic acid methyl ester* (CAS No: 70630-17-0).....	9.0%
Other Ingredients	86.5%
Total.....	100.0%

*Also known as mefenoxam.

Platinum Ridomil Gold is a soluble liquid containing approximately 0.406 pounds of thiamethoxam active ingredient per gallon and 0.812 pounds of mefenoxam active ingredient per gallon.

U. S. Standard Measure

KEEP OUT OF REACH OF CHILDREN

WARNING

See directions for use in attached booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes substantial but temporary eye injury. **Harmful** if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling. Remove clothing and wash before reuse.

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<p>HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call 1-800-888-8372</p>	
<p>NOTE TO PHYSICIAN There is no specific antidote if Platinum Ridomil Gold is ingested. Induce emesis or lavage stomach. Treat symptomatically.</p>	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to inter-tidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff (See manual at the following internet address:

<http://www.nrcs.usda.gov/water/quality/common/pestmgt/files/core4.html>).

Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

Spray Drift Advisory

Do not allow this product to drift.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

For minor spills, leaks, etc., follow precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

EPA Reg. No. 100-OTU

EPA Est.

Platinum™ Ridomil Gold® trademark of Syngenta

©2001 Syngenta

Syngenta Crop Protection, Inc.

Greensboro, North Carolina 27409
www.syngenta-us.com

SCP

(Typed by Fred Pearson)

[Z/LABELP-W/PLATINUM RIDOMIL GOLD] - ccg - 5/11/01