

State Prior to Use of This Product. Product of India Cypermethrin¹: $(\pm)\alpha$ -cyano-(3-phenoxyphenyl) methyl(±)-cis, trans-3-(2,2-dichloroethenyl)-2,2-60.0% 176272 100.0% 1 pound Net Weight

*Cis/trans ratio: 45/55±10

'Synthetic pyrethroid

Active Ingredient:

Other Ingredients:

Total:


Formulated in the USA

SCP 990A-L1B 0804

syngenta

10/12/2004

8



	(Pyrethroid Insecticide) • If pers		 Move person to fresh air. If person is not breathing, call 911 or an
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. 		 ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
	Call a poison control center or doctor for treatment advice.	Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
lf on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of soap and water for 15-20 minutes. Call a poison control center or doctor for treatment advice.	HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident, Call 1-800-888-8372	
If swallowed	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 	Hazards to H Causes subst on clothing, safety glass through the may cause all	HONARY STATEMENTS humans and Domestic Animals WARNING/AVISO antial but temporary eye injury. Do not get in eyes Wear protective eyewear (goggles, face shield, es). Harmful if swallowed, inhaled or absorb skin. Prolonged or frequently repeated skin cont lergic reactions in some individuals. Wash thoroug d water after handling. Remove contaminated clo

.

3

+

2)|0

PRECAUTIONARY STATEMENTS (continued)

Environmental Mazards This product is extremely book to fish and equatic invertebrates. Do not apply directly to water or to an example a sub-present or to intertitial areas below the mean high water mark. Do not contaminate water when desting equipment or disposing of equipment water water. Apply this product only as spectfied on this label, Care should be used when making applications to avoid household pets, particularly fish and reptile pets.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Syngenta Crop Protection, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Syngenta and Seller harmless for any claims relating to such factors.

4

Syngenta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Syngenta, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WAR-RANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICU-LAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

IT IS SYNGENTA'S AND SELLER'S INTENTION THAT in no event shall Syngenta or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR. AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Syngenta and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Syngenta.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Prohibitions

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container.

Storage

Mix only as needed. Do not store diluted material. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, sweep up and remove to chemical waste storage area until proper disposal can be made.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Completely empty jar into application equipment. Triple rinse (or equivalent); then puncture and offer for recycling or dispose of empty jar in a sanitary landfill or by other procedures allowed by state and local authorities.

GENERAL INFORMATION

For crack and crevice or spot treatment for residual and contact pest control in, on, or around buildings and structures and their immediate surroundings and on modes of transport. Permitted areas of use include, but are not limited to, industrial buildings, houses, apartment buildings, laboratories, buses, greenhouses and the nonfood/feed areas of stores, warehouses, vessels, railcars, trucks, trailers, aircraft (cargo and non-cabin areas only), schools, nursing homes, hospitals (non-patient areas), restaurants, hotels and food manufacturing, processing and servicing establishments.

Authorized by the USDA for use in inedible product areas in Federally inspected meat, poultry and egg processing plants.

Demon WP Insecticide is intended for dilution with water for spray application. Do not use strainers finer than 50 mesh size. Fill applicator tank with the desired volume of water and add Demon WP Insecticide. Close and agitate before use in order to insure proper suspension of the wettable powder. Shake or reagitate application equipment before use if application is interrupted. Make up only amount of treatment volume as required. Do not store dilution in application equipment overnight. Repeat treatments as necessary, but not more than once every two weeks.

APPLICATION PROCEDURES

Recommended Application Rates:

Pests	Concentration of Active Ingredient	Dilution Rate
Ants Aphids ³ Assassin Bugs ³	0.1%	1 scoop²/ 1 gal. water
Boxelder Bugs California Oakworm ³		
Centipedes Chiggers ^{3,4}		
Chinch Bugs ³ Cockroaches (Maintenance) Crickets		
Earwigs Elm Leaf Beetle ³	$\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{i=1}^{n-1}\sum_{j=1}^{n-1}\sum_{$	
Firebrats Fleas		
Flies ¹ Millipedes Mites ³		
Mosquitoes Pilibugs		
Silverfish Sowbugs		
Termites ⁵	· ·	

·~+ -

Pests	Concentration of Active ingredient	Dilution Rate
Ants ³ Bees Carpenter Ants Carpenter Bees Cockroaches (Clean-out) Fire Ants (Mounds) ³ Scorpions Sod webworms ³ Spiders Ticks Wasps	0.2%	2 scoops ² / 1 gal. water

¹Not for indoor use in California for fly control. ²1 scoop = 0.33 oz. of Demon WP (equivalent to 2 level tablespoons). ³Not approved for use in California. ⁴Outdoor use only. ⁵Above ground only.

Cockroaches, Spiders, Crickets, Scorpions, Silverfish, Ticks, and Firebrats

Apply as a coarse, low pressure treatment to areas where these pests hide, such as baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, cabinets, sinks, furnaces, stoves, behind and under refrigerators, the underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices; also see Outdoor Surfaces Use.

Ants

Apply to any trails, around doors and windows and similar areas where ants (including carpenter ants) may be found. Apply perimeter barrier treatments to prevent infestation as described below; also see Outdoor Surfaces Use.

Bees, Carpenter Bees, and Wasps

Application to nests should be made in late evening when insects are at rest. Thoroughly treat nest and entrance, surfaces, and surrounding areas where these insects alight also see Outdoor Surfaces Use.

Assassin Bugs, Boxelder Bugs, Sowbugs, Pillbugs, Millipedes, Elm Leaf Beetle, and Centipedes

Apply around doors and windows and similar areas where these, or similar perimeter, pests may be found or where they may enter premises. Apply to baseboards, storage areas and other locations. Apply barrier treatments to prevent infestation as described below; also see Outdoor Surfaces Use.

Bees, Flies, Mosquitoes, and Wasps

Apply directly to walls, ceilings, window screens and other resting areas as a residual surface treatment. May be used inside residential buildings as well as in and around carports, garages and storage sheds; also see Outdoor Surfaces Use.

Termites Above Ground

For control of termites, subterranean aerial colonies, Formosan aerial colonies or drywood termites in localized areas of infested wood in structures, apply a 0.1% emulsion to voids and galleries in damaged wood and in spaces between wooden members of a structure and between wood and foundations where wood is vulnerable. Application may be made to inaccessible areas by drilling and then injecting the emulsion with a crack and crevice injector into the damaged wood or void spaces. Application to attics, crawl spaces, unfinished basements, or man-made voids may be made with a coarse fan spray of 0.1% emulsion to control workers and winged reproductive forms of termites in mud shelter tubes. This type of application is not intended to be a substitute for soil treatment for extensive infestation of drywood termites or other wood-infesting insects. This treatment is intended to kill localized infestations of workers and winged reproductive forms of termites.

For termites active inside trees, utility poles and/or fence posts, drill to find the interior infested cavity and inject 0.1% emulsion using treatment tool with a splash back guard.

Termite carton nests in trees or building voids may be injected with 0.25-0.50% emulsion using a pointed injection tool. Multiple injection points to varying depths may be necessary. It is desirable to physically remove carton nest material from building voids when such nests are found. For enhanced penetration of unfinished wood or other porous materials, utilize the emulsifiable concentrate version of Demon (Demon EC or Demon TC).

Food Handling Establishments (For Use in Non-Food Areas Only)

Demon WP insecticide may be used as a general spot or crack and crevice treatment in the non-food areas of food handling establishments.

Examples of non-food areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Do not use Demon WP Insecticide in food areas of food handling establishments. Food areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) also would be considered a food area.

Residential Food Areas

In the home all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Outdoor Surfaces Use

For control of ants, aphids, assassin bugs, boxelder bugs, bees, centipedes, chiggers, chinch bugs, cockroaches, crickets, earwigs, firebrats, fleas, filies, millipedes, mites, mosquitoes, scorpions, silverfish, sowbugs, pillbugs, spiders, ticks, wasps and other similar perimeter insect pests. Apply with either hand or power application equipment as a residual spray to surfaces of buildings, porches, screens, window frames, eaves, patios, lawns, refuse dumps, garages and other similar areas where these pests are found. Also, may be used as a spot treatment to lawns in landscaped areas in and around residential, commercial, institutional and industrial buildings and similar areas where these insect pests are active.

Fire Ants (Mounds)

For individual fire ant mounds, apply Demon WP as a drench. Gently sprinkle (watering can) a total of 1 gal. over the surface of mounds up to 12 inches in diameter and 2 gals. for larger mounds. Thoroughly wet mound and surrounding area. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration. If possible, applications should be made on warm days after recent rainfall. All colonies in the vicinity should be treated, including those that have not yet constructed a mound.

Scorpions

Treat and remove accumulations of lumber, firewood and other materials which serve as harborage sites. Apply as a coarse spray treatment to the point of drip.

Carpenter Ants

Apply as a coarse wet spray treatment, but avoid runoff around doors, door frames, windows and similar areas where these ants enter premises or hide.

Sod Webworms

8

Apply as a spot treatment when pests first appear (do not apply

as a broadcast spray). Retreat when needed, but not more than once every two weeks. Delay watering or mowing the treated area for 12-24 hours after treatment. Use 1-5 gal. of solution per 800-1,600 sq. ft. when making application.

Barrier Treatment

To help prevent infestation of buildings, apply to a band of soil and vegetation 6-10 ft. wide around and adjacent to the building. Also, treat the building foundation to a height of 2-3 ft. where pests are active and may find entrance. Apply as a coarse spray treatment to thoroughly and uniformly wet the band area, using 1-5 gal. of treatment solution applied to 800-1600 sq. ft.

GENERAL RESTRICTIONS AND PRECAUTIONS

Do not apply this product in or on conduits, motor housings, junction boxes, switch boxes, or other live electrical equipment because of possible shock hazard. Let treated surfaces/areas dry before allowing humans and pets to contact treated surfaces/areas. For best results, thoroughly wash out sprayer and screen with water and detergent before using Demon WP Insecticide. Do not use this product with oil. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

Demon WP will not stain or cause damage to any painted or varnished surface, plastic, fabric or other surface when water applied alone causes no damage.

DO NOT APPLY THIS PRODUCT TO EDIBLE CROPS.

During any application to ceilings of a structure, cover surface below with plastic shielding or similar material.

Do not use as a space spray.

Use only in well ventilated areas,

Do not apply in any rooms when patients, the elderly, or the infirm are actively using the room for sleeping, convalescing, eating or other similar activities.

Do not apply to classrooms when in use.

Do not apply to areas of institutions (including libraries, sports facilities, etc.) when occupants are present in the immediate treatment area.

Do not use in warehouses where raw or cured tobacco is stored.

Do not use in warehouses while raw agricultural commodities for food or feed are being stored.

Do not use in greenhouses on plants grown for food or feed, for sale or other commercial use, or for research purposes.

Do not allow applications to contact water inhabited by fish, such as in aquariums and ornamental fish ponds that are in/around structures being treated.

Do not apply to aircraft cabins.

