



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

August 16, 2010

Mr. Tom Parshley
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419

Subject: Label Notification(s) for Pesticide Registration Notice 2008-1
Updating the Environmental Hazard Language per EPA letter
Dated June 4, 2009

Dear Mr. Parshley:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2008-1 dated May 19, 2010 for the following product(s):

| | |
|---|------------------------------|
| Scimitar WP Insecticide in Water-Soluble Packets | EPA Reg. No. 100-1065 |
| Demand CS Insecticide | EPA Reg. No. 100-1066 |
| Scimitar CS Insecticide | EPA Reg. No. 100-1078 |
| Demand Pestab Insecticide | EPA Reg. No. 100-1082 |
| Scimitar GC Insecticide | EPA Reg. No. 100-1088 |
| Demand EZ Insecticide | EPA Reg. No. 100-1232 |
| Scimitar GR Insecticide | EPA Reg. No. 100-1238 |
| Demand G Insecticide | EPA Reg. No. 100-1240 |

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2008-1 and finds that the label change(s) requested falls within the scope of PRN 98-10. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact Regina Foushee-Smith at 703-605-0780.

Sincerely,

A handwritten signature in black ink that reads "Richard Gebken".

Richard J. Gebken
Product Manager
Insecticide Branch
Registration Division (7505P)

2/18

Please read instructions on reverse before completing form.



United States
Environmental Protection Agency
Washington, DC 20460

| | |
|-------------------------------------|--------------|
| <input type="checkbox"/> | Registration |
| <input type="checkbox"/> | Amendment |
| <input checked="" type="checkbox"/> | Other |

OPP Identifier Number

Notification

Application for Pesticide - Section I

| | | |
|---|---------------------------------------|---|
| 1. Company/Product Number 100-1088 | 2. EPA Product Manager Mark Suarez | 3. Proposed Classification <input type="checkbox"/> None <input checked="" type="checkbox"/> Restricted |
| 4. Company/Product (Name) Scimitar® GC Insecticide | PM# 13 | |
| 5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419 <input type="checkbox"/> Check if this is a new address | | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. NOTIFICATION Product Name AUG 16 2010 |

Section - II

| | |
|--|--|
| <input type="checkbox"/> Amendment - Explain below. | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application. |
| <input checked="" type="checkbox"/> Notification - Explain below. | <input type="checkbox"/> Other - Explain below. |

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)

Notification of label change per June 4, 2009, letter from Environmental Protection Agency. This notification is consistent with the guidance in the June 4, 2009, letter and the requirements of EPA's regulations at 40 CFR part 156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C Sec. 1001 to willfully make any false statements to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR part 156 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

| | | | | | |
|---|--|---|--|--|---|
| 1. Material This Product Will Be Packaged In: | | | | 2. Type of Container | |
| Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | Metal <input type="checkbox"/> | | <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____ |
| *Certification must be submitted | | If "Yes" Unit Packaging wgt. No. per Container | If "Yes" Unit Packaging wgt. No. per container | | |
| 3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container | | 4. Size(s) Retail Container 1 qt. | | 5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product | |
| 6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Other <input type="checkbox"/> Pressure Sensitive <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled | | | | | |

Section - IV

| | | | | | |
|--|--|------------------------------------|---------------------------------------|---|---|
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) | | | | | |
| Name Tom Parshley | | Title Senior Regulatory Manager | | Telephone No. (Include Area Code) 336-632-7207 | |
| Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. | | | | | 6. Date Application Received (Stamped) RECEIVED ON JUN 16 2010 |
| 2. Signature | | | 3. Title Senior Regulatory Manager | | |
| 4. Typed Name Tom Parshley | | | 5. Date May 19, 2010 | | |

Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300
www.syngenta.com



FedEx

May 19, 2010

Document Processing Desk (NOTIF-PYRETHROID)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

**SUBJECT: SCIMITAR® GC INSECTICIDE
EPA REG. NO. 100-1088
.....NOTIFICATION OF LABEL CHANGE PER JUNE 4, 2009, LETTER FROM
ENVIRONMENTAL PROTECTION AGENCY**

Syngenta Crop Protection, Inc. is submitting Notification for Scimitar GC Insecticide, EPA Reg. No. 100-1088. Syngenta is amending the Environmental Hazards and Directions for Use sections of the label by Notification according to the directions stated in the June 4, 2009, letter from EPA.

Attached are:

- One copy of the label with the changes clearly marked
- One unmarked copy of the label
- A CD of the unmarked copy of the label for Electronic Comparison and Review
- Completed EPA Form 8570-1.

Thank you in advance for approving this Notification. If you have any questions, please contact me at 336-632-7207.

Sincerely,

Tom Parshley
Senior Regulatory Manager

Enclosures

AUG 16 2010

4
18

RESTRICTED USE PESTICIDE
Due to Toxicity to Fish and Aquatic Organisms
 FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS
 OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE
 USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

Scimitar® GC

Insecticide

For Use to Control Insect Pests On Golf Course and Athletic Field Turf; Sod Farms; Ornamentals Grown in Commercial Greenhouses, Shadehouses and Nurseries; and on Ornamentals and Lawns in Landscaped Areas Around Residential, Institutional, Public, Commercial and Industrial Buildings, Parks, Recreational Areas, Golf Courses and Athletic Fields

Active Ingredient:

Lambda-cyhalothrin*

[1α(S*),3α(Z)]-(±)-cyano(3-phenoxyphenyl)methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate

9.7%

Other Ingredients:

90.3%

Total

100.0%

*Synthetic pyrethroid.

Contains 0.88 lbs. active ingredient per gallon or 0.22 lbs. a.i. per quart.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1088

EPA Est. No. XXXXX

Product of XXXXX

Formulated in the USA

SCP 1088A-M(Notif Pyrethroid Occupational)

Net contents

| FIRST AID | |
|---|--|
| If on skin or clothing | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| If in eyes | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Do not give any liquid to the person.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person. |
| If inhaled | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. | |
| HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-800-888-8372 | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist or vapors. Harmful if inhaled. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash clothing before reuse.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2 to 30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, Category F, such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, or Viton® \geq 14 mils
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is extremely toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run-off to water bodies or drainage systems. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on this label. Care should be used when making applications to avoid household pets, particularly fish and reptile pets.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, estuaries, and commercial fish farm ponds. In the state of New York, do not apply within 100 ft. of coastal marshes or streams that drain into marshes. When treating tall trees (>15 ft. from the ground) with high pressure sprays or during any application with air assisted equipment (mist blowers), do not apply within 150 ft. of aquatic areas.

Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to the use of this product.

Physical or Chemical Hazards

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

- Do not water treated area to the point of run-off.
- Do not make applications during rain.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - 1 Treatment to soil or vegetation around structures;
 - 2 Applications to lawns, turf, and other vegetation;
 - 3 Applications to building foundations, up to a maximum height of 3 feet.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, Category F, such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, or Viton® \geq 14 mils
- Shoes plus socks

Scimitar GC may be used for applications to ornamentals grown in commercial greenhouses, shadehouses, and nurseries, and turf grown on sod farms or for commercial seed production.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep adults, children, and pets off treated areas until spray has dried following the application.

Scimitar GC may be used for applications to maintain indoor or outdoor areas where turf and ornamentals are grown, such as residential landscaped areas and non-residential landscapes around institutional, public, commercial, and industrial buildings, parks, recreational areas, golf courses, and athletic fields.

Scimitar GC may also be used for applications to golf course fairways, greens, greens aprons, and tee areas.

IMPORTANT: Time application to flowering plants during periods when pollinating insects are not present, such as early morning or late evening.

Do not apply this product through any type of irrigation system.

Do not apply this product to edible crops or crops grown for food/feed.

Do not apply this product by aerial application.

Application

Scimitar GC insecticide mixes readily with water and may be used in all types of application equipment. Mix product with the required amount of water and apply as a dilute application to the point of runoff. Apply product using spray nozzles, which produce a coarse droplet size. Formation of very small droplets may be minimized by appropriate nozzle selection and by avoiding excessive spray pressure. When applying to hard-to-wet foliage, such as holly, pine, or ivy, the addition of a spreader-sticker may enhance knockdown and residual activity. If concentrate or mist-type application equipment is used, use an equivalent amount of product on the area treated as would be used in a dilute application. Phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed. It is advised to pretreat a selection of ornamentals if local use experience is unavailable.

SPRAY DRIFT PRECAUTIONS (sod farms and grass seed farms only)

BUFFER ZONES

Vegetative Buffer Strip

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing Scimitar GC insecticide onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers:

Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21 pp.
www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf

In the State of New York, a 25 ft. vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for ground application.

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast)

Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes; natural ponds; estuaries; and commercial fish ponds).

SPRAY DRIFT REQUIREMENTS (sod farms and grass seed farms only)

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition.

Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

Do not make ground applications into temperature inversions.

Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only medium or coarser spray nozzles (for ground application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

MIXING

Scimitar GC Insecticide is intended for dilution with water for spray application and may be used in all types of application equipment. Fill application tank with $\frac{1}{2}$ – $\frac{3}{4}$ volume of water. Adjust water pH with a buffering agent if necessary to achieve a pH of 5-7. Slowly add Scimitar GC to applicator tank water with maximum agitation. If other chemicals are added to the applicator tank, add Scimitar GC last. If mixed with EC formulations or oils, use within 24 hours. Fill tank to desired volume and continue to agitate while making applications. Shake or re-agitate application equipment before use if application is interrupted. Make up only amount of application volume as required.

**Scimitar GC Mixing Chart for Ornamental Insect Pest Control
(Scimitar GC to add per spray tank)**

| Rate of Scimitar GC | 25 gals. | 50 gals. | 100 gals. | 200 gals. | 300 gals. |
|---------------------|-------------|-------------|-------------|--------------|--------------|
| 1.5 fl. oz. | 0.4 fl. oz. | 0.8 fl. oz. | 1.5 fl. oz. | 3.0 fl. oz. | 4.5 fl. oz. |
| 3.0 fl. oz. | 0.8 fl. oz. | 1.5 fl. oz. | 3.0 fl. oz. | 6.0 fl. oz. | 9.0 fl. oz. |
| 5.0 fl. oz. | 1.3 fl. oz. | 2.5 fl. oz. | 5.0 fl. oz. | 10.0 fl. oz. | 15.0 fl. oz. |

**Scimitar GC Mixing Chart for Turf Insect Pest Control
(Scimitar GC to add per 100 gallon spray tank)**

| Rate of Scimitar GC | 2 gals. | 4 gals. | 6 gals. | 8 gals. | 10 gals. |
|---------------------|--------------|-------------|-------------|-------------|-------------|
| 5 fl. oz./A | 5.7 fl. oz. | 2.9 fl. oz. | 1.9 fl. oz. | 1.4 fl. oz. | 1.2 fl. oz. |
| 10 fl. oz./A | 11.5 fl. oz. | 5.7 fl. oz. | 3.8 fl. oz. | 2.9 fl. oz. | 2.3 fl. oz. |
| | | | | | |

Conversion Rate: 1 fluid ounce (fl. oz.) equals 29 milliliters (mL).

Compatibility

Scimitar GC has been found to be compatible with most commonly used fungicides, miticides, liquid fertilizers, and other insecticides. Check physical compatibility using the correct proportion of products in a small jar, if local experience is unavailable.

Insect Pests Controlled In Ornamentals by Scimitar GC Insecticide

| Use | Pest | Amount of Scimitar GC |
|--|---|---|
| <p>Ornamentals in Greenhouses, Shadehouses, and Nurseries</p> <p>Ornamentals (Including Trees, Shrubs, Flowers, Evergreens, Foliage Plants, and Ground covers) in Residential Landscaped Areas and Landscaped Areas around Institutional, Public, Commercial, and Industrial Buildings, Parks, Recreational Areas, Golf Courses, and Athletic Fields</p> | <p>Ants (Including Imported fire ants)</p> <p>Armyworms</p> <p>Azalea caterpillars</p> <p>Aphids</p> <p>Bagworms</p> <p>Black vine weevils (adult)</p> <p>Boxelder bugs</p> <p>Budworms</p> <p>California oakworms</p> <p>Cankerworms</p> <p>Cockroaches</p> <p>Crickets</p> <p>Cutworms</p> <p>Eastern tent caterpillars</p> <p>Elm leaf beetles</p> <p>European sawflies</p> <p>Fall webworms</p> <p>Flea beetles</p> <p>Forest tent caterpillars</p> <p>Gypsy moth larvae</p> <p>Japanese beetles (adult)</p> <p>June beetles (adult)</p> <p>Lace bugs</p> <p>Leaf-feeding caterpillars</p> <p>Leafhoppers</p> <p>Leafminers (adult)</p> <p>Leaf rollers</p> <p>Leaf skeletonizers</p> <p>Midges</p> <p>Mosquitoes</p> <p>Oleander moth larvae</p> <p>Pillbugs</p> <p>Pine sawflies</p> <p>Pine shoot beetles</p> <p>Pinetip moths</p> <p>Plant bugs</p> <p>Root weevils</p> <p>Sawflies</p> <p>Scale insects (crawlers)</p> <p>Spiders</p> <p>Spittlebugs</p> <p>Striped beetles</p> <p>Striped oakworms</p> <p>Thrips</p> <p>Tip moths</p> <p>Tussock moth larvae</p> <p>Wasps</p> | <p>1.5-5 fl. oz./100 gals. or 44-148 mL/100 gals.</p> |

| Use | Pest | Amount of Scimitar GC |
|--------------------|--|--|
| <i>(continued)</i> | Broadmites Brown softscales California redscales (crawler) Clover mites Mealybugs Pine needlescales (crawler) Spider mites Whiteflies | 3-5 fl. oz./100 gals. or 88-148 mL/100 gals. |

Start application to ornamentals prior to the establishment of high insect pest populations. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Apply at 7-day intervals if treatment is necessary.

Do not apply more than 0.36 lb. of the active ingredient (52.4 fl. oz. of concentrate) per acre per year.

Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at specified rates may enhance the control of insects on certain species of ornamentals having waxy, hard-to-wet foliage.

For spot treatments, use 0.5 fl. oz. Scimitar GC per 1-2.5 gals. of water.

Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.

Note: While phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed, certain cultivars may be sensitive to the final spray solution. Prespray a selection of ornamental plants and observe them for 7-10 days prior to treating large areas if local use experience is unavailable.

Scale

Thoroughly cover the plant with Scimitar GC spray, including trunks, stems, twigs, and foliage for control of scale insects.

Bagworm

Apply Scimitar GC when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young.

Insect Pests Controlled on Turfgrass by Scimitar GC Insecticide

| Use | Pest | Amount of Scimitar GC |
|--|--|--|
| Sod Farms and Grass Seed Farms Lawns around Residential, Institutional, Public, Commercial, and Industrial Buildings, Parks, Recreational Areas, Golf Courses, and Athletic Fields Golf Course and Athletic Field Turf | Ants (Including Imported fire ants) Armyworms Centipedes Crickets Cutworms Earwigs Fleas (adult) Grasshoppers Japanese beetles (adult) Millipedes Mites Pillbugs Sod webworms Sow bugs Ticks (including species which transmit Lyme disease) | 3.4-7 ml/1,000 sq. ft. or 5-10 fl. oz./A |
| | Bluegrass billbugs (adult) Black turfgrass ataenius (adult) Chiggers Fleas (adult) Grub (suppression) Hyperodes weevils (adult) Mole crickets (nymphs and young adults) | 7 mL/1,000 sq. ft. or 10 fl. oz./A |

Start application to turf prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Apply at 7-day intervals if treatment is necessary.

Do not apply more than 0.36 lb. of active ingredient (52.4 fl. oz. of concentrate) per acre per year.

For spot treatments, use 0.5 fl. oz. of Scimitar GC per 1-2.5 gals. of water.

Do not apply when turfgrass is waterlogged or when soils are saturated with water (i.e., will not accept irrigation).

Keep children and pets off treated areas until spray has dried following the application.

Surface Insect Control (armyworm, cutworms, fleas, etc.)

Apply Scimitar GC at specified rates in 2-5 gals. of water per 1,000 sq. ft. The use of a spreader-sticker may be useful if high rainfall amounts are forecast, otherwise the addition of adjuvants is not necessary under normal conditions for surface insect control in turf. Delay watering or mowing for 12-24 hours for optimum control of surface-feeding insect pests.

Thatch Inhabiting Insect Control (chinch bugs, billbugs, etc.)

Apply Scimitar GC at specified rates in 2-10 gals. of water per 1,000 sq. ft. The use of a nonionic wetting agent, penetrant, or similar adjuvant is recommended. Lightly irrigate after application with up to ½ inch of water to move the Scimitar GC into the thatch layer. If irrigation is not available, then use high water application rates for optimum results.

Subsurface Insect Control (mole crickets, grubs, etc.)

Apply Scimitar GC at specified rates in 4-10 gals. of water per 1,000 sq. ft. The use of a nonionic wetting agent, penetrant, or similar adjuvant is strongly recommended. Use the highest water application rates possible with your sprayer. Apply Scimitar GC to turf wet with dew, rain, or irrigation. Water-in immediately after application with ¼–½ inch of water for optimum results.

Fire Ant Control

Treat individual mounds with a drench application using a watering can. Use 0.5 fl. oz. of Scimitar GC per 2.5 gals. of water. Thoroughly soak each mound and a 3 ft. diameter circle around each mound. Gently apply the mixture to avoid disturbing the mound. Disturbing the mound may cause the ants to migrate and reduce the effectiveness of the treatment. For best results, apply in early morning or late evening hours. Make additional treatments as needed, but not more than every 7 days.

Mosquito Control

Apply as a spray around landscape plantings, turf, and building foundations to control mosquitoes. Apply Scimitar GC at specified rates in 2-5 gals. of water per 1,000 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide Storage

Keep container closed when not in use. Do not store near food or feed. Shake well before use. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste storage area until proper disposal can be made.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Completely empty container into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of empty container in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Scimitar® and the Syngenta logo are trademarks of a Syngenta Group Company
©2009 Syngenta

Viton® is a registered trademark of E.I. DuPont de Nemours and Company

| |
|---|
| <p>For non-emergency (e.g., current product information) call Syngenta Crop Protection at 1-800-334-9481.</p> |
|---|

Manufactured for:
Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1088A-M(Notif Pyrethroid Occupational)

SCI GC 1088A-M(Notif Pyrethroid Occupational) CLEAN – pl – 5/5/10
000100-01088.20100519.SCIMITAR-GC_NOTIF_MAY2010.pdf