



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

July 28, 2025

Monique Inforzato  
Monique.Inforzato@syngenta.com  
SYNGENTA CROP PROTECTION, LLC

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment - Revise label and add azoxystrobin registration review  
Product Name: HERITAGE FUNGICIDE  
Admin Number: 100-1093  
EPA Receipt Date: 11/10/2017  
Action Case Number: 00605744

Dear Monique Inforzato:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.


Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have questions, please contact Elizabeth Andrews via email at [andrews.elizabeth@epa.gov](mailto:andrews.elizabeth@epa.gov).

Sincerely,

*Kable Bo Davis*

Kable Bo Davis, Senior Advisor  
FB, RD  
Office of Pesticide Programs



[MASTER]

|                     |              |           |                  |
|---------------------|--------------|-----------|------------------|
| <b>AZOXYSTROBIN</b> | <b>GROUP</b> | <b>11</b> | <b>FUNGICIDE</b> |
|---------------------|--------------|-----------|------------------|

## Heritage® Fungicide

Broad-spectrum fungicide for the prevention and control of diseases of turf and ornamentals, and fruit and nut trees, and vegetable and herb plants for retail sale to consumers.

### Active Ingredient:

Azoxystrobin: methyl (*E*)-2-[2-[6-(2-cyanophenoxy)  
pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate\* ..... 50%

Other Ingredients: ..... 50%

Total: ..... 100%

Heritage Fungicide is a water-dispersible granule (WG) containing 0.5 lb azoxystrobin per lb product.

\*IUPAC

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

See additional precautionary statements and directions for use inside booklet.

Reformulation is prohibited. See individual container labels for repackaging limitations.

EPA Reg No. 100-1093

EPA Est.

---

Net Contents

| <b>FIRST AID</b>   |   |
|--|---|
| <b>If on skin or clothing</b>  | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| <b>If in eyes</b>  | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul> |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment.   |   |
| <b>SYNGENTA HOTLINE NUMBER</b><br>For 24-Hour Medical Emergency Assistance (Human or Animal) or<br>Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),<br>Call<br>1-800-888-8372 |   |

---

## **PRECAUTIONARY STATEMENTS**

---

### **Hazards to Humans and Domestic Animals**

#### **CAUTION**

HARMFUL IF ABSORBED THROUGH SKIN. CAUSES MODERATE EYE IRRITATION. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### **Personal Protective Equipment (PPE)**

#### **Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

In addition,

- **Mixers/loaders** supporting aerial or chemigation applications must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.
- Mixers/loaders supporting groundboom applications must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.
- Mixers/loaders/applicators using mechanically pressurized handwands, except when applying to Christmas tree farms, nursery ornamentals, landscaping, must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters

### **Respirator fit testing, medical qualification, and training**

Using a program that conforms to OSHA's requirements (see 29 CFR Part 1910.134), employers must verify that any handler who uses a respirator is:

- Fit-tested and fit-checked,
- Trained, and
- Examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn. A qualified medical practitioner is a physician or other licensed health care professional who will evaluate the ability of a worker to wear a respirator. The initial evaluation consists of a questionnaire that asks about medical conditions (such as a heart condition) that would be problematic for respirator use. If concerns are identified, then additional evaluations, such as a physical exam, might be necessary. The initial evaluation must be done before respirator use begins. Handlers must be reexamined by a qualified medical practitioner if their health status or respirator style or use-conditions change.

Upon request by local/state/federal/tribal enforcement personnel, employers must provide documentation demonstrating how they have complied with these requirements.

### **User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **Engineering Controls**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS. Human flagging is prohibited.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for “applicators and other handlers” and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

#### **User Safety Recommendations**

##### **Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash thoroughly with soap and water after handling.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **Environmental Hazards**

This pesticide is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

#### **Groundwater Advisory**

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### **Surface Water Advisory**

This product may impact surface water quality due to run-off of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a high potential for reaching surface water via run-off for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features, such as ponds, streams and springs, will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from run-off water and sediment. Run-off of this product also will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify state and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

---

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

---

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.



## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

---

### AGRICULTURAL USES

---

Use to prevent and control diseases of: turf produced on sod farms; and ornamentals, fruit and nut trees, and vegetable and herb plants grown for retail sale to consumers.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

---

## NON-AGRICULTURAL USES

---

Use to prevent and control diseases of turf and ornamentals on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area being treated must be vacated by unprotected persons.

Do not treat areas while unprotected humans or domestic animals are present in the treatment areas. Because certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

Do not allow entry into treatment area until area that was treated with Heritage Fungicide is dry.

---

## PRODUCT INFORMATION

---

Heritage Fungicide is a broad-spectrum, systemic fungicide that prevents or controls the listed pathogens that cause foliar, stem, and root diseases, including leaf and stem blights, leaf spots, patch diseases, mildews, anthracnose, fairy rings, molds, and rusts of turfgrass, ornamentals, vegetables, and herbs. Heritage Fungicide may be applied in alternating application programs or in tank mixes with other registered plant protection products. All applications must be made according to the use directions that follow.

Heritage Fungicide is a member of Syngenta's Plant Performance™ product line that can improve plant vigor and quality. The additional benefits are due to positive effects on plant physiology, which can vary according to plant species and growing environment.

---

## USE RESTRICTIONS

---

**DO NOT** apply more than 10 lb of Heritage Fungicide (5 lb ai) per acre per year to crops grown outdoors.

**DO NOT** apply more than 1.1 lb of Heritage Fungicide (0.55 lb ai) per acre per application.

**DO NOT** graze or feed clippings from treated turf areas to animals.

## PHYTOTOXICITY

Heritage Fungicide is highly phytotoxic to apple and certain crabapple and flowering cherry varieties. Use caution to prevent injury to these trees.

**DO NOT** use spray equipment that has been used to apply Heritage Fungicide to spray apple, crabapple or flowering cherry trees. Even trace amounts can cause unacceptable phytotoxicity in certain varieties.

**DO NOT** apply Heritage Fungicide where spray drift may reach apple, crabapple, or flowering cherry trees.

**DO NOT** apply when weather conditions favor drift from treated areas to a non-target aquatic habitat.

## SPRAY DRIFT MANAGEMENT

### SPRAY DRIFT

#### **Aerial Applications:**

- Do not release spray at a height greater than 10 ft above the ground or crop canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzles that deliver Medium to coarse spray droplets in accordance with ASABE Standard S-572.1.
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Applicators must use  $\frac{1}{2}$  swath displacement upwind at the downwind edge of the field.
- Do not apply during temperature inversions.

#### **Groundboom Applications:**

- User must only apply with the release height recommended by the manufacturer, but no more than 4 ft above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 mph at the application site.
- Do not apply during temperature inversions.

## **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under favorable environmental conditions.

#### **Controlling Droplet Size – Ground Boom**

- **Volume** – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

#### **Controlling Droplet Size – Aircraft**

- **Adjust Nozzles** – Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **BOOM HEIGHT – Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **RELEASE HEIGHT – Aircraft**

Higher release heights increase the potential for spray drift.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be

indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

#### **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

#### **AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**

#### **INTEGRATED PEST (DISEASE) MANAGEMENT (IPM)**

Heritage Fungicide should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development, such as planting of resistant cultivars, removal of plant debris to reduce inoculum, and water management, should be followed. The following sections in this label identify specific IPM recommendations for each crop. Consult your local agricultural, turf and ornamental authorities for additional IPM strategies established for your area. Heritage Fungicide may be used in State Agricultural Extension advisory (disease forecasting) programs that recommend application timing based on environmental factors favorable for disease development.

## RESISTANCE MANAGEMENT

For resistance management, Heritage Fungicide contains a Group 11/[azoxystrobin] fungicide. Any fungal population may contain individuals naturally resistant to Heritage Fungicide and other Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Heritage Fungicide or other Group 11 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local Syngenta Representative, retailer, or extension specialist for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Syngenta at 1-866-Syngent(a) (866-796-4368). You can also contact your pesticide distributor or university extension specialist to report resistance.

Since pathogens differ in their potential to develop fungicide resistance, use the resistance management strategies for each disease given in the **SPECIFIC USE DIRECTIONS** section in this label.

Syngenta Crop Protection encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

## MIXING INSTRUCTIONS

To prepare spray solution, partially fill the spray tank with clean water and begin agitation. Add the specified amount of Heritage Fungicide to the tank, allowing adequate time for good mixing. Add an adjuvant if appropriate. Finish filling the tank to the desired volume to obtain the proper spray concentration. Maintain agitation throughout the spraying operation. **DO NOT** allow spray mixture to stand overnight or

for prolonged periods. Mix only the amount of spray required for your immediate use. Sprayers should be thoroughly cleaned immediately after application.

Heritage Fungicide is compatible with commonly used fungicides, liquid fertilizers, herbicides, insecticides and biological control products. If tank mixes are desired, products should be added to the spray tank in the following order: Heritage Fungicide, other WG or dry flowable formulations, wettable powders and flowable (aqueous suspension) products. Observe all directions, precautions, and limitations on labeling of all products used. Consult compatibility charts or your local or state agricultural or turf authorities for compatibility information.

**Adjuvants:** When an adjuvant is to be used with this product, use an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program.

**DO NOT** mix Heritage Fungicide in the spray tank with pesticides, surfactants or fertilizers unless compatibility charts or your own prior use has shown that the combination is physically compatible, effective and non-injurious under your conditions of use. If physical compatibility is unknown, the following procedure should be followed: Pour the specified proportions of the products into a suitable container of water, mix thoroughly and allow it to stand at least 20 minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible.

Heritage Fungicide is incompatible with many fertilizers when low water volumes are used for in-furrow applications. Cold temperatures and water quality exacerbate these compatibility problems. Conduct a physical compatibility test as described in the paragraph as above before making a field application.

**NOTE:** Heritage Fungicide has demonstrated some phytotoxic effects when mixed with products that are formulated as EC's. These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, silicone-containing adjuvants have also contributed to phytotoxicity.

---

## APPLICATION INSTRUCTIONS

---

Heritage Fungicide should be applied prior to disease development at the rates and timings given in this label. Apply at the higher rates in the rate range and/or shorter spray intervals under conditions of heavy infection pressure, on highly susceptible varieties or when environmental conditions are conducive for disease development. Use of Heritage Fungicide as a "rescue" (late curative or eradicator) treatment is not recommended and may not give satisfactory disease control.

Heritage Fungicide may be applied with various types of spray equipment commonly used for making ground and aerial applications. For ground applications, apply

Heritage Fungicide in a volume of water sufficient to provide good plant canopy penetration and coverage. For aerial applications, apply Heritage Fungicide in a minimum of two gallons of water per acre. Ground application is preferred as it typically provides better canopy penetration and coverage.

Proper adjustments and calibration of spraying equipment are essential for optimal disease control. If you have questions about calibration, contact a State Extension Service specialist, the equipment manufacturer or other experts.

### **Directions for Use through Sprinkler and Drip Chemigation Systems**

**Spray Preparation:** Chemical tank and injector system should be thoroughly cleaned. Flush system with clean water.

### **Use Precautions for Sprinkler and Drip Irrigation Applications:**

**Drip and Micro Irrigation:** Heritage Fungicide may be applied through drip and micro irrigation systems to potted ornamentals or to bedded, field-grown ornamentals for soil-borne disease control. The soil or potting media should have adequate moisture capacity prior to drip application.

Terminate drip irrigation at fungicide depletion from the main feed supply tank or after 6 hours from start, whichever is shorter. For maximum efficacy, subsequent irrigation (water only) should be delayed for at least for 24 hours following drip application.

**Sprinkler Irrigation:** Apply this product through sprinkler irrigation systems including center pivot, motorized boom, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system except as specified on this label.

Apply with center-pivot or continuous-move equipment distributing ½ acre-inch or less during treatment. In general, use the least amount of water required for proper distribution and coverage. If stationary systems (solid set, handlines or wheel lines other than continuous-move) are used, this product should be injected into no more than the last 20-30 minutes of the set. **DO NOT** apply when winds are greater than 10-15 mph to avoid drift or wind skips. Plant injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform treated water. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.

If you have questions about calibration, contact the State Extension Service specialist, equipment manufacturers or other experts.

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.



- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

### **Specific Instructions for Public Water Systems**

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There must be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

---

## DIRECTIONS FOR TURF

---

Heritage Fungicide controls the listed pathogens that cause foliar, stem, and root diseases, including leaf and stem blights, leaf spots, patch diseases, mildew, molds and rusts of turfgrass. Heritage Fungicide may be used on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, athletic fields, and sod farms.

### Integrated Pest (Disease) Management (IPM)

Sound turf management resulting in healthy, vigorous turf is the foundation of a good IPM program. Cultural practices, such as proper choice of turf variety, nutrient management, proper cutting height, thatch management, and proper watering, drainage, and moisture stress management, should be integrated with the use of fungicides to increase turf vigor and reduce the susceptibility to disease. Immunoassay detection kits and extension service diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

### Resistance Management

**DO NOT** spray more than two sequential Heritage Fungicide applications for Gray Leaf Spot or *Pythium* spp. control. For all other diseases when Gray Leaf Spot and *Pythium* spp. are not present, **DO NOT** spray more than three sequential applications of Heritage Fungicide.

### Application Directions

Heritage Fungicide should be applied prior to disease development. Mix Heritage Fungicide with the required amount of water and apply as a dilute spray in 2-4 gallons of water per 1,000 square feet (87-174 gallons per acre). Repeat applications at specified intervals. For spot treatments, use 0.2 oz of Heritage Fungicide per 1-3 gallons of water.

## TURF USE RESTRICTIONS

**DO NOT** apply more than 10 lb of Heritage Fungicide/Acre (5 lb ai/Acre) per year to crops grown outdoors.

For applications with handheld equipment, **DO NOT** exceed 0.0025 lb ai/gal.

For applications with mechanically pressurized handwand equipment to golf course (tees, greens, fairways) or landscaping turf, **DO NOT** exceed 0.95 lb ai/A.

Aerial and/or chemigation application to sod is prohibited

**DO NOT** apply to golf course turf by air.

## Soil Injection Application

Heritage Fungicide may be applied through a liquid fungicide injector for the control of ectotrophic root diseases such as summer patch and take-all patch. Use Heritage Fungicide only in liquid injection equipment specifically designated for pesticide use.

Apply Heritage Fungicide at 0.2 to 0.4 oz per 1,000 sq ft. Spray carrier volume should fall within 30-150 gallons of water per 1,000 sq ft. Use an injection hole spacing of 1 inch by 1 inch for optimum control. Do not exceed injection depth of 2 inches. Use a one-inch depth for optimum results. Application timing should follow disease control strategies used for normal broadcast spray programs.

## Application when Establishing Turfgrass from Seed or in Overseeding of Dormant Turfgrass

Heritage Fungicide may be used for control of certain turfgrass diseases associated with turfgrass establishment from seed or during overseeding of dormant turfgrass. Heritage Fungicide may be safely applied before or after seeding or at seedling germination and emergence to ryegrass, bentgrass, bluegrass (including *Poa trivialis*), and fescue turfgrass types. Optimum application timing is during seeding. See **Application Directions** section.

Dollar Spot: Heritage Fungicide does not control dollar spot. Mix Heritage Fungicide with another fungicide labeled for dollar spot control when environmental conditions are conducive for disease development. Heritage Fungicide is tank-mix compatible with many fungicides that control dollar spot.

**TABLE 1: Directions for Application for Turf Diseases**

| Target Diseases   | Use Rate<br>(oz product/<br>1,000 sq ft) | Application<br>Interval<br>(days) | Application Instructions*   |
|---|--|-----------------------------------|---|
| Anthrachnose<br>( <i>Colletotrichum cereale</i> )                             | 0.2-0.4                                  | 14-28                             | Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development. |
| Take-all root rot<br>Bermudagrass Decline<br>( <i>Gaeumannomyces</i><br>spp.) | 0.4                                      | 28                                | Use preventatively. Begin applications when conditions are favorable for disease infection prior to disease symptom development.  |
| Brown Patch<br>( <i>Rhizoctonia solani</i> )                                  | 0.2-0.4                                  | 14-28                             | Apply when conditions are favorable for disease development.  |

| Target Diseases   | Use Rate<br>(oz product/<br>1,000 sq ft) | Application<br>Interval<br>(days) | Application Instructions*  |
|---|--|-----------------------------------|--|
| Brown Ring Patch<br>( <i>Waitea circinata</i> )   | 0.2-0.4                                  | 14-28                             | Apply when conditions are favorable for disease development  |
| Yellow Patch<br>( <i>Rhizoctonia cerealis</i> )   | 0.2-0.4                                  | 14-28                             | Make one or two applications in fall or when conditions are favorable for disease development.   |
| Fairy Ring<br>( <i>Lycoperdon</i> spp.,<br><i>Agrocybe pediades</i> ,<br><i>Arachnion</i> spp., <i>Bovista</i><br>spp., and <i>Vascellum</i><br>spp.) | 0.4                                      | 28                                | Apply preventatively or as soon as possible after fairy ring symptoms develop. Add the specified rate of a wetting agent to the final spray and water-in immediately with 1/8 to 1/4 inches irrigation. Fairy ring symptoms may take 2 to 3 weeks to disappear following curative application. Reapplication after 28 days may be required in some cases. Severely damaged or thin turf may require reseeding. |
| Microdochium Patch<br>( <i>Monographella nivalis</i> )  | 0.2-0.4                                  | 14-28                             | Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.  |
| Gray Leaf Spot<br>( <i>Pyricularia oryzae</i> )   | 0.2-0.4                                  | 14-28                             | Begin applications before disease is present and continue applications while conditions are favorable for disease development.   |
| Gray Snow Mold<br>Typhula blight<br>( <i>Typhula incarnata</i> , and<br><i>T. ishikariensis</i> )   | 0.4                                      | 10-28                             | Make two applications of 0.4 oz spaced 10-28 days apart in late fall just before snow cover. When disease pressure is high, tank-mix Heritage Fungicide with another fungicide registered for control of snow mold.  |
| Leaf and Sheath Spot<br>( <i>Waitea zea</i> )   | 0.4                                      | 14-28                             | Apply when disease conditions are favorable for disease development.   |
| Leaf Rust, Stem Rust,<br>Stripe Rust<br>( <i>Puccinia</i> spp.)   | 0.2-0.4                                  | 14-28                             | Begin applications when conditions are favorable for disease infection, prior to disease symptom development.  |
| Leaf Spot<br>( <i>Bipolaris</i> spp.)   | 0.2-0.4                                  | 14-21                             | Apply when conditions are favorable for disease development.   |
| Melting Out<br>( <i>Drechslera poae</i> )   | 0.2-0.4                                  | 14-21                             | Apply when conditions are favorable for disease development.   |
| Necrotic Ring Spot<br>( <i>Ophiosphaerella korrae</i> )   | 0.2-0.4                                  | 14-28                             | Apply when conditions are favorable for disease development.   |
| Pink Patch<br>( <i>Limonomyces roseipellis</i> )  | 0.2-0.4                                  | 14-28                             | Apply when conditions are favorable for disease development.   |
| Pink Snow Mold<br>( <i>Monographella nivalis</i> )  | 0.4                                      | 10-28                             | Make two applications of 0.4 oz spaced 10-28 days apart in late fall just before snow cover. When disease pressure is high, tank-mix Heritage Fungicide with another fungicide registered for control of snow mold.  |

| Target Diseases   | Use Rate<br>(oz product/<br>1,000 sq ft) | Application<br>Interval<br>(days) | Application Instructions*  |
|---|--|-----------------------------------|--|
| Powdery Mildew<br>( <i>Erysiphe graminis</i> )  | 0.2-0.4                                  | 14-28                             | Begin applications when conditions are favorable for disease infection, prior to disease symptom development.  |
| Pythium Blight<br>Pythium Root Rot<br>( <i>Pythium</i><br><i>aphanidermatum</i> ,<br><i>Pythium</i> spp.) | 0.2-0.4                                  | 10-14                             | Use preventatively. Begin applications before disease is present. During periods of prolonged favorable conditions, treat on the 10-day application interval. For use on newly seeded as well as established turf. |
| Pythium Root Dysfunction<br>( <i>Pythium volutum</i> )  | 0.4                                      | 21-28                             | Apply preventatively when mean daily soil temperatures are between 55 F and 70 F. Irrigate with 0.1 to 0.2 inches within 24 hours after application to facilitate movement into the root zone.                     |
| Red Thread<br>( <i>Laetisaria fuciformis</i> )  | 0.2-0.4                                  | 14-28                             | Apply when conditions are favorable for disease development.   |
| Large Patch<br>( <i>Rhizoctonia solani</i> )  | 0.2-0.4                                  | 14-28                             | Make one or two applications in fall or when conditions are favorable for disease development. Spring applications may also be required in some locations or when disease pressure is high.                        |
| Southern Blight<br>( <i>Sclerotium rolfsii</i> )  | 0.2-0.4                                  | 14-28                             | Apply when conditions are favorable for disease development.   |
| Summer Patch<br>( <i>Magnaporthiopsis poae</i> )  | 0.2-0.4                                  | 14-28                             | Apply when conditions are favorable for disease development.   |
| Take-all patch<br>( <i>Gaeumannomyces graminis</i> )  | 0.2-0.4                                  | 28                                | Begin applications when conditions are favorable for disease infection, prior to disease symptom development. Make two applications 28 days apart in the spring and two applications 28 days apart in the fall.    |
|   |  |                                   |  |

\*Do not apply more than two sequential applications of Heritage Fungicide for control of Gray Leaf Spot and *Pythium* spp. For all other diseases when Gray Leaf Spot and *Pythium* spp. are not present, do not apply more than three sequential applications of Heritage Fungicide.

**TABLE 2: Heritage Fungicide Rate Conversion Chart for Turf**

| Oz Product/<br>1,000 sq ft | Oz AI/<br>1,000 sq ft | Oz Product/<br>Acre | Pound Product/<br>Acre |
|----------------------------|-----------------------|---------------------|------------------------|
| 0.20                       | 0.10                  | 8.7                 | 0.5                    |
| 0.30                       | 0.15                  | 13.1                | 0.8                    |
| 0.40                       | 0.20                  | 17.4                | 1.1                    |

**TABLE 3: Amount of Heritage Fungicide to Mix 100 Gallons for Turf Applications**

| Heritage Fungicide Use Rate | Spray Volume (gallons/1,000 sq ft) |             |             |
|-----------------------------|------------------------------------|-------------|-------------|
|                             | 2.0 gallons                        | 3.0 gallons | 4.0 gallons |
| 0.20 oz                     | 10 oz                              | 6.7 oz      | 5 oz        |
| 0.40 oz                     | 20 oz                              | 13.3 oz     | 10 oz       |

## **DIRECTIONS FOR ORNAMENTALS**

Heritage Fungicide controls listed pathogens that cause foliar, aerial, and root diseases, including: leaf, tip, and flower blights; leaf spots; downy mildew; powdery mildew; anthracnose; and rusts of ornamental plants. Heritage Fungicide may be used to control certain diseases of container, bench, flat, plug, bed, succulents or field-grown ornamentals in greenhouses, shade- and lath-houses, outdoor nurseries, retail nurseries, interiorscapes, and other landscape areas.

### **Integrated Pest (Disease) Management (IPM)**

Heritage Fungicide should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, winter and/or spring pruning, plant debris management and proper timing and placement of irrigation. Immunoassay detection kits and diagnostic services can assist in the early and accurate identification of causal organisms and corresponding selection of the proper fungicide when required.

### **Resistance Management**

**DO NOT** make more than three (3) sequential applications of Heritage Fungicide before alternating with a fungicide of a different mode of action. A sound resistance management program would include blocks of three Heritage Fungicide applications separated by blocks of two alternate fungicide applications. **DO NOT** alternate Heritage Fungicide with other strobilurin fungicides.

### **Application Directions**

Apply Heritage Fungicide as a foliar or soil broadcast, drench, or banded spray targeted at the foliage or crown of the plant. Apply to the point of runoff to ensure complete coverage of the target plant. Good coverage and wetting of foliage is necessary for best control. Refer to the label for specific use directions for control of certain diseases. Repeat applications at specified intervals using an appropriate resistance management program.

Apply Heritage Fungicide at listed use rates. The addition of an adjuvant at the specified use rate may enhance coverage on hard-to-wet plant foliage. Under light to

moderate disease pressure, use the lower listed rates and shorter listed application interval. Under environmental conditions conducive to severe disease development, use the higher rates and shorter listed application interval. Use of Heritage Fungicide as a "rescue" (late curative or eradicator) treatment is not recommended and may not result in satisfactory disease control.

## **ORNAMENTALS USE RESTRICTIONS**

**DO NOT** apply more than 0.75 lb ai/A for field-grown or nursery ornamentals by groundboom or chemigation.

For applications with handheld equipment to nursery ornamentals, broadcast application or landscaping (plants, flowers, trees and turf), **DO NOT** exceed 0.0025 lb ai/gal.

For application with handheld equipment to Christmas tree, **DO NOT** exceed 0.00125 lb ai/gal.

For application with mechanically pressurized handwand to greenhouse ornamentals, **DO NOT** exceed 0.0025 lb ai/gal.

**DO NOT** apply greater than 2 pints of solution per square foot for drench and crown applications.

**DO NOT** exceed 600 gallons spray volume per acre for foliar applications.

**DO NOT** tank-mix Heritage Fungicide with other fungicides, insecticides, herbicides, fertilizers, or adjuvants unless local experience indicates that the tank mix will not injure ornamental plants.

**DO NOT** apply Heritage Fungicide to apple or cherry trees (including flowering and ornamental varieties, such as Yoshino) due to possible phytotoxicity.

**DO NOT** use spray equipment that has applied Heritage Fungicide for use in these sensitive crops due to possible phytotoxicity from residue remaining in the sprayer.

**DO NOT** apply more than 10 lb of Heritage Fungicide/Acre (5 lb ai/Acre) per year to crops grown outdoors.

## **Drench Application**

Heritage Fungicide may be applied to control soilborne, seedling, and crown diseases of ornamentals as a preventative drench treatment prior to infection. Good coverage of



the pre-infection area (root zone, root ball, and crown) is necessary for satisfactory control. Heritage Fungicide may be applied by drench to container-grown ornamentals. Make a drench application prior to infection as healthy roots are necessary to optimize product uptake and systemic translocation to optimize disease protection. Due to the systemic activity of Heritage Fungicide, suppression of certain foliar diseases has been observed in plants treated with drench applications.

Use caution before applying Heritage Fungicide as a drench to small bedding plants in the seedling/plug stage due to possible phytotoxicity. A limited quantity of plants should be tested prior to full-scale application.

### **Chemigation: Use through Sprinkler and Drip Irrigation Systems**

Heritage Fungicide may be applied through sprinkler, drip, or other micro irrigation systems to potted ornamentals or to bedded, field-grown ornamentals for soil-borne disease control. Apply Heritage Fungicide as a preventative treatment. The soil or potting media should have adequate moisture capacity prior to drip application.

Terminate irrigation at fungicide depletion from the main feed supply tank or after 6 hours from start, whichever is shorter. For maximum efficacy, subsequent irrigation (water only) should be delayed for at least for 24 hours following application.

### **PHYTOTOXICITY**

Application of Heritage Fungicide to crabapple may cause phytotoxicity. Heritage Fungicide may be applied to specific tolerant varieties of crabapple. Heritage Fungicide has been shown to be safer when applied to the species and varieties listed in **Table 12**. Due to the large number of species, and varieties of crabapple, it is impossible to test every one for tolerance to Heritage Fungicide. The user should conduct small-scale testing to ensure plant safety prior to large-scale commercial use on plant species and varieties not listed on this label.

When used in accordance with the label directions, Heritage Fungicide will provide control of the diseases of ornamental plants in the following table.

**TABLE 4: Foliar Plant Diseases Controlled**

(Heritage Fungicide is sold in several product container sizes. For product container sizes of 8 oz and larger, follow the mixing instructions in the second column. For the 4 oz product container size, follow the mixing instructions in the third column in the following table.)

| <b>DISEASE (Pathogen)</b>  | <b>Application Instructions</b>   |   |
|--|---|---|
|  | 8 oz and larger<br>product container sizes<br>(oz product per 100 gallons)  | 4 oz product container size<br>(oz product per 50 gallons)  |
| <b>CONIFER BLIGHTS</b>   |   |   |
| Phomopsis Blight<br>( <i>Phomopsis juniperovora</i> )  | Apply 1-4 oz every 7-28 days.   | Apply 0.5-2 oz every 7-28 days.   |
| Tip Blight<br>( <i>Sirococcus strobilinus</i> )  | Apply 1-4 oz every 7-28 days.   | Apply 0.5-2 oz every 7-28 days.   |
| <b>LEAF BLIGHTS/LEAF SPOTS</b>   |   |   |
| Alternaria Leaf Spot<br>( <i>Alternaria</i> spp.)  | Apply 1-8 oz every 7-28 days.   | Apply 0.5-4 oz every 7-28 days.   |
| Anthrachnose<br>( <i>Colletotrichum</i> spp.,<br><i>Elsinoe</i> spp.)  | Apply 1--8 oz every 7-28 days.  | Apply 0.5-4 oz every 7-28 days.   |
| Cercospora Leaf Spot<br>( <i>Cersospora</i> spp.)  | Apply 1-4 oz every 7-28 days.   | Apply 0.5-2 oz every 7-28 days.   |
| Cylindrocladium leaf<br>spot/stem canker<br>( <i>Cylindrocladium</i> spp.)   | Apply 4-8 oz every 7-14 days.   | Apply 2-4 oz every 7-14 days.   |
| Downy Mildew<br>(including <i>Peronospora</i> spp.,<br><i>Plasmopara</i> spp.,<br><i>Bremiella</i> spp., <i>Bremia</i> spp.) | Apply 2-4 oz every 7-21 days during periods of active plant growth and prior to dormancy or severe infection. Use lower use rates for herbaceous seedlings. | Apply 1-2 oz every 7-21 days during periods of active plant growth and prior to dormancy or severe infection. |
| Entomosporium Leaf Spot<br>( <i>Entomosporium</i> spp.)<br>( <i>Diplocarpon</i> spp.)  | Apply 1-4 oz every 7-28 days.   | Apply 0.5-2 oz every 7-28 days.   |
| Iris Leaf Spot<br>( <i>Mycosphaerella</i> spp.)  | Apply 2-4 oz every 7-21 days.   | Apply 1-2 oz every 7-21 days.   |
| Leaf spot<br>( <i>Cladosporium</i> spp.,<br><i>Septoria</i> spp. and<br><i>Stemphylium</i> spp.)                             | Apply 1-4 oz every 7-28 days.   | Apply 0.5-2 oz every 7-28 days.   |

| DISEASE (Pathogen)   | Application Instructions  |   |
|--|---|---|
|  | 8 oz and larger product container sizes (oz product per 100 gallons)  | 4 oz product container size (oz product per 50 gallons)   |
| Rose Blackspot<br>( <i>Diplocarpon rosea</i> )   | Apply 4-8 oz every 7-14 days. Under severe disease conditions or if disease is already present, apply Heritage Fungicide at the highest listed rate and shortest application interval. Heritage Fungicide may be tank-mixed with another rose blackspot fungicide. Do not exceed 24 oz/A per application. | Apply 2-4 oz every 7-14 days. Under severe disease conditions or if disease is already present, apply Heritage Fungicide at the highest listed rate and shortest application interval. Heritage Fungicide may be tank-mixed with another rose blackspot fungicide. Do not exceed 24 oz/A per application. |
| Myrothecium leaf spot<br>( <i>Myrothecium</i> spp.)  | Apply 2-4 oz every 7-21 days.   | Apply 1-2 oz every 7-21 days.   |
| Scab<br>( <i>Venturia inaequalis</i> ,<br><i>Sphaceloma poinsettiae</i> ,<br><i>Elsinöe australis</i> )  | Apply 1-4 oz every 10-28 days. Do not apply to apple trees. For crabapples, see <b>Table 12</b> for tolerant varieties.   | Apply 0.5-2 oz every 10-28 days. Do not apply to apple trees. For crabapples, see Table 12 for tolerant varieties.  |
| Marssonina Leaf Spot<br>( <i>Marssonina</i> spp.)  | Apply 1-4 oz every 14-28 days.  | Apply 0.5-2 oz every 14-28 days.  |
| <b>POWDERY MILDEW</b>  |   |   |
| <i>Erysiphe</i> spp.,<br><i>Microsphaera</i> spp.,<br><i>Podosphaera</i> spp., <i>Oidium</i> spp.,<br><i>Podosphaera</i> spp.,<br><i>Uncinula</i> spp.   | Apply 1-4 oz every 7-28 days. Do not make more than 2 sequential applications before rotating to another class of fungicide.  | Apply 0.5-2 oz every 7-28 days. Do not make more than 2 sequential applications before rotating to another class of fungicide.  |
| <b>RUSTS</b>   |   |   |
| Needle Rust<br>( <i>Melampsora occidentalis</i> )  | Apply 1-4 oz every 7-28 days.   | Apply 0.5-2 oz every 7-28 days.   |
| Other Rusts<br>( <i>Phragmidium</i> spp.,<br><i>Puccinia</i> spp.,<br><i>Gymnosporangium</i> spp.,<br><i>Coleosporium</i> spp.,<br><i>Uromyces</i> spp.) | Apply 1-4 oz every 7-28 days.   | Apply 0.5-2 oz every 7-28 days.   |
| <b>FLOWER BLIGHTS</b>  |   |   |
| Anthrachnose<br>( <i>Collectotrichum</i> spp.,<br><i>Elsinoe</i> spp.)   | Apply 1-4 oz every 7-28 days.   | Apply 0.5-2 oz every 7-28 days.   |
| Botrytis Blight<br>( <i>Botrytis cinerea</i> )   | Apply 4-8 oz every 7-21 days. For suppression only. Do not exceed 24 oz/acre.   | Apply 2-4 oz every 7-21 days. For suppression only. Do not exceed 24 oz/acre.   |

| DISEASE (Pathogen)   | Application Instructions   |  |
|--|--|--|
|  | 8 oz and larger<br>product container sizes<br>(oz product per 100 gallons) | 4 oz product container size<br>(oz product per 50 gallons) |
| <b>SHOOT/STEM DISEASES</b>   |  |  |
| Aerial/Shoot Blight<br>( <i>Phytophthora</i> spp.)   | Apply 1-4 oz every 7-28 days.  | Apply 0.5-2 oz every 7-28 days.                            |
| <p><b>DO NOT</b> apply more than 0.75 lb ai/A for field-grown or nursery ornamentals by groundboom or chemigation.</p> <p>For applications with handheld equipment to nursery ornamentals, broadcast application or landscaping (plants, flowers, trees and turf), <b>DO NOT</b> exceed 0.0025 lb ai/gal.</p> <p>For application with mechanically pressurized handwand to greenhouse ornamentals, <b>DO NOT</b> exceed 0.0025 lb ai/gal</p> <p><b>DO NOT</b> exceed 600 gallons spray volume per acre for foliar applications.</p> <p><b>DO NOT</b> apply more than 10 lb of Heritage Fungicide/Acre (5 lb ai/Acre) per year to crops grown outdoors.</p> |  |  |

**TABLE 5: Soilborne Diseases Controlled – Directed Spray**

| Pathogen  | Application Instructions   |  |
|---|--|--|
|   | 8 oz and larger<br>product container sizes<br>(oz product per 100 gallons) | 4 oz product container size<br>(oz product per 50 gallons) |
| <i>Fusarium</i> spp.<br><i>Rhizoctonia solani</i><br><i>Athelia rolfsii</i><br>( <i>Sclerotium rolfsii</i> )<br><i>Sclerotinia</i> spp. | Apply 1-4 oz every 7-21 days.  | Apply 0.5-2 oz every 7-21 days.                            |

**TABLE 6: Soilborne Diseases Controlled – Drench and Drip Irrigation**

See **Drench Application** section of **ORNAMENTAL DIRECTIONS FOR USE** for additional drench directions.

| <b>Pathogen</b>  | <b>Use Rate</b><br>(oz product/100 gallons) | <b>Application Instructions</b>   |
|--|---|---|
| <i>Fusarium</i> spp.<br><i>Rhizoctonia solani</i><br><i>Athelia rolfsii</i><br>( <i>Sclerotium rolfsii</i> ) | 0.2-1.2 oz                                  | Apply 1-2 pints of the solution per square foot surface area every 7-28 days.   |
| <i>Sclerotinia</i> spp.  | 1.2 oz                                      | Apply 1-2 pints of the solution per square foot surface area every 7-28 days. Apply for control of <i>Sclerotinia</i> by drench application only. |
| <b>DO NOT</b> apply greater than 2 pints of solution per square foot for drench and crown applications.      |   |   |

**TABLE 7: Soilborne Diseases Controlled – Banded Applications**

| <b>Pathogen</b>   | <b>Use Rate</b><br>(oz product/1,000 row feet) | <b>Application Instructions</b>   |
|---|--|---|
| <i>Fusarium</i> spp.<br><i>Rhizoctonia solani</i><br><i>Athelia rolfsii</i><br>( <i>Sclerotium rolfsii</i> )<br><i>Sclerotinia</i> spp. | 0.2-0.4 <sup>1</sup> oz                        | Apply as a banded spray, 7 inches or less in width, directed to the soil using single or multiple spray nozzles adjusted to provide thorough coverage of lower stems <sup>2</sup> and the soil surface. |

<sup>1</sup>When applications are applied to crops grown using 22-inch row spacing, the maximum application rate is 0.35 oz/1,000 row feet.

<sup>2</sup>Applications that come into contact with the foliage are considered foliar applications for resistance management.

**TABLE 8: Soilborne Diseases Controlled – In-furrow Application**

| <b>Pathogen</b>  | <b>Use Rate</b><br>(oz product/1,000 row feet) | <b>Application Instructions</b>  |
|--|--|--|
| <i>Fusarium</i> spp.<br><i>Rhizoctonia solani</i><br><i>Athelia rolfsii</i><br>( <i>Sclerotium rolfsii</i> ) | 0.2-0.3 oz <sup>1</sup>                        | Mount the spray nozzle to allow the spray mixture to be applied directly into the furrow and before the propagated unit (seed, seed pieces, bulbs or corms) are covered by soil. Apply the higher listed rate when current or expected weather conditions are conducive for disease development. |
| <b>Suppression Only:</b><br><i>Pythium</i> spp.  |  |  |

<sup>1</sup>Apply product in 3 to 15 gallons of water per 1,000 row feet.

**TABLE 9: Soilborne Diseases Controlled – Broadcast Application**

| Pathogen  | Use Rate<br>(oz product/A) | Application Instructions  |
|---|----------------------------|---|
| <i>Fusarium</i> spp.<br><i>Rhizoctonia solani</i><br><i>Athelia rolfsii</i><br>( <i>Sclerotium rolfsii</i> )<br><i>Sclerotinia</i> spp. | 2-16 oz                    | Apply as a preventative broadcast application. Soil or potting media should have adequate moisture capacity prior to application if applied by overhead irrigation. |

**TABLE 10: Foliar and Soil Diseases Suppressed – Drench and Drip Applications**

| Disease/Pathogen  | Use Rate<br>(oz product/100 gallons) | Application Instructions  |
|---|--------------------------------------|---|
| Rusts<br>Powdery Mildew<br><i>Pythium</i> spp.  | 0.45-1.2 oz                          | Apply 1-2 pints of the solution per square foot surface area every 7-28 days. |
| <b>DO NOT</b> apply greater than 2 pints of solution per square foot for drench and crown applications. |                                      |   |

## Plant Safety

Heritage Fungicide has been shown to be safe when applied to the ornamental plants listed in **Tables 11 and 12** when applied according to listed application methods, rates, and timings. Due to the large number of species and varieties of ornamental and nursery plants, it is impossible to test every one for tolerance to Heritage Fungicide. Neither the manufacturer nor the seller has determined whether or not Heritage Fungicide can be used safely on ornamental and nursery plants not specified on this label. The user should conduct small-scale testing to ensure plant safety prior to large-scale commercial use on varieties not listed on this label. Note: Applications to petunia and impatiens in bloom may occasionally cause discoloration to the flowers.

**DO NOT** tank-mix Heritage Fungicide with other fungicides, insecticides, herbicides, fertilizers, or adjuvants unless local experience indicates that the tank mix is safe to ornamental plants.

**DO NOT** apply Heritage Fungicide to certain crabapple or cherry trees (including flowering or ornamental varieties, such as Yoshino) due to possible phytotoxicity (see **Table 13** for complete list). **DO NOT** use spray equipment that has been used to apply Heritage Fungicide for use in these sensitive crops due to possible phytotoxicity.

**TABLE 11: Tolerant Plants Listed by Common Name**

| COMMON NAME                                      | BOTANICAL NAME                                  |
|--|---|
| Abelia   | <i>Abelia</i> spp.                              |
| Andromeda, Japanese                              | <i>Pieris japonica</i>                          |
| Arborvitae                                       | <i>Thuja</i> spp.                               |
| Aspen Trees                                      | <i>Populus</i> spp.                             |
| Aster  | <i>Aster</i> spp.                               |
| Aucuba, Japanese                                 | <i>Aucuba japonica</i>                          |
| Azalea, Glacier                                  | <i>Rhododendron</i> spp.                        |
| Azaleas  | <i>Rhododendron</i> spp.                        |
| Balsam   | <i>Impatiens</i> spp.                           |
| Barberry   | <i>Berberis thunbergii</i>                      |
| Begonia (except Rieger begonia)                  | <i>Begonia</i> spp.                             |
| Birch, River                                     | <i>Betula nigra</i>                             |
| Black-Eyed Susan                                 | <i>Rudbeckia hirta</i>                          |
| Blanket Flower                                   | <i>Gaillardia</i> spp.                          |
| Bougainvillea                                    | <i>Bougainvillea</i> spp.                       |
| Boxwood  | <i>Buxus sempervirens</i>                       |
| Buddleia   | <i>Buddleia davidii</i>                         |
| Bugle  | <i>Ajuga reptans</i>                            |
| Bugleweed  | <i>Ajuga reptans</i>                            |
| Burning Bush                                     | <i>Euonymus alatus</i>                          |
| Butterfly Bush                                   | <i>Buddleia davidii</i>                         |
| Cactus, Holiday                                  | <i>Schlumbergera</i>                            |
| Caladium   | <i>Caladium</i> spp.                            |
| Camellia   | <i>Camellia japonica</i>                        |
| Carnation  | <i>Dianthus caryophyllus</i>                    |
| Ceanothus  | <i>Ceanothus</i> spp.                           |
| Cedar, Atlas                                     | <i>Cedrus atlantica</i>                         |
| Cedar, Red                                       | <i>Juniperus virginiana</i>                     |
| Cedar, Western Red                               | <i>Thuja plicata</i>                            |
| Cedar, White                                     | <i>Cedrus</i> spp.                              |
| Christmas Trees                                  | See Fir, Douglas; Fir, Fraser; and Pine, Scotch |
| Chrysanthemum                                    | <i>Chrysanthemum</i> spp.                       |
| Cinquefoil                                       | <i>Potentilla</i> spp.                          |
| Clethra  | <i>Clethra alnifolia</i>                        |
| Coleus   | <i>Plectranthus</i> spp.                        |
| Cotoneaster, Creeping                            | <i>Cotoneaster adpressus</i>                    |
| Cotoneaster, Variegated Rockspray                | <i>Cotoneaster horizontalis</i>                 |
| Crabapple (See <b>Table 12</b> for variety list) | <i>Malus</i> spp.                               |
| Cranesbill                                       | <i>Geranium</i> spp.                            |

| COMMON NAME                    | BOTANICAL NAME                       |
|--------------------------------|--------------------------------------|
| Crapemyrtle                    | <i>Lagerstroemia indica</i>          |
| Cyclamen                       | <i>Cyclamen</i> spp.                 |
| Cyperus                        | <i>Cyperus</i> spp.                  |
| Cypress, Sawara                | <i>Chamaecyparis pisifera</i>        |
| Cypress, Leyland               | <i>Chamaecyparis</i> spp.            |
| Daisy, Gerber                  | <i>Gerbera jamesonii</i>             |
| Daisy, Transvaal               | <i>Gerbera jamesonii</i>             |
| Dogwood                        | <i>Cornus</i> spp.                   |
| Dogwood                        | <i>Cornus florida</i>                |
| Dogwood, Pink                  | <i>Cornus</i> spp.                   |
| Dumb-Cane                      | <i>Dieffenbachia</i> spp.            |
| Euonymus, Dwarf Winged         | <i>Euonymus alata</i>                |
| Euonymus, Evergreen            | <i>Euonymus japonicus</i>            |
| Evergreen, Chinese             | <i>Aglaonema</i> spp.                |
| Fatsia, Japanese               | <i>Fatsia japonica</i>               |
| Fig                            | <i>Ficus</i> spp.                    |
| Fir, Douglas                   | <i>Pseudotsuga</i> spp.              |
| Fir, Fraser                    | <i>Abies fraseri</i>                 |
| Fir, Noble                     | <i>Abies procera</i>                 |
| Floss Flower                   | <i>Ageratum</i> spp.                 |
| Forsythia                      | <i>Forsythia viridissima</i>         |
| Foxglove                       | <i>Digitalis</i> spp.                |
| Gardenia                       | <i>Gardenia jasminoides</i>          |
| Geranium                       | <i>Pelargonium</i> spp.              |
| Grass                          | <i>Pennisetum alopecuroides</i>      |
| Grass, Dwarf Pampas            | <i>Phalaris</i> spp.                 |
| Grass, Pampas                  | <i>Cortaderia selloana</i>           |
| Hawthorn, Indian               | <i>Rhaphiolepis indica</i>           |
| Heather                        | <i>Erica dareyensis</i>              |
| Hemlock                        | <i>Tsuga</i> spp.                    |
| Hemlock, Western               | <i>Tsuga heterophylla</i>            |
| Hibiscus                       | <i>Hibiscus moscheutos</i>           |
| Hibiscus                       | <i>Hibiscus rosa-sinensis</i>        |
| Holly                          | <i>Ilex</i> spp.                     |
| Hosta                          | <i>Hosta</i> spp.                    |
| House Leek                     | <i>Sempervivum</i> spp.              |
| Hydrangea                      | <i>Hydrangea</i> spp.                |
| Hydrangea, French              | <i>Hydrangea macrophylla</i>         |
| Impatiens <sup>1,2</sup>       | <i>Impatiens</i> spp. <sup>1,2</sup> |
| Iris (Bulbous, Spanish, Dutch) | <i>Iris xiphium</i>                  |
| Iris, African                  | <i>Dietes iridiodes</i>              |



| COMMON NAME          | BOTANICAL NAME                    |
|----------------------|-----------------------------------|
| Iris, Butterfly      | <i>Dietes iridiodes</i>           |
| Ivy, Algerian        | <i>Hedera algeriensis</i>         |
| Ivy, English         | <i>Hedera helix</i>               |
| Ivy, Swedish         | <i>Plectranthus</i> spp.          |
| Japanese Pittosporum | <i>Pittosporum tobira</i>         |
| Juniper              | <i>Juniperus procumbens</i>       |
| Juniper              | <i>Juniperus scopulorum</i>       |
| Juniper              | <i>Juniperus</i> spp.             |
| Larkspur             | <i>Delphinium</i> spp.            |
| Laurel               | <i>Laurus nobilis</i>             |
| Laurel, Australian   | <i>Pittosporum</i> spp.           |
| Laurel, Japanese     | <i>Aucuba japonica</i>            |
| Lilac, California    | <i>Ceanothus</i> spp.             |
| Lilac, Wild          | <i>Ceanothus sanguineus</i>       |
| Lily, Asiatic        | <i>Lilium</i> spp.                |
| Lily, Peace          | <i>Spathiphyllum floribundium</i> |
| Lily-Turf            | <i>Liriope muscari</i>            |
| Live-Forever         | <i>Sempervivum</i> spp.           |
| Magnolia             | <i>Magnolia</i> spp.              |
| Magnolia, Saucer     | <i>Magnolia soulangiana</i>       |
| Magnolia, Southern   | <i>Magnolia grandiflora</i>       |
| Maple, Japanese      | <i>Acer palmatum</i>              |
| Maple, Sugar         | <i>Acer saccharum</i>             |
| Marigold             | <i>Tagetes</i> spp.               |
| Mugwort              | <i>Artemisia</i> spp.             |
| Nandina              | <i>Nandina domestica</i>          |
| Oak, Pin             | <i>Quercus palustris</i>          |
| Oak, Red             | <i>Quercus falcata</i>            |
| Oleander             | <i>Nerium oleander</i>            |
| Orpine               | <i>Sedum</i> spp.                 |
| Palm, Date           | <i>Phoenix dactylifera</i>        |
| Palm, Parlor         | <i>Chamaedora elegans</i>         |
| Palm, Queen          | <i>Syagrus romanzoffianum</i>     |
| Palm, Robellini      | <i>Phoenix roebelenii</i>         |
| Palm, Sago           | <i>Caryota urens</i>              |
| Pansy <sup>1</sup>   | <i>Viola</i> spp. <sup>1</sup>    |
| Paper Plant          | <i>Fatsia japonica</i>            |
| Pear, Bradford's     | <i>Pyrus calleryana</i>           |
| Periwinkle           | <i>Vinca</i> spp.                 |
| Petunia <sup>2</sup> | <i>Petunia</i> spp.               |

| COMMON NAME          | BOTANICAL NAME   |
|----------------------|--|
| Philodendron         | <i>Philodendron</i> spp.   |
| Phlox                | <i>Phlox</i> spp.  |
| Photinia, Red-Tip    | <i>Photinia glabra</i>   |
| Pine                 | <i>Pinus</i> spp.  |
| Pine, Black          | <i>Pinus nigra</i>   |
| Pine, Eastern White  | <i>Pinus strobus</i>   |
| Pine, Mugo           | <i>Pinus mugo</i>  |
| Pine, Scotch         | <i>Pinus sylvestris</i>  |
| Pink                 | <i>Dianthus</i> spp.   |
| Plum, Flowering      | <i>Prunus</i> spp.   |
| Plum, Purple-Leaf    | <i>Prunus</i> spp.   |
| Poinsettia           | <i>Euphorbia</i> spp.  |
| Poplar               | <i>Populus trichocarpa</i>   |
| Pothos               | <i>Epipremnum</i> spp.   |
| Primrose             | <i>Primula</i> spp.  |
| Pussy's Foot         | <i>Ageratum</i> spp.   |
| Redbud, Western      | <i>Cercis occidentalis</i>   |
| Rhododendron         | <i>Rhododendron</i> spp.   |
| Ribbon Grass         | <i>Setaria</i> spp.  |
| Rose of Sharon       | <i>Hibiscus syriacus</i>   |
| Rose                 | <i>Rosa</i> spp.   |
| Rose-Bay             | <i>Nerium oleander</i>   |
| Rosemary (Prostrate) | <i>Rosmarinus</i> spp.   |
| Rubber Plant, Baby   | <i>Peperomia</i> spp.  |
| Rubber Tree          | <i>Brassaia actinophylla</i>   |
| Sage                 | <i>Salvia</i> spp.   |
| Sand cherry          | <i>Prunus pumila</i>   |
| Snap-Dragon          | <i>Antirrhinum</i> spp.  |
| Snowball             | <i>Ceanothus</i> spp.  |
| Spirea               | <i>Spirea budalda</i>  |
| Spirea               | <i>Spirea japonica</i>   |
| Spruce, Blue         | <i>Picea pungens</i>   |
| Spruce, Norway       | <i>Picea abies</i>   |
| Spruce, White        | <i>Picea glauca</i>  |
| Starwort             | <i>Aster</i> spp.  |
| Stonecrop            | <i>Sedum</i> spp.  |
| Succulent            | Succulents encompass a wide variety of genera, including but not limited to Echeveria, Crassula, Haworthia, Gasteria, Agave, Euphorbia, Senecio, Sedum, and Sempervivum. |
| Sweet Alyssum        | <i>Lobularia maritima</i>  |

| COMMON NAME      | BOTANICAL NAME               |
|------------------|------------------------------|
| Thyme, Creeping  | <i>Thymus serpyllum</i>      |
| Umbrella Tree    | <i>Brassaia actinophylla</i> |
| Verbena          | <i>Verbena</i> spp.          |
| Vervain          | <i>Verbena</i> spp.          |
| Viburnum         | <i>Viburnum</i> spp.         |
| Vinca            | <i>Catharanthus roseus</i>   |
| Viola            | <i>Viola</i> spp.            |
| White Alder      | <i>Clethra</i> spp.          |
| Weigela, Pink    | <i>Weigela florida</i>       |
| Willow, Virginia | <i>Itea virginica</i>        |
| Winterberry      | <i>Ilex</i> spp.             |
| Wormwood         | <i>Artemisia</i> spp.        |
| Yaupon           | <i>Ilex</i> spp.             |
| Yew, Spreading   | <i>Taxus baccata</i>         |
| Yucca            | <i>Yucca</i> spp.            |
| Zebra Plant      | <i>Aphelandra</i> spp.       |
| Zinnia           | <i>Zinnia</i> spp.           |

<sup>1</sup> Do not exceed 2 oz/100 gallons on these species.

<sup>2</sup> Heritage Fungicide may occasionally cause discoloration of flowers when applied directly to blooms of certain plant species. Not all varieties and colors have been evaluated.

**TABLE 12: Tolerant Varieties of Ornamental Crabapple Species (Genus *Malus*)**

|                                     |   |                |  |
|-------------------------------------|---|----------------|--|
| Callaway                            | Golden Raindrops                                      | Mary Potter    | Selkirk                                |
| Carmine ( <i>M. atrosanguinea</i> ) | Hopa  | Molten Lava    | Sentinel                               |
| Candymint Sargent                   | Indian Magic  | New Centennial | Silver Moon                            |
| Christmas Holly                     | Island  | Ormiston Roy   | Silverdrift                            |
| David                               | Jackii ( <i>M. baccata</i> var. jackii)               | Pink Satin     | Sinai Fire                             |
| Dolgo                               | Japanese Flowering Crabapple ( <i>M. floribunda</i> ) | Prairie Maid   | Sugar Tyme                             |
| Donald Wyman                        | Katherine   | Prairiefire    | Van Eseltine                           |
| Dorothea                            | Lancelot  | Profusion      | White Angel                            |
| Doubloons                           | Louisa  | Ralph Shay     | Wild crabapple ( <i>M. coronaria</i> ) |
| Eleyi                               | <i>Malus x zumi</i> var. Calocarpa                    | Red Baron      | Winter Gold                            |
| Evereste                            | <i>M. sargentii</i>                                   | Red Jade       |  |
| Eyelynn                             | Manchurian ( <i>M. baccata</i> var. mandshurica)      | Sargent        |  |

**TABLE 13: Plants Sensitive to Heritage Fungicide  
(Do not apply Heritage Fungicide to these species or varieties.)**

| COMMON NAME                                      | BOTANICAL NAME                               |
|--|--|
| Crabapple - Flame variety                        | <i>Malus</i> spp. 'Flame'                    |
| Crabapple – Brandywine variety                   | <i>Malus</i> spp. 'Brandywine'               |
| Crabapple – Novamac variety                      | <i>Malus</i> spp. 'Novamac'                  |
| Cherry, Flowering – Yoshino variety              | <i>Prunus</i> x <i>yedoensis</i>             |
| Leatherleaf Fern and Other Ferns for cut foliage | <i>Rumohra adianformis</i> and other species |
| Privet   | <i>Ligustrum</i> spp.                        |

## CONIFERS INCLUDING CHRISTMAS TREES AND COMMERCIAL PRODUCTION ROSES

Heritage Fungicide may be used to control certain diseases on conifers and commercial production roses in indoor and outdoor production and landscape situations. See the **DIRECTIONS FOR ORNAMENTALS** section above for more detailed directions for use in landscape situations.

[On the following **Specific Use Directions for Conifer and Commercial Rose Production** table, use the **oz product/A** rate (column 3) for products 8 oz or larger. Use the **oz product/0.5 A** rate (column 4) for 4 oz products.]

**TABLE 14: Specific Use Directions for Conifer and Commercial Rose Production**

| Crop  | Target Diseases   | Use Rate<br>oz product/A<br>(lb ai/A) | Use Rate<br>oz product/0.5<br>A<br>(lb ai/A) | Application Instructions  |
|---|---|---------------------------------------|--|---|
| Conifers<br>including<br>Christmas<br>Trees | Diplodia tip blight<br>( <i>Diplodia pinea</i> )            | 3.2-8.0<br>(0.10-0.25)                | 1.6-4.0<br>(0.10-0.25)                       | <p><b>Integrated Pest (Disease) Management:</b> Heritage Fungicide should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance and removal of plant debris in which inoculum may overwinter.</p> <p><b>Resistance Management:</b> Do not apply more than four sequential applications of Heritage Fungicide before alternating with a fungicide that is not in Group 11. Do not make more than eight applications of Heritage Fungicide per acre per year.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout the disease development period at 7- to 21-day intervals following the resistance management guidelines.</p> <p>Applications may be</p> |
|   | Lophodermium needlecast<br>( <i>Lophodermium pinastri</i> ) |                                       |  |   |
|   | Swiss needlecast<br>( <i>Phaeocryptopus gaumannii</i> )     |                                       |  |   |

| Crop | Target Diseases | Use Rate<br>oz product/A<br>(lb ai/A) | Use Rate<br>oz product/0.5<br>A<br>(lb ai/A) | Application Instructions   |
|------|-----------------|---------------------------------------|--|--|
|      |                 |                                       |  | made by ground, air or chemigation. An adjuvant may be added at specified rates. |

**Specific Use Restrictions: DO NOT** apply more than 4.0 pounds product/acre (2.0 lb ai/A/year).

**Maximum application rates for Christmas tree using any handheld equipment: DO NOT** exceed 0.00125 lb ai/gal for foliar applications.

|  |  |                        |                        |   |
|--|--|------------------------|------------------------|---|
| <b>Roses<br/>(commercial<br/>production)</b> | Black spot<br>( <i>Diplocarpon rosea</i> )<br>Cercospora Leaf spot<br>( <i>Cercospora</i> spp.)<br>Downy Mildew<br>( <i>Peronospora sparsa</i> )<br>Powdery Mildew<br>( <i>Podosphaera pannosa</i> )<br>Rust<br>( <i>Phragmidium mucronatum</i> , <i>P. tuberculatum</i> , and other<br><i>Phragmidium</i> spp.)<br>Septoria Leaf Spot<br>( <i>Septoria rosea</i> )<br>Alternaria Leaf Spot<br>( <i>Alternaria alternata</i> ) | 1.6-8.0<br>(0.05-0.25) | 0.8-4.0<br>(0.05-0.25) | <p><b>Integrated Pest (Disease) Management:</b><br/>Heritage Fungicide should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, winter and/or spring pruning, plant residue management and proper timing and placement of irrigation.</p> <p><b>Resistance Management:</b> Do not make more than four sequential applications of Heritage Fungicide before alternating with a fungicide that is not in Group 11. Do not make more than eight applications per acre per year.</p> <p>Begin Heritage Fungicide application prior to disease development and continue throughout the disease development period on 7- to 21-day intervals following the resistance management guidelines.</p> <p>Applications may be made by ground, air or chemigation. An adjuvant may be added at specified</p> |
|--|--|------------------------|------------------------|---|

| Crop | Target Diseases | Use Rate<br>oz product/A<br>(lb ai/A) | Use Rate<br>oz product/0.5<br>A<br>(lb ai/A) | Application Instructions |
|------|-----------------|---------------------------------------|--|--------------------------|
|      |                 |                                       |  | rates.                   |

**Specific Use Restrictions:** Do not apply more than 4.0 pounds product/acre (2.0 lb ai/A/year).

**DO NOT** apply more than 0.75 lb ai/A for field-grown or nursery ornamentals by groundboom or chemigation.

For applications with handheld equipment, **DO NOT** exceed 0.0025 lb ai/gal.

For application with mechanically pressurized wand to greenhouse ornamentals, **DO NOT** exceed 0.0025 lb ai/gal

**DO NOT** exceed 600 gallons spray volume per acre for foliar applications.

**PLANTS GROWN FOR FRUIT AND NUTS**

**RESTRICTION:** Heritage may be applied to fruit and nut trees, fruiting vines, brambles and bushberries grown in ornamental production nurseries and intended for resale to the consumer market. In some instances, immature and or/inedible berries, fruits, or nuts may be present on the plant at the time of the application but are not intended for immediate harvest or consumption.

*[On the following Fruit and Nut tables, use the **oz product/A** rate (column 3) for products 8 oz or larger. Use the **oz product/0.5 A** rate (column 4) for 4 oz products. Use the **oz product/1,000 sq ft** rate (column 5) for all product sizes.]*



**TABLE 15: Specific Use Directions for Almonds**

| Crop    | Target Diseases   | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application Instructions  |
|---------|---|--|---|---|---|
| Almonds | Alternaria leaf and fruit spot<br>( <i>Alternaria alternata</i> )<br>Anthracnose<br>( <i>Colletotrichum acutatum</i> )<br>Leaf Blight<br>( <i>Seimatosporium lichenicola</i> )<br>Leaf rust<br>( <i>Tranzschelia discolor</i> )<br>Scab<br>( <i>Cladosporium carpophilum</i> )<br>Shothole<br>( <i>Wilsonomyces carpophilus</i> ) | 3.2-8.0<br>(0.1-0.25)                    | 1.6-4.0<br>(0.1-0.25)                           | 0.075-0.18                                      | <p><b>Resistance Management:</b> Do not apply more than two sequential applications of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout the disease development period following the resistance management guidelines. Applications may be made by ground, air (minimum 15 GPA) or chemigation. Heritage Fungicide may be applied by air only at growth stages prior to and including 5 weeks after petal fall. An adjuvant may be added at specified rates.</p> <p>For anthracnose, scab and shothole, begin applications prior to disease development and continue at 7- to 14-day intervals throughout the disease development period.</p> |
|         | Brown Rot Blossom Blight<br>( <i>Monilinia laxa</i> ,<br><i>M. fructicola</i> )   | 6.4 – 8.0<br>(0.2-0.25)                  | 3.2-4.0<br>(0.2-0.25)                           | 0.15-0.18                                       | For blossom blight, begin applications at early bloom and continue through petal fall.  |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- Do not apply within 28 days of harvest (28-day PHI).

**TABLE 16: Specific Use Directions for Bananas and Plantains**

| Crop                         | Target Diseases  | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application Instructions   |
|------------------------------|--|--|---|---|--|
| <b>Bananas<br/>Plantains</b> | Black Sigatoka<br>( <i>Mycosphaerella<br/>fijiensis</i> )<br>Yellow Sigatoka<br>( <i>Mycosphaerella<br/>musicola</i> ) | 2.9-4.3<br>(0.09-<br>0.135)              | 1.45-2.15<br>(0.09-<br>0.135)                   | 0.07-0.1  | <p><b>Integrated Pest (Disease) Management:</b> Heritage Fungicide should be integrated into an overall disease management strategy that includes canopy management through removal of suckers, proper plant spacing, selection of varieties with disease tolerance, removal of plant debris in which inoculum overwinters, and good surface water drainage.</p> <p><b>Resistance Management:</b> Follow the resistance management guidelines in the <b>Resistance Management</b> section. Do not apply more than two sequential applications of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout the disease development period every 12-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> |

**Specific Use Restrictions:**

- Do not apply more than 1.08 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 17: Specific Use Directions for Berries, Bushberry Subgroup**

| Crop  | Target Diseases  | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application Instructions  |
|---|--|--|---|---|---|
| <b>Berries<br/>Bushberry<br/>subgroup</b><br><br>Blueberry<br>Currant<br>Elderberry<br>Gooseberry<br>Huckleberry<br>Lingonberry<br>Juneberry<br>Salal<br><br>including all<br>cultivars<br>and/or<br>hybrids of<br>these      | Alternaria Leaf<br>Spot and Fruit<br>Rot<br>( <i>Alternaria</i> spp.)<br>Anthracnose fruit<br>rot<br>( <i>Colletotrichum<br/>gloeosporoides</i> )<br>Botryosphaeria<br>canker<br>( <i>Botryosphaeria</i><br>spp.)<br>Mummyberry<br>( <i>Vaccinium</i><br>spp.)<br>Phomopsis stem<br>canker<br>( <i>Phomopsis<br/>vaccinii</i> )<br>Powdery mildew<br>( <i>Podosphaera</i><br>spp.)<br>Septoria blight<br>( <i>Septoria</i> spp.) | 3.2-8.0<br>(0.1-0.25)                    | 1.6-4.0<br>(0.1-0.25)                           | 0.08-<br>0.18                                   | <b>Resistance Management:</b><br>Follow the resistance<br>management guidelines in<br>the <b>Resistance<br/>Management</b> section. Do<br>not apply more than two<br>sequential applications of<br>Heritage Fungicide or other<br>Group 11 fungicides before<br>alternation with a fungicide<br>that is not in Group 11.<br><br>Begin Heritage Fungicide<br>applications prior to disease<br>development and continue<br>throughout the disease<br>development period on a 7-<br>to 14- day schedule,<br>following the resistance<br>management guidelines.<br>Applications may be made<br>by ground, air or<br>chemigation. An adjuvant<br>may be added at specified<br>rates. |
| <b>Specific Use Restrictions:</b> <ul style="list-style-type: none"> <li>Do not apply more than 0.75 lb ai/A per year of azoxystrobin-containing products.</li> <li>May be applied the day of harvest (0-day PHI).</li> </ul> |  |  |   |   |   |

**TABLE 18: Specific Use Directions for Berries, Caneberry Subgroup**

| Crop   | Target Diseases  | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application Instructions  |
|--|--|--|---|---|---|
| <b>Berries</b><br><b>Caneberry</b><br><b>subgroup</b><br><br>Blackberry<br>Bingleberry<br>Boysenberry<br>Dewberry<br>Lowberry<br>Marionberry<br>Olallieberry<br>Youngberry<br>Loganberry<br>Red and<br>black<br>raspberry<br><br>including all<br>cultivars<br>and/or<br>hybrids of<br>these | Anthracnose<br>( <i>Sphaceloma</i><br><i>necator</i> )<br>( <i>Elsinoe veneta</i> )<br>Botryosphaeria<br>canker<br>( <i>Botryosphaeria</i><br><i>dothidea</i> )<br>Colletotrichum rot<br>( <i>Colletotrichum</i><br><i>gloeosporioides</i> )<br>Leaf spot<br>( <i>Septoria rubi</i> )<br>( <i>Sphaerulina</i><br><i>rubi</i> )<br>Powdery mildew<br>( <i>Podosphaera</i><br><i>macularis</i> )<br>Rosette or double<br>blossom of<br>blackberries<br>( <i>Cercospora</i><br><i>rubi</i> )<br>Spur blight<br>( <i>Didymella</i><br><i>applanata</i> ) | 3.2-8.0<br>(0.1-0.25)                    | 1.6-4.0<br>(0.1-0.25)                           | 0.08-<br>0.18                                   | <b>Integrated Pest (Disease)</b><br><b>Management:</b> Heritage<br>Fungicide should be<br>integrated into an overall<br>disease management<br>strategy that includes<br>varieties with disease<br>tolerance, proper timing of<br>irrigation and removal of<br>plant debris in which<br>inoculum overwinters.<br><br><b>Resistance Management:</b><br>Follow the resistance<br>management guidelines in<br>the <b>Resistance</b><br><b>Management</b> section. Do<br>not apply more than two<br>sequential applications of<br>Heritage Fungicide or<br>other Group 11 fungicides<br>before alternation with a<br>fungicide that is not in<br>Group 11.<br><br>Begin applications at onset<br>of disease and continue as<br>required until harvest.<br>Make applications on a 7-<br>to 14-day schedule.<br>Applications may be made<br>by ground, air or<br>chemigation. Use a<br>minimum water volume of<br>10 gallons per acre by<br>ground and a minimum of<br>3 gallons by air. |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 19: Specific Use Directions for Citrus Fruit**

| Crop  | Target Diseases  | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application<br>Instructions   |
|---|--|--|---|---|---|
| <b>Citrus Fruit</b><br><br>Calamondin<br>Citron<br>Grapefruit<br>Kumquat<br>Lemon<br>Lime<br>Mandarin<br>Orange (sour<br>and sweet)<br>Pummelo<br>Satsuma<br>mandarin<br>Tangerine<br><br>including all<br>cultivars<br>and/or<br>hybrids of<br>these | Albinism<br>( <i>Alternaria alternata</i><br>pv. <i>citri</i> )<br>Alternaria leaf and<br>fruit spot<br>( <i>Alternaria citri</i> )<br>Cercospora leaf spot<br>( <i>Cercospora</i> spp.)<br>Diplodia stem-end rot<br>( <i>Lasiodiplodia</i><br><i>theobromae</i> )<br>Greasy spot<br>( <i>Mycosphaerella</i><br><i>citri</i> )<br>Melanose<br>( <i>Diaporthe citri</i> )<br>Penicillium Decays<br>Green mold,<br>Whisker mold,<br>suppression of Blue<br>mold<br>( <i>Penicillium</i> spp.)<br>Phomopsis stem-end<br>rot<br>( <i>Phomopsis citrii</i> )<br>Post-bloom fruit drop<br>(PFD)<br>( <i>Colletotrichum</i><br><i>acutatum</i> )<br>Powdery mildew<br>( <i>Erysiphe</i> spp.)<br>Scab<br>( <i>Elsinoe fawcettii</i> ) | 6.4-8.0<br>(0.2-0.25)                    | 3.2-4.0<br>(0.2-0.25)                           | 0.15-<br>0.18                                   | <b>Integrated Pest<br/>(Disease)</b><br><b>Management:</b> Heritage<br>Fungicide should be<br>integrated into an<br>overall disease<br>management strategy<br>that includes selection<br>of varieties with disease<br>tolerance, removal of<br>plant debris in which<br>inoculum overwinters,<br>and proper timing of<br>irrigation.<br><br><b>Resistance</b><br><b>Management:</b> Follow<br>the resistance<br>management guidelines<br>in the <b>Resistance</b><br><b>Management</b> section.<br>Do not apply more than<br>two sequential<br>applications of Heritage<br>Fungicide or other<br>Group 11 fungicides<br>before alternation with a<br>fungicide that is not in<br>Group 11. Do not make<br>more than four (4)<br>applications of Heritage<br>Fungicide or other<br>Group 11 fungicide per<br>disease development<br>period.<br><br>Begin Heritage<br>Fungicide applications<br>prior to disease<br>development and<br>continue throughout the<br>disease development |

|  |  |  |  |  |   |
|--|--|--|--|--|---|
|  |  |  |  |  | period on 7- to 21-day intervals following the resistance management guidelines. Under conditions that favor severe disease epidemics, apply Heritage Fungicide at the highest listed rate. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. Use a horticultural spray oil to improve control of greasy spot. |
|--|--|--|--|--|---|

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 20: Specific Use Directions for Grapes**

| Crop   | Target Diseases   | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application Instructions  |
|--|---|--|---|---|---|
| <b>Grapes</b><br><br>including<br>Muscadines | Black rot<br>( <i>Guignardia<br/>bidwellii</i> )<br>Downy Mildew<br>( <i>Plasmopara<br/>viticola</i> )<br>Phomopsis<br>cane and<br>leaf spot<br>( <i>Phomopsis<br/>viticola</i> )<br>Powdery<br>mildew<br>( <i>Uncinula<br/>necator</i> )<br><br><b>Suppression<br/>Only:</b><br>Botrytis bunch<br>rot<br>( <i>Botrytis<br/>cinerea</i> ) | 5.1-8.0<br>(0.16-<br>0.25)               | 2.55-4.0<br>(0.16-0.25)                         | 0.11-<br>0.18                                   | <p><b>Integrated Pest (Disease) Management:</b> Heritage Fungicide should be integrated into an overall disease management strategy that includes canopy management through pruning and thinning, proper selection of varieties with disease tolerance, proper timing and placement of irrigation and removal of plant debris in which inoculum overwinters.</p> <p><b>Resistance Management:</b> Follow the resistance management guidelines in the <b>Resistance Management</b> section. Do not apply more than two sequential foliar applications of Heritage Fungicide or other Group 11 fungicides before alternating with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout the disease development period every 10-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p><b>ATTENTION</b><br/>Heritage Fungicide is extremely phytotoxic to certain apple varieties. DO NOT spray Heritage Fungicide where spray drift may reach apple trees.</p> |

|  |  |  |  |  |   |
|--|--|--|--|--|---|
|  |  |  |  |  | <p><b><u>DO NOT</u></b> use spray equipment which has been previously used to apply Heritage Fungicide to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.</p> <p>AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.</p> |
|--|--|--|--|--|---|

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- Do not apply within 14 days of harvest (14-day PHI).



**TABLE 21: Specific Use Directions for Pecans**

| Crop   | Target Diseases  | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application Instructions  |
|--------|--|--|---|---|---|
| Pecans | Anthracnose<br>( <i>Glomerella<br/>cingulata</i> )<br>Scab<br>( <i>Cladosporium<br/>caryigenum</i> ) | 3.2-6.4<br>(0.10-<br>0.20)               | 1.6-3.2<br>(0.10-0.20)                          | 0.08-<br>0.15                                   | <p><b>Integrated Pest (Disease) Management:</b> Heritage Fungicide should be integrated into an overall disease management strategy that includes selection of varieties with tolerance to disease and removal of plant debris in which inoculum overwinters.</p> <p><b>Resistance Management:</b> Follow the resistance management guidelines in the <b>Resistance Management</b> section. Do not apply more than two sequential applications of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout the disease development period on 7- to 21-day intervals following the resistance management guidelines.</p> <p>Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> |

**Specific Use Restrictions:**

- Do not apply more than 1.2 lb ai/A per year of azoxystrobin-containing products.
- Do not apply within 45 days of harvest (45-day PHI).

**TABLE 22: Specific Use Directions for Pistachios**

| Crop              | Target Diseases   | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application Instructions   |
|-------------------|---|--|---|---|--|
| <b>Pistachios</b> | <p>Alternaria late blight<br/>(<i>Alternaria alternata</i>)</p> <p>Botryosphaeria panicle and shoot blight<br/>(<i>Botryosphaeria dothidea</i>)</p> <p>Septoria leaf spot<br/>(<i>Septoria pistaciarum</i>)</p> | 3.2-8.0<br>(0.10-0.25)                   | 1.6-4.0<br>(0.10-0.25)                          | 0.08-0.18                                       | <p><b>Integrated Pest (Disease) Management:</b> Heritage Fungicide should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance and removal of plant debris in which inoculum overwinters.</p> <p><b>Resistance Management:</b> Follow the resistance management guidelines in the <b>Resistance Management</b> section. Do not apply more than two sequential applications of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout the disease development period on 7- to 21-day intervals following the resistance management guidelines.</p> <p>Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- Do not apply within 7 days of harvest (7-day PHI).

**TABLE 23: Specific Use Directions for Stone Fruit**

| Crop   | Target Diseases  | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application<br>Instructions  |
|--|--|--|---|---|--|
| <b>Stone<br/>Fruit</b><br><br>Apricot<br>Cherry,<br>sweet<br>Cherry,<br>tart<br>Nectarine<br>Peach<br>Plum<br>Plumcot<br>Prune | Alternaria spot and fruit<br>rot<br>( <i>Alternaria alternata</i> )<br>Anthracnose<br>( <i>Colletotrichum<br/> prunicola</i> ,<br><i>C. gloeosporioides</i> )<br>Leaf rust<br>( <i>Tranzschelia discolor</i> )<br>Powdery mildew<br>( <i>Podosphaera<br/> pannosa</i> , <i>Podosphaera<br/> clandestina</i> )<br>Scab<br>( <i>Cladosporium<br/> carpophilum</i> )<br>Shot hole<br>( <i>Wilsonomyces<br/> carpophilus</i> ) | 3.2-8.0<br>(0.1-0.25)                    | 1.6-4.0<br>(0.1-0.25)                           | 0.08-<br>0.18                                   | <b>Resistance<br/> Management:</b> Follow<br>the resistance<br>management<br>guidelines in the<br><b>Resistance<br/> Management</b> section.<br>Do not apply more<br>than two sequential<br>applications of<br>Heritage Fungicide or<br>other Group 11<br>fungicides before<br>alternation with a<br>fungicide that is not in<br>Group 11.<br><br>For brown rot blossom<br>blight, begin<br>applications at early<br>bloom and continue<br>through petal fall. For<br>brown rot on fruit,<br>Heritage Fungicide<br>may be applied to fruit<br>up to the day of<br>harvest. For scab,<br>begin applications at<br>petal fall and continue<br>at 7- to 14-day<br>intervals. For all other<br>diseases, begin<br>application at the onset<br>of disease as a<br>protectant fungicide<br>and continue on a 7- to<br>14-day schedule. For<br>peaches only, 5-8 oz.<br>of Heritage Fungicide<br>may be used for scab<br>control. |
|  | Brown rot blossom blight<br>and fruit rot<br>( <i>Monilinia fructicola</i> , <i>M.<br/> laxa</i> )   | 6.4-8.0<br>(0.2-0.25)                    | 3.2-4.0<br>(0.2-0.25)                           | 0.15-<br>0.18                                   | Applications may be  |

|  |  |  |  |  |                                       |
|--|--|--|--|--|---------------------------------------|
|  |  |  |  |  | made by ground, air or<br>chemigation |
|--|--|--|--|--|---------------------------------------|

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 24: Specific Use Directions for Strawberry**

| Crop       | Target Diseases  | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application Instructions  |
|------------|--|--|---|---|---|
| Strawberry | Anthracnose<br>( <i>Colletotrichum fragariae</i> )<br>Powdery mildew<br>( <i>Podosphaera macularis</i> )<br><br><b><u>Suppression</u></b><br><b><u>only:</u></b> Botrytis<br>on the foliage<br>( <i>Botrytis cinerea</i> ) | 3.2-8.0<br>(0.1-0.25)                    | 1.6-4.0<br>(0.1-0.25)                           | 0.08-<br>0.18                                   | <b>Resistance Management:</b><br>Follow the resistance<br>management guidelines in<br>the <b>Resistance</b><br><b>Management</b> section. Do<br>not apply more than two<br>sequential applications of<br>Heritage Fungicide or other<br>Group 11 fungicides before<br>alternation with a fungicide<br>that is not in Group 11.<br><br>Begin Heritage Fungicide<br>applications prior to disease<br>development and continue<br>throughout the disease<br>development period on a 7-<br>to 10-day schedule, following<br>the resistance management<br>guidelines. Applications may<br>be made by ground, air or<br>chemigation. An adjuvant<br>may be added at specified<br>rates. |

**Specific Use Restrictions:**

- Do not apply more than 1.0 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 25: Specific Use Directions for Tree Nuts (except Almonds, Pecans, and Pistachios)**

| Crop   | Target Diseases   | Use Rate<br>oz<br>product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A<br>(lb ai/A) | Use<br>Rate<br>oz<br>product/<br>1,000 sq<br>ft | Application Instructions   |
|--|---|--|---|---|--|
| <b>Tree Nuts</b><br><br>Beechnut<br>Brazil nut<br>Butternut<br>Cashew<br>Chestnut<br>Chinquapin<br>Filbert<br>Hickory<br>Macadamia<br>Walnut<br><br>Almonds,<br>Pecans,<br>Pistachios:<br>see specific<br>use<br>instructions. | Alternaria leaf<br>and fruit spot<br>( <i>Alternaria<br/> alternata</i> )<br>Anthracnose<br>( <i>Colletotrichum<br/> acutatum</i> ,<br><i>Glomerella<br/> cingulata</i> )<br>Eastern filbert<br>blight<br>( <i>Anisogramma<br/> anomale</i> )<br>Late blight<br>( <i>Alternaria<br/> alternata</i> )<br>Scab<br>( <i>Cladosporium<br/> carpophilum</i> )<br>Septoria leaf<br>spot<br>( <i>Septoria<br/> pistaciarum</i> )<br>Shothole<br>( <i>Wilsonomyces<br/> carpophilus</i> ) | 3.2-6.4<br>(0.10-<br>0.20)               | 1.6-3.2<br>(0.10-0.20)                          | 0.08-<br>0.15                                   | <b>Resistance Management:</b><br>Follow the resistance<br>management guidelines in<br>the <b>Resistance<br/> Management</b> section. Do<br>not apply more than two<br>sequential applications of<br>Heritage Fungicide or other<br>Group 11 fungicides before<br>alternation with a fungicide<br>that is not in Group 11.<br><br>Begin Heritage Fungicide<br>applications prior to disease<br>development and continue<br>throughout the disease<br>development period<br>following the resistance<br>management guidelines.<br>Applications may be made<br>by ground, air or<br>chemigation. An adjuvant<br>may be added at specified<br>rates.<br><br>For all other diseases, begin<br>applications prior to disease<br>development and continue at<br>7- to 21-day intervals<br>throughout the disease<br>development period. |
|  | Blossom blight<br>( <i>Monilinia laxa</i> ,<br><i>M. fructicola</i> )   | 6.4<br>(0.20)                            | 3.2<br>(0.20)                                   | 0.15  | For blossom blight, begin<br>applications at early bloom<br>and continue through petal<br>fall. <b>Do not</b> make more than<br>six applications of Heritage<br>Fungicide or other strobilurin<br>fungicide per acre per year.   |

**Specific Use Restrictions:**

- Do not apply more than 1.2 lb ai/A per year of azoxystrobin-containing products.
- Do not apply within 45 days of harvest (45-day PHI).

**TABLE 26: Specific Use Instructions for Tropical Fruit**

| Crop  | Target Diseases   | Use Rate<br>oz product/A<br>(lb ai/A) | Use Rate<br>oz<br>product/0.5<br>A (lb ai/A) | Use Rate<br>oz<br>product/1,000<br>sq ft | Application<br>Instructions   |
|---|---|---------------------------------------|--|--|---|
| <b>Tropical Fruit</b><br>Acerola<br>Atemoya<br>Avocado<br>Biriba<br>Canistel<br>Cherimoya<br>Custard apple<br>Feijoa<br>Guava<br>Llana<br>Jaboticaba<br>Jackfruit<br>Longan<br>Loquat<br>Lychee<br>Mango<br>Papaya<br>Passionfruit<br>Pawpaw<br>Persimmon<br>Pulasan<br>Pummello<br>Rambutan<br>Sapodilla<br>Sapote, black<br>Sapote, mamey<br>Sapote, white<br>Soursop<br>Star apple<br>Starfruit<br>Sugar apple<br>Spanish lime<br>Tamarind<br>Uniq fruit | Alternaria leaf and fruit spot ( <i>Alternaria</i> spp.)<br>Anthracnose ( <i>Colletotrichum</i> spp.)<br>Cercospora leaf spot ( <i>Cercospora</i> spp.)<br>Powdery mildew ( <i>Erysiphe</i> spp.)<br>Rust ( <i>Puccinia</i> spp.) | 3.2-8.0<br>(0.1-0.25)                 | 1.6-4.0<br>(0.1-0.25)                        | 0.08-0.18                                | <b>Resistance Management:</b><br>Follow the resistance management guidelines in the <b>Resistance Management</b> section. Do not apply more than two sequential applications of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.<br><br>Begin Heritage Fungicide applications prior to disease development and continue throughout the disease development period on a 10- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

## VEGETABLE AND HERB PLANTS

**RESTRICTION:** Apply to vegetable and herb plants grown for retail sale to consumers only.

**TABLE 27: Specific Use Directions for Asparagus Plants**

| Crop      | Target Diseases   | Use Rate<br>oz product/<br>1,000 sq ft | Use Rate<br>oz product/<br>5,000 sq ft | Application Instructions   |
|-----------|---|--|--|--|
| Asparagus | Stemphyllium<br>purple spot<br>( <i>Stemphyllium<br/>vesicarium</i> ) | 0.075-0.18                             | 0.375-0.9                              | <p><b>Resistance Management:</b> Do not apply more than one application of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout plant production on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by foliar sprays, including chemigation. An adjuvant may be added at specified rates. Use a minimum of 10 gallons of water per acre.</p> |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- Do not apply within 100 days of harvest (100-day PHI).
- A use rate of 0.18 oz product/1,000 sq ft is equal to 0.25 lb ai/A.



**TABLE 28: Specific Use Directions for Brassica Head and Stem Subgroup Plants**

| Crop   | Target Diseases   | Use Rate<br>oz product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz product/<br>5,000 sq ft | Application Instructions   |
|--|---|---|--|--|
| <b>Brassica Head and stem subgroup</b><br><br>Broccoli<br>Chinese broccoli [gai lon]<br>Brussels sprouts<br>Cabbage<br>Chinese cabbage [napa]<br>Chinese mustard cabbage [gai choy]<br>Cauliflower<br>Cavalo broccolo<br>Kohlrabi<br><br>including all cultivars and/or hybrids of these | Alternaria leaf spot ( <i>Alternaria</i> spp.)<br><br>Downy mildew ( <i>Peronospora parasitica</i> )<br><br>Pin rot ( <i>Alternaria</i> spp.) | 0.08-0.18<br>(0.1-0.25)                             | 0.4-0.9                                | <b>Resistance Management:</b><br>Do not apply more than one application of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.<br><br>Begin Heritage Fungicide applications prior to disease development and continue throughout plant production on a 7- to 14-day schedule, following the resistance management guidelines.<br><br>Applications may be made by foliar sprays including chemigation. An adjuvant may be added at specified rates. Use a minimum of 10 gallons of water per acre. |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 29: Specific Use Directions for Brassica Leafy Greens Subgroup Plants**

| Crop   | Target Diseases  | Use Rate<br>oz<br>product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz<br>product/<br>5,000 sq ft | Application Instructions  |
|--|--|--|---|---|
| <b>Brassica<br/>Leafy greens<br/>subgroup</b><br><br>Broccoli raab<br>Cabbage,<br>Chinese<br>Collards<br>Kale<br>Mizuna<br>Mustard greens<br>Mustard spinach<br>Rape greens<br><br>including all<br>cultivars and/or<br>hybrids of these | Black spot<br>( <i>Alternaria</i><br>spp.)<br>Cercospora leaf<br>spot<br>( <i>Cercospora</i><br>spp.)<br>White rust<br>( <i>Albugo</i><br><i>candida</i> ) | 0.08-0.18<br>(0.1-0.25)                                | 0.4-0.9                                   | <p><b>Resistance Management:</b> Follow the resistance management guidelines in the <b>Resistance Management</b> section. Do not apply more than one application of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout plant production on a 7- to 14-day schedule, following the resistance management guidelines.</p> <p>Applications may be made by foliar sprays including chemigation. An adjuvant may be added at specified rates. Use a minimum of 10 gallons of water per acre.</p> |

**Specific Use Restrictions:**

- Do not apply more than 0.75 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 30: Specific Use Directions for Bulb Vegetable Plants**

| Crop  | Target Diseases  | Use Rate<br>oz product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz.<br>product/<br>5,000 sq ft | Application Instructions   |
|---|--|---|--|--|
| <b>Bulb Vegetables</b><br><br>Garlic<br>Leek<br>Onion, bulb<br>Onion, green<br>Welsh onion<br>Shallot | <b>Foliar Diseases</b><br>Cladosporium leaf blotch<br>( <i>Cladosporium allii</i> )<br>Purple blotch<br>( <i>Alternaria porri</i> )<br>Rust<br>( <i>Puccinia allii</i> )<br>White rot<br>( <i>Sclerotium cepivorum</i> ) | 0.08-0.15<br>(0.1-0.20)                             | 0.4-0.75                                   | <b>Resistance Management:</b> Do not apply more than one application of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.<br><br>For downy mildew, make preventative applications on a 5- to 7-day schedule. For all other diseases, begin Heritage Fungicide applications prior to disease development and continue throughout plant production every 7-14 days, following the resistance management guidelines. |
|   | Botrytis leaf blight<br>( <i>Botrytis aclada</i> )<br>Downy mildew<br>( <i>Peronospora destructor</i> )  | 0.11-0.18<br>(0.15-0.25)                            | 0.55-0.9                                   | Applications may be made by foliar sprays including chemigation. An adjuvant may be added at specified rates.<br><br>Mixtures of Heritage Fungicide with insecticides and silicone adjuvants should be tested for crop safety before application to the crop.  |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 31: Specific Use Directions for Celery Plants**

| Crop   | Target Diseases   | Use Rate<br>oz product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz<br>product/<br>5,000 sq ft | Application Instructions   |
|--------|---|---|---|--|
| Celery | <p>Early blight<br/>(<i>Cercospora apii</i>)<br/>Late blight<br/>(<i>Septoria apicola</i>)</p> <p>For additional<br/>diseases, see <b>Leafy<br/>Vegetables</b>.</p> | <p>0.11-0.18<br/>(0.15-0.25)</p>                    | <p>0.55-0.9</p>                           | <p><b>Resistance Management:</b> Do not apply more than one application of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout plant production every 7-14 days, following the resistance management guidelines.</p> <p>Applications may be made by foliar sprays including chemigation. An adjuvant may be added at specified rates.</p> |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0 day PHI).

**TABLE 32: Specific Use Directions for Cucurbit Plants**

| Crop  | Target Diseases  | Use Rate<br>oz<br>product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz<br>product/<br>5,000 sq<br>ft | Application Instructions  |
|---|--|--|--|---|
| <b>Cucurbits</b><br><br>Cantaloupe<br>Chayote<br>Chinese-waxgourd<br>Cucumber<br>Gourds<br>Honeydew<br>Melons<br><i>Momordica</i> spp.<br>(bitter melon, balsam apple)<br>Muskmelon<br>Watermelon<br>Pumpkin<br>Squash<br>Zucchini<br><br>including cultivars and/or hybrids of these | Anthracnose<br>( <i>Colletotrichum lagenarium</i> )<br>Belly Rot<br>( <i>Rhizoctonia solani</i> )<br>Downy Mildew<br>( <i>Pseudoperonospora cubensis</i> )<br>Gummy Stem Blight<br>( <i>Didymella bryoniae</i> )<br>Leaf spots<br>( <i>Alternaria</i> spp., <i>Cercospora</i> spp.)<br>Myrothecium canker<br>( <i>Myrothecium roridum</i> )<br>Plectosporium blight<br>( <i>Plectosporium tabacinum</i> )<br>Powdery Mildew<br>( <i>Podosphaera xanthii</i> ,<br><br><i>Erysiphe cichoracearum</i> )<br>Ulocladium leaf spot<br>( <i>Ulocladium cucurbitae</i> ) | 0.08-0.18<br>(0.10-0.25)                               | 0.4-0.9                                      | <p><b>Resistance Management:</b> Do not apply more than one application of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11. Do not make more than four (4) foliar applications of Heritage Fungicide or other Group 11 fungicides per crop per acre per year.</p> <p>For both downy and powdery mildew, make preventative applications on a 5- to 7-day schedule. For belly rot control, make the first application at the 1- to 3-leaf crop stage with a second application just prior to vine tip over or 10-14 days later, whichever occurs first. For all other diseases, begin Heritage Fungicide applications prior to disease development and continue throughout plant production every 7-14 days, following the resistance management guidelines.</p> <p>Applications may be made by foliar sprays, including chemigation.</p> <p>An adjuvant may be added at specified rates. However, do not tank-mix Heritage Fungicide with COC, MSO or silicon adjuvants.</p> <p>Do not tank-mix Heritage Fungicide with malathion, methomyl, potassium laurate, or dicloran containing products.</p> |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.

- Do not apply within 1 day of harvest (1-day PHI).

**TABLE 33: Specific Use Directions for Herb and Spice Plants**

| Crop  | Target Diseases   | Use Rate<br>oz product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz product/<br>5,000 sq ft | Application<br>Instructions  |
|---|---|---|--|--|
| <b>Herbs &amp; Spices<br/>(except basil and<br/>black pepper)</b><br>Allspice; Angelica;<br>Anise (seed); Anise,<br>star; Annatto; Balm;<br>Borage; Burnet;<br>Camomile; Caper<br>(buds); Caraway;<br>Caraway, black;<br>Cardamom; Cassia<br>(buds); Catnip;<br>Celery seed; Chervil<br>(dried); Chive; Chive,<br>Chinese; Cinnamon;<br>Clary; Clove (buds);<br>Coriander (cilantro or<br>Chinese parsley)<br>(leaf); Coriander<br>(seed); Costmary;<br>Culantro (leaf and<br>seed); Cumin; Curry<br>(leaf); Dill (seed);<br>Dillweed; Fennel,<br>common; Fennel,<br>Florence (seed);<br>Fenugreek; Grains of<br>paradise;<br>Horehound; Hyssop;<br>Juniper (berry);<br>Lavender;<br>Lemongrass; Lovage<br>(leaf and seed);<br>Mace; Marigold;<br>Marjoram; Mustard<br>(seed), Nasturtium;<br>Nutmeg; Oregano;<br>Parsley (dried);<br>Pennyroyal; Pepper,<br>white; Poppy seed;<br>Rosemary; Rue;<br>Saffron; Sage;<br>Savory, summer and<br>winter; Sweet bay;<br>Tansy; Tarragon; | Alternaria leafspot<br>( <i>Alternaria</i> spp.)<br>Corynespora blight<br>( <i>Corynespora</i><br><i>cassicola</i> )<br>Downy mildew<br>(except Basil)<br>( <i>Plasmopara</i> spp.,<br><i>Peronospora</i> spp.)<br>Dill blight<br>( <i>Cercosporidium</i><br><i>punctum</i> )<br>Phoma blight<br>( <i>Passalora puncta</i> )<br>Powdery mildew<br>( <i>Erysiphe</i> spp.,<br><i>Podosphaera</i> spp.) | 0.08-0.18<br>(0.1-0.25)                             | 0.4-0.9                                | <b>Resistance<br/> Management:</b> Do not<br>apply more than two<br>sequential applications<br>of Heritage Fungicide or<br>other Group 11<br>fungicides before<br>alternation with a<br>fungicide that is not in<br>Group 11.<br><br>Begin Heritage<br>Fungicide applications<br>at the onset of disease<br>development and<br>continue throughout<br>plant production on a 7-<br>day schedule, following<br>the resistance<br>management<br>guidelines.<br><br>Applications may be<br>made by foliar sprays.<br>An adjuvant may be<br>added at specified<br>rates. Use a minimum<br>of 30 gallons of water<br>per acre. |

|  |  |  |  |  |
|--|--|--|--|--|
| Thyme; Vanilla;<br>Wintergreen;<br>Woodruff;<br>Wormwood |  |  |  |  |
|--|--|--|--|--|

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).
- Do not apply by aerial application.

**TABLE 34: Specific Use Directions for Basil Plants**

| Crop  | Target Diseases                                      | Use Rate<br>oz product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz product/<br>5,000 sq ft | Application Instructions  |
|-------|--|---|--|---|
| Basil | Downy mildew<br>( <i>Peronospora<br/>belbahrii</i> ) | 0.18<br>(0.25)                                      | 0.9                                    | <p><b>Plug Production*</b>: Apply to emerged plants in plug production trays prior to disease development. Apply uniformly to foliage using a minimum of 3.4 gallons of water/5,000 sq ft (30 gallons/A). Make no more than one application during the plug production phase. Follow the Heritage Fungicide application with alternative chemistries on a weekly schedule, implementing a preventative integrated disease management program.</p> <p><b>Finish Production**</b>: Apply to plants following transplant of plugs to trays, pots or containers in which plants are grown to finish. Apply uniformly to foliage using a minimum of 3.4 gallons of water/5,000 sq ft (30 gallons/A). Make no more than one application during the finish production phase. Follow the Heritage Fungicide application with alternative chemistries on a weekly schedule, implementing a preventative integrated disease management program. For specific resistance management programs, contact your state Extension specialist.</p> |

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

\* Plug production refers to the production of a young plant grown from seed in a multi-celled germination tray for a short period of time. After growing to a desired size, the plug is then transplanted in a larger pot or container to grow to a larger size suitable to sell.

\*\* Finish production refers to the production of a finished plant grown to a desired size suitable to sell in a garden center, large format retailer, or other retailers selling plants to consumers for home and garden plantings.



**TABLE 35: Specific Use Directions for Leafy Vegetable Plants (except Brassica)**

| Crop  | Target Diseases  | Use Rate<br>oz product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz product/<br>5,000 sq ft | Application Instructions  |
|---|--|---|--|---|
| <b>Leafy Vegetables (except Brassica)</b><br><br>Amaranth<br>Arugula<br>Cardoon<br>Celery<br>Celtuce<br>Chervil<br>Chrysanthemum, edible<br>Coriander, leaves (Cilantro)<br>Corn salad<br>Cress<br>Dandelion<br>Dock<br>Endive<br>Fennel<br>Lettuce, head and leaf<br>Orach<br>Parsley<br>Purslane<br>Radicchio<br>Rhubarb<br>Spinach<br>Swiss Chard<br><br>including cultivars and/or hybrids of these | <b>Foliar Diseases</b><br>Alternaria leaf spot ( <i>Alternaria sonchi</i> , <i>A. spp.</i> )<br>Anthracnose ( <i>Microdochium panattonianum</i> , <i>Colletotrichum dematium</i> )<br>Cercospora leaf spot ( <i>Cercospora spp.</i> )<br>Septoria leaf spot ( <i>Septoria petroselini</i> )<br>White rust ( <i>Albugo occidentalis</i> ) | 0.08-0.18<br>(0.1-0.25)                             | 0.4-0.9                                | <b>Resistance Management:</b><br>Do not apply more than one application of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.<br><br>For both downy and powdery mildew, make preventative applications on a 5- to 7-day schedule. For all other diseases, begin Heritage Fungicide applications prior to disease development and continue throughout plant production every 7-14 days following the resistance management guidelines.   |
|   | Downy mildew ( <i>Bremia lactucae</i> )<br>Powdery mildew ( <i>Erysiphe cichoracearum</i> )  | 0.15-0.18<br>(0.2-0.25)                             | 0.75-0.9                               | Applications may be made by foliar sprays including chemigation. An adjuvant may be added at specified rates.<br><br><b>ATTENTION:</b> Applications of Heritage Fungicide to leafy vegetable foliage have contributed to phytotoxicity under certain circumstances. Proceed with caution with regard to tank mixes and adjuvants when treating all leafy vegetables with Heritage Fungicide. Heritage Fungicide must not be tank-mixed on leaf lettuce with permethrin, fosetyl-al, lambda-cyhalothrin, or another product that may increase the penetration of Heritage Fungicide into the |

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  | leaf surface, such as, but not limited to, silicone wetters. |
|--|--|--|--|--|

**Specific Use Restrictions:**

- Do not apply more than 1.5 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 36: Specific Use Directions for Mint Plants**

| Crop            | Target Diseases  | Use Rate<br>oz product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz product/<br>5,000 sq ft | Application Instructions  |
|-----------------|--|---|--|---|
| Mint<br>(Fresh) | Powdery mildew<br>( <i>Erysiphe</i> spp.)<br>Rust<br>( <i>Puccinia<br/>menthae</i> ) | 0.08-0.18<br>(0.1-0.25)                             | 0.4-0.9                                | <p><b>Resistance Management:</b> Do not apply more than two sequential applications of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout plant production on a 7- to 10-day schedule, following the resistance management guidelines.</p> <p>Applications may be made by foliar sprays including chemigation. An adjuvant may be added at specified rates.</p> |

**Specific Use Restrictions:**

- Do not apply more than 0.75 lb ai/A per year of azoxystrobin-containing products.
- For fresh mint, may be applied the day of harvest (0-day PHI).

**TABLE 37: Specific Use Directions for Pepper and Other Fruiting Vegetable Plants (except Cucurbits)**

| Crop   | Target Diseases  | Use Rate<br>oz product/<br>1,000 sq ft<br>(lb ai/A) | Use Rate<br>oz<br>product/<br>5,000 sq<br>ft | Application Instructions   |
|--|--|---|--|--|
| <b>Peppers and other Fruiting Vegetables (except cucurbits)</b><br>Bell Pepper<br>Non-Bell Pepper<br>Sweet Non-Bell Pepper<br>Eggplant<br>Groundcherry<br>Okra<br>Pepino<br>Tomatillo<br><br>See specific directions for use for Tomatoes. | Anthracnose ( <i>Colletotrichum</i> spp.)<br>Cercospora leaf spot ( <i>Cercospora capsici</i> )<br>Downy mildew ( <i>Peronospora tabacini</i> )<br>Powdery mildew<br><i>Leveillula taurica</i> | 0.08-0.18<br>(0.1-0.25)                             | 0.4-0.9                                      | <b>Resistance Management:</b><br>Follow the resistance management guidelines in the <b>Resistance Management</b> section. Do not apply more than one application of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.<br><br>Begin Heritage Fungicide applications prior to disease development and continue throughout plant production on a 7- to 14-day schedule, following the resistance management guidelines.<br><br>Applications may be made by foliar sprays including chemigation. An adjuvant may be added at specified rates. |

**Specific Use Restrictions:**

- Do not apply more than 1.0 lb ai/A per year of azoxystrobin-containing products.
- May be applied the day of harvest (0-day PHI).

**TABLE 38: Specific Use Directions for Tomato Plants**

| Crop     | Target Diseases   | Use Rate<br>oz product/1,000<br>sq ft<br>(lb ai/A) | Use Rate<br>oz product/<br>5,000 sq ft | Application Instructions  |
|----------|---|--|--|---|
| Tomatoes | Anthracnose<br>( <i>Colletotrichum<br/>coccodes</i> )<br>Black Mold<br>( <i>Alternaria<br/>alternata</i> )<br>Buckeye Rot<br>( <i>Phytophthora<br/>spp.</i> )<br>Early Blight<br>( <i>Alternaria solani</i> )<br>Powdery Mildew<br><i>Leveillula<br/>taurica</i><br>Septoria Leaf<br>spot<br>( <i>Septoria<br/>lycopersici</i> )<br>Target spot<br>( <i>Corynespora<br/>cassicola</i> ) | 0.024-0.08<br>(0.08-0.10)                          | 0.12-0.4                               | <p><b>Resistance Management:</b><br/>Do not apply more than one application of Heritage Fungicide or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Begin Heritage Fungicide applications prior to disease development and continue throughout plant production following the resistance management guidelines. For late blight, apply Heritage Fungicide at 5- to 7-day intervals. For all other tomato diseases, apply Heritage Fungicide on 7- to 21-day intervals.</p>   |
|          | Late Blight<br>( <i>Phytophthora<br/>infestans</i> )  | 0.08<br>(0.10)                                     | 0.4                                    | <p>Applications may be made by foliar sprays including chemigation.</p> <p>Under certain environmental conditions (particularly high temperatures), Heritage Fungicide in combination with high rates of silicone-containing or oil-containing (petroleum or crop) additives or adjuvants may cause injury. <b>Do not</b> exceed 0.125% adjuvant (v/v). Consult a Syngenta representative for more information concerning additives or adjuvants.</p> <p>A tank mixture with dimethoate may cause crop injury.</p> <p>Do not use adjuvants or tank mix Heritage Fungicide with any emulsifiable concentrate (EC) product.</p> |

**Specific Use Restrictions:**

- 
- Do not apply Heritage Fungicide until 35 days after seeding or 21 days after transplanting plugs to larger pots or containers.
  - Do not apply more than 0.6 lb ai/A per year of azoxystrobin-containing products.
  - May be applied the day of harvest (0-day PHI).
- 

**TABLE 39: Heritage Rate Conversion Chart**  
(For use with 4 oz package size only)

| <b>oz product/A</b> | <b>oz product/<br/>1,000 sq ft</b> | <b>Treated Acres/<br/>4 oz product</b> |
|---------------------|------------------------------------|--|
| 1.0                 | 0.025                              | 4.0                                    |
| 1.5                 | 0.035                              | 2.7                                    |
| 2.0                 | 0.05                               | 2.0                                    |
| 2.5                 | 0.06                               | 1.6                                    |
| 3.0                 | 0.07                               | 1.3                                    |
| 3.5                 | 0.08                               | 1.1                                    |
| 4.0                 | 0.09                               | 1.0                                    |
| 4.5                 | 0.1                                | 0.9                                    |
| 5.0                 | 0.11                               | 0.8                                    |
| 5.5                 | 0.13                               | 0.73                                   |
| 6.0                 | 0.14                               | 0.67                                   |
| 6.5                 | 0.15                               | 0.62                                   |
| 7.0                 | 0.16                               | 0.57                                   |
| 7.5                 | 0.17                               | 0.53                                   |
| 8.0                 | 0.18                               | 0.5                                    |
| 8.7                 | 0.2                                | 0.46                                   |
| 13.1                | 0.3                                | 0.31                                   |
| 17.4                | 0.4                                | 0.23                                   |

## **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

### **Pesticide Storage**

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, sweep and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

### **Pesticide Disposal**

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### **Container Handling [(less than or equal to 50 pounds)]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling [(bags)]**

Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

Heritage®, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company.

©20XX Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-866-796-4368.

Manufactured for:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

Heritage Fungicide 1093 MAS 0316 AMEND-E 0519-CL – jab- 7/23/25  
000100-01093.20190515E.HERITAGE FUNGICIDE.AMEND.MAY2019-CL