

PRODUCT INFORMATION



Syngenta Crop Protection, Inc.
Greensboro, NC 27409
www.syngenta-us.com

RESTRICTED USE PESTICIDE
Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABEL FOR KARATE® INSECTICIDE WITH ZEON™ TECHNOLOGY

Insecticide
EPA Reg. No. 100-1097

Supplemental Directions for Use on Non-Cropland (Excluding Public Land) and Sugarcane

Active Ingredient:

Lambda-cyhalothrin

[1α(S*),3α(Z)]-(±)-cyano-(3-phenoxyphenyl)methyl-
3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-

dimethylcyclopropanecarboxylate 22.8%

Other Ingredients: 77.2%

Total: 100.0%

Karate Insecticide with Zeon Technology contains 2.08 lbs. of active ingredient per gal. and is a capsule suspension.

Contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN.

ACCEPTED
AUG - 5 2003
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 100-1097


WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

SCP 1097A-S1 0703



DIRECTIONS FOR USE

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
Non-Cropland (Excluding Public Land)	See Crop Outlets	See Crop Outlets	See Crop Outlets	<ul style="list-style-type: none"> • Spray non-cropland adjacent to agricultural areas to control migratory insects, which may threaten crops. • Follow General Use Directions, rates and spray recommendations found elsewhere in this label for the adjacent crop outlet and target pests. • Use highest labeled rates for dense/large foliage, high insect populations and larger larval stages. • Repeat as necessary to maintain control. • Do not exceed 0.2 lb. ai (0.8 pt.) /A per year. • Do not graze livestock in treated areas.

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
Sugarcane	Sugarcane Borer ¹ Rice Borer ¹ Sugarcane Beetle (Adult) ² Yellow Sugarcane Aphid ³	0.025-0.04	1.6-2.56	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. • Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 2 gal. of water /A. • Do not apply within 21 days of harvest. • Do not apply more than 0.16 lb. ai (1.28 pt.) /A per season. <p>¹For control before the larva bores into the plant stalk. ²Suppression only of beetles active above ground. ³See Resistance statement under General Directions for Use.</p>

Karate®, Zeon™ and the Syngenta logo are trademarks of a Syngenta Group Company
 © 2003 Syngenta

Syngenta Crop Protection, Inc.
 Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1097A-S1 0703
 Kar 1097A-S1 0703-Ig-7-30-03

