

Please read instructions on reverse before completing form.

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number NOTIFICATION
--	--	--	---------------------------------------

Application for Pesticide - Section I

1. Company/Product Number 100-1108	2. EPA Product Manager (acting) Dan Rosenblatt	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) TOUCHDOWN® 5 HERBICIDE	PM# Team 25	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. Registrations & Regulatory Affairs 410 Swing Road PO Box 18300 Greensboro NC 27409-8300 <input checked="" type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.	NOTIFICATION APR 27 2001
--	---	---

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. The following changes are being made via this notification: 1) Company name and address have been updated to reflect Syngenta Crop Protection, Inc., 2) EPA Reg. No. changed to new Company number, 3) The copyright date reflects Syngenta., 4) Trademark statements have been updated to reflect Syngenta for those products for which Syngenta holds the trademark. 5) The Internet address has been changed to reflect Syngenta. 6) Other places in the label which refer to the company name have been updated.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
*Certification must be submitted		If "Yes" Unit Packaging wgt. No. per Container	If "Yes" Unit Packaging wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Martina A. Haw	Title Regulatory Assistant II	Telephone No. (Include Area Code) 302 / 476-2373
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Martina Haw</i>	3. Title Regulatory Assistant II	
4. Typed Name Martina A. Haw	5. Date April 18, 2001	

PAPERWORK REDUCTION ACT NOTICE. Public reporting burden for this collection of information is estimated to average 0.85 hour per response, including time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Chief, Information Policy Branch, (2136), U.S. Environmental Protection Agency, 401 M Street, SW, Washington, DC 20460.

INSTRUCTIONS This form is to be used for all applications for new registration, end use reregistration, amendment, resubmission, to applications for notifications, final printed labeling, reregistration, etc. In order to process an application for a new registration submitted on this form, the following material must accompany the application:

1. Certification with Respect to Citation of Data (EPA Form 8570-29). [If not exempted by 40 CFR 152.81 (b) (4)];
2. Confidential Statement of Formula (EPA Form 8570-4);
3. Formulator's Exemption Statement (EPA Form 8570-27);
4. Five copies of draft labeling;
5. Three copies of any data submitted;
6. Authorization letter where applicable;
7. Matrices where applicable.

Submission of Labeling - Labeling should first be submitted in the form of draft labels with all applications for new registration. Such draft labels may be in the form of typed label text on 8.5 x 11 inch paper for submission or a mockup of the proposed label. If prepared for mockup, it should be constructed in a way as to facilitate storage in an 8.5 x 11 inch file. Mockup labels significantly smaller than 8.5 x 11 inches should be mounted on 8.5 x 11 inch paper for submission. Submission of Data - Data submitted in support of this application must be submitted in accordance with PR Notice 86-5.

SPECIFIC INSTRUCTIONS: Please read the instructions listed below before completing this application. First determine the type of registration action, listed in Block A, for which you are submitting this application. For applications submitted in connection with New Registration actions, Sections I, III, and IV must be completed by the applicant. For applications submitted in connection with amended reregistration actions, resubmissions, notifications, reregistrations, etc., Sections I, II, and IV must be completed by the applicant.

Block A - Check the appropriate action for which you are submitting this form.

SECTION I - This section must be completed, as applicable, for all registration actions.

1. Company/Product Number - Insert your Company Number, if one has been assigned by EPA. This number may have been assigned to you as a basic registrant, a distributor, or as an establishment. If your product is registered, insert the Product Number.
2. EPA Product Manager - If known, fill in the name and PM number of the EPA Product Manager.
3. Proposed Classification - Specify the proposed classification of this product.
4. Product Name - Enter the complete product name of this pesticide as it will appear on the label. The name must be specific to this product only. Duplication of names is not permitted among products of the same company. Do not include any brand name or company line designations.
5. Name and Address of Applicant - The name of the firm or person and address shown in your application is the person or firm to whom the registration will be issued. If you are acting in behalf of another party, you must submit authorization from that party to act for them in registration matters. An applicant not residing in the United States must have an authorized agent residing in the United States to act for them in all registration matters. The name and complete mailing address of such an agent must accompany this application.
6. Expedited Review - FIFRA section 3 (c) 3 (B) provides for expedited review of applications for registration, or amendments to existing registrations, that are similar or identical to other pesticide products that are currently registered with the EPA. In order for your application to be eligible for expedited review, you must provide us with the EPA Registration Number and product name of the product you believe is similar to or identical to your product. The product must be similar or identical in both formulation and labeled uses.

SECTION II - This section must be completed for all applications submitted to amend the registration only of a currently registered product (Amendment), for a resubmission in response to an Agency letter, for notifications to the Agency, for the submission of final printed labeling, for reregistration and for any other action that pertains to a specific EPA-registered product. This section is not to be used for a new application for registration.

1. Subject of submission - Check the applicable block and provide the Agency letter date if appropriate. Provide a brief explanation of the purpose(s) for the submission, such as "the addition of a site, pest or crop (specify)"; "amend the Confidential Statement of Formula by..."; "reregistration submission"; "general label revision of use directions." Attach a separate page if additional space is needed.

SECTION III - (Packaging and Container Information) - This Section must be completed for all applications submitted in connection with now registration or applicable amendments.

1. Type of Packaging - Check the appropriate block if your product will be packaged in the indicated packaging types. Indicate the size of the individual packets and number per retail container.
2. Type of Retail Container - Indicate type of container in which product will be marketed.
3. Location of Net Contents - Indicate the location of the net contents information for your product.
4. Size(s) of Retail Container - Specify the net contents of all retail containers for your product.
5. Location of Use Directions - Indicate the location of the use directions for your product.
6. Manner in which label is affixed to product - Indicated the method product label is attached to retail container.

SECTION IV (Contact Point) - This Section must be completed for all applications for Registration actions, i.e., now products registration, resubmission, "me-too," reregistration, etc.

- 1-5. Self-explanatory.
6. EPA Use Only

NOTIFICATION

APR 27 2001



3 8 43

TOUCHDOWN® 5 Herbicide

NONSELECTIVE FOLIAR SYSTEMIC HERBICIDE FOR WEED CONTROL

ACTIVE INGREDIENT:	
Sulfosate	48.6%
INERT INGREDIENTS:	
Total	100.0%

Contains 5 pounds active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements and directions
for use (inside booklet; on back panel; on side panel)

NOTIFICATION

APR 27 2001

EPA Reg. No. 100-1108
EPA Est. No.

Product of Belgium
Formulated and Packaged in USA.

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com


SCP 100-xxxx

Net Weight/U.S. Standard Measure

TABLE OF CONTENTS

	Page
STATEMENT OF PRACTICAL TREATMENT	
PRECAUTIONARY STATEMENTS	
Personal Protective Equipment	
User Safety Recommendations	
Environmental Hazards	
Drift	
DIRECTIONS FOR USE	
AGRICULTURAL USE REQUIREMENTS	
NONAGRICULTURAL USE REQUIREMENTS	
STORAGE AND DISPOSAL	
GENERAL INFORMATION	
GENERAL USE PRECAUTIONS	
APPLICATION DIRECTIONS	
Timing	
Rates	
Spray Additives	
Surfactants/Wetting Agents	
Ammonium Sulfate	
Tankmixes with Residual Herbicides	
Application Equipment & Techniques	
Ground	
Air	
Spot Treatments	
Wiper Application	
CROPS	
BERRIES, FRUITS, NUTS, VEGETABLES, AND VINES	
Method of Application	
General Use Precautions	
Tank Mixtures with Residual Herbicides and 2,4-D	
Hard to Control Weed Recommendations in Citrus (Florida and Texas Only)	
Perennial Grass Suppression (Chemical Mowing) of Orchard Floors	
CONSERVATION COMPLIANCE/CONSERVATION RESERVE PROGRAM (CRP)	
CORN (FIELD, POPCORN, AND SEED CORN)	
Method of Application	
General Use Precautions	
Tank Mixtures for Corn	
FALLOWLAND AND POSTHARVEST USE	
Method of Application	
Chemical Fallow-Ecofallow	
Postharvest Chemical Fallow	
Aid to Tillage	
Fallow Beds/Stale Seed Beds	
Postharvest Use	
General Use Precautions	
FARMSTEAD (NONCROP)	

SOYBEANS
 Method of Application
 Preharvest
 General Use Precautions
 Tank Mixtures for Soybeans (Preplant/Preemergence)
 SOYBEANS, GLYPHOSATE-TOLERANT (including ROUNDUP READY® soybeans)
 Method of Application
 General Use Precautions
 Preplant/Preemergence
 Postemergence
 Annual Weeds
 Perennial Weeds
 Tank Mixtures
 Preharvest
 WHEAT
 Method of Application
 General Use Precautions
 Tank Mixtures
 WEEDS CONTROLLED
 North/ South Region Annual Weed Control - Table 1
 Western Region Annual Weed Control - Table 2
 Annual Weed Control - Tankmix with Dicamba or 2,4-D - Table 3
 Perennial and Woody Plant Weed Control-Alone or with Dicamba or 2,4-D - Table 4
 STORAGE AND DISPOSAL



STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

HOTLINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal)
or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident)
Call
1-800-888-8372

PRECAUTIONARY STATEMENTS

Hazards to Humans And Domestic Animals**CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: long sleeved shirt and long pants, socks and shoes, and waterproof gloves.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

In case of spill, ISOLATE the spill. Absorb spill with inert absorbent material such as clay or Fuller's earth. Sweep up used absorbent and place in an appropriate chemical waste container. Flush spill area with water. Observe all local, State, and Federal laws and regulations regarding disposal, spill, cleanup, removal, or discharge.

DRIFT: Caution must be taken when applying TOUCHDOWN 5 to avoid drift or contact with nontarget plant species. Such contact may result in plant injury.

8 2 43

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Syngenta Crop Protection, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Syngenta and Seller harmless for any claims relating to such factors.

Syngenta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Syngenta, and Buyer and User assume the risk of any such use. **SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

In no event shall Syngenta or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Syngenta and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of Syngenta.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Waterproof gloves.
- Shoes plus socks.

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not contaminate water, food, or feed by storage and disposal. Keep container tightly closed when not in use. For help with any spill, leak, fire, or accident involving this material, call 1-800-888-8372.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

FOR RECYCLABLE/REFILLABLE CONTAINERS:

Before refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices. After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container.

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

GENERAL INFORMATION

TOUCHDOWN 5 is a nonselective foliar systemic herbicide used to control a broad spectrum of emerged grass and broadleaf weeds, both annual and perennial, in:

- berries, fruits, nuts, vegetables, and vines specified on this label
- conservation compliance/conservation reserve program (CRP)
- corn
- fallowland and postharvest
- soybeans, including glyphosate-tolerant
- wheat
- certain noncrop areas around the farm specified on this label

TOUCHDOWN 5 is formulated as a liquid concentrate which contains 5 pounds active ingredient per gallon.

GENERAL USE PRECAUTIONS

- TOUCHDOWN 5 requires actively growing green plant tissue to function. Application to drought-stressed weeds or weeds with little green foliage (i.e., mowed or cut weeds), may result in reduced weed control.
- TOUCHDOWN 5 does not provide residual control of weeds. Weeds emerging after application will require retreatment.
- Heavy rainfall or irrigation shortly after application may require retreatment.
- Tillage or mowing within 3 days following application may reduce weed control.
- Do not apply this product through any type of irrigation system.
- DO NOT spray if conditions of thermal inversion exist, or if wind direction and speed may cause spray to drift onto adjacent nontarget areas. Drift minimization is the responsibility of the applicator. Consult with local and State agricultural authorities for information regarding avoiding or minimizing spray drift.
- It is recommended that the spray system be thoroughly cleaned with water and a commercial tank cleaner after each use.
- Stainless steel, plastic-lined steel, plastic or fiberglass containers are acceptable for mixing and storing TOUCHDOWN 5. TOUCHDOWN 5 should not be mixed or stored in galvanized steel or unlined steel containers. TOUCHDOWN 5 is not compatible with carbon steel.
- Any crop not listed on this label may be planted back into TOUCHDOWN 5 treated areas 35 days after application.
- Do not exceed a total of 6.4 pints of TOUCHDOWN 5 per acre per year in corn.
- Do not exceed a total of 12.8 pints of TOUCHDOWN 5 per acre per year in all other uses on this label except corn.
- Do not graze or harvest treated cover crops for feed.

APPLICATION DIRECTIONS

Timing

TOUCHDOWN 5 should be applied to actively growing emerged weeds. Annual weeds of 6 inches or less in height are typically the easiest to control. Generally, more effective control of perennial weeds is achieved at the flowering or seedhead stage. Refer to the **WEEDS CONTROLLED** section, beginning on Page _____, for specific application timing.

When annual weeds have been mowed or grazed, wait for 3-4 inches of new growth to appear prior to application. When perennial weeds have been mowed or grazed, allow new growth to reach recommended stage prior to application.

Rates

Follow recommended rates for TOUCHDOWN 5 listed in the **WEEDS CONTROLLED** section, beginning on Page _____. Use the higher label rates when weeds are dense or large. Also, use higher application volumes and pressures when weed vegetation is dense. The following table will assist in rate conversions:

TOUCHDOWN 5 RATE CONVERSION TABLE			
LBS A/A	FL OZS/A	PINTS/A	ACRES/ GALLON
0.5	12.8	0.8	10
0.625	16.0	1.0	8
0.75	19.2	1.2	6.7
1.0	25.6	1.6	5
1.5	38.4	2.4	3.3
2.0	51.2	3.2	2.5
3.0	76.8	4.8	1.7
4.0	102.4	6.4	1.25
6.0	153.6	9.6	0.8
8.0	204.8	12.8	0.63

SPRAY ADDITIVES

Surfactants/Wetting Agents

A nonionic surfactant (NIS) or wetting agent (approved for use on growing crops) may be used at levels up to 0.25% v/v (1 quart NIS or wetting agent/100 gallons) of finished spray volumes. All nonionic surfactants or wetting agents should contain at least 75% active ingredient.

Ammonium Sulfate (AMS)

Control of annual and perennial weeds with TOUCHDOWN 5 may be improved by adding dry ammonium sulfate at 1 - 2% by weight or 8.5 - 17 pounds per 100 gallons of water. Liquid formulations of AMS may be used at an equivalent rate. Do not reduce use rates of TOUCHDOWN 5 when using AMS.

TANKMIXES WITH RESIDUAL HERBICIDES

Refer to crop sections for recommended tankmixes. Tankmixes of TOUCHDOWN 5 with other pesticides, fertilizers, or any other additives except as specified on this label or other approved Syngenta supplemental labeling may result in tankmix incompatibility or unsatisfactory performance. It is recommended that the compatibility of any tankmix combination be tested on a small scale such as a jar test before actual tank mixing.

Always refer to labels of other pesticide products for mixing directions and precautions which may differ from those outlined here. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing

Tank Mixing Recommendations:

1. Fill spray tank ½ full with clean water.
2. Begin tank agitation and continue throughout mixing and spraying.
3. Add ammonium sulfate (if used).
4. Add dry formulations (WP, DF, etc.) to tank.
5. Add liquid formulations (SC, EC, L, etc.) to tank.
6. Add TOUCHDOWN 5.
7. Add nonionic surfactant/wetting agent (if used).
8. Fill remainder of spray tank.

APPLICATION EQUIPMENT & TECHNIQUES

- Avoid drift. Do not apply in low level inversion conditions, when winds are gusty or under any other conditions which favor drift. Inversions are characterized by stable air and increasing temperatures with height above the ground. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer. Drift may cause damage to any vegetation contacted to which treatment is not intended.
- Drift control additives may be used with TOUCHDOWN 5. Read and follow the manufacturer's directions for use.
- All aerial equipment must be properly maintained and washed to remove product residues after use.

Broadcast Applications

Ground

Applications should be made in 3 to 40 gallons of water per acre.

When foliage is dense, spray volume should be increased to ensure coverage of the target weeds. Flat-fan nozzles will result in the most effective application of TOUCHDOWN 5. Spray boom and nozzle heights must be adjusted to provide coverage of target weed. Flood nozzles may result in reduced weed control due to inadequate coverage.

Air

Applications should be made in 3 to 15 gallons of water per acre.

Spray should be released at the lowest height consistent with effective weed control and flight safety. Applications more than 10 feet above the canopy should be avoided.

Use the largest droplet size consistent with good weed control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding inappropriate spray boom pressure. Solid stream or low shear nozzles may be utilized to reduce small droplet formation. These nozzles direct the fluid parallel to the existing airflow to reduce shear effects. Other techniques may include reducing the fan angle of flat fan nozzles if used, or reducing the deflector plate angle if deflector type nozzles are used. Ensure the spray is released at an appropriate distance below the airfoil.

For best results, each specific aerial application vehicle used should be quantifiably pattern tested for aerial application of TOUCHDOWN 5 herbicide initially and every year thereafter. To minimize drift, it is suggested aerial application equipment produce the following minimum spray deposition characteristics:

Volume Median Diameter (VMD) > 400 microns
Volume Diameter (VD) {0.9} > 200 microns

Spot Treatments

For annual weeds less than 6 inches, use a 0.4 to 0.75% v/v solution. For annual weeds over 6 inches, use a 0.75 to 1.5% v/v solution. Use a 1 to 2% v/v solution for most perennials (see Table 4 for specific rates and timing). When using motorized spot spray equipment (rider bar), use a 3% v/v solution. See Spot Spray Dilution Table below for rates of TOUCHDOWN 5/volume of finished spray solution. Spray the solution on actively growing weeds until uniformly wet but not to the point of runoff. Retreat 14-21 days later if regrowth occurs.

TOUCHDOWN 5 Herbicide Spot Spray Dilution Table

Solution Strength	To Make This Volume			
	1 gallon	10 gallons	25 gallons	100 gallons
0.4%	0.5 fl oz	5 fl oz	0.8 pints	3.2 pints
0.75%	1 fl oz	0.6 pints	1.5 pints	3 quarts
1%	1.3 fl ozs	0.8 pints	2 pints	1 gallon
1.25%	1.6 fl ozs	1 pint	2.5 pints	5 quarts
1.5%	2 fl ozs	1.25 pints	3 pints	6 quarts
2%	2.6 fl ozs	1.5 pints	4 pints	2 gallons
3%	4 fl ozs	2.5 pints	6 pints	3 gallons

Wiper Application

TOUCHDOWN 5 may be applied using a wiper or "wick" applicator for selective control or suppression of annual and perennial weeds which become taller than the crop or desirable vegetation. Mix 1 quart of TOUCHDOWN 5 in 1 gallon of water. Precautions should be taken to avoid contact with crops or desirable vegetation. Equipment should be operated at speeds of 5 mph or less. Use slower speeds where weeds are dense. For improved control, make two applications in opposite directions. Do not use wiper applications in corn or in bearing grape vineyards.

CROPS

This section is organized alphabetically by crop categories. There may be several crops listed in a crop category.

Any crop not listed on this label may be planted back into TOUCHDOWN 5 treated areas 35 days after TOUCHDOWN 5 application.

BERRIES , FRUITS, NUTS, VEGETABLES, AND VINES

TOUCHDOWN 5 may be used on both bearing and nonbearing crops listed below.

BEARING CROPS

Almond
Apple
Apricot
Beechnut
Brazil nut
Butternut
Calamondin
Cashew
Cherry (sweet, sour, tart)
Chestnut
Chinquapin
Chironja
Citron
Citrus Citron
Crabapple
Filbert (Hazelnut)
Grapefruit
Grapes (all)
Hickory nut
Kumquat
Lemon
Lime
Loquat
Macadamia
Mandarin
Mayhaw
Nectarine
Orange (all)
Oriental Pear
Peach
Pear
Pecan
Plum (all)
Plumcot
Prune
Pummelo
Quince
Satsuma Mandarin
Tangelo
Tangerine
Tangor
Walnut (Black, English)

NONBEARING CROPS

TOUCHDOWN 5 can be used 1 year or more prior to harvest in these crops.

Acerola (West Indian Cherry)
Artichoke
Asparagus
Atemoya
Avocado
Banana
Blackberry
Blueberry
Boysenberry
Breadfruit
Canistel
Carambola
Cherimoya
Cocoa Bean
Coconut
Coffee
Cranberry
Currant
Date
Dewberry
Elderberry
Fig
Genip
Ginseng
Gooseberry
Guava
Huckleberry
Jaboticaba
Jackfruit
Jojoba
Kiwi
Loganberry
Longan
Lychee
Mango
Olallieberry
Olive
Papaya
Passion Fruit
Persimmon
Pineapple
Pistachio
Plantain
Pomegranate
Raspberry (black, red)
Sapodilla
Sapote
Soursop
Sugar Apple
Tamarind
Tea
Youngberry

METHOD OF APPLICATION: Preplant; preemergence: directed spray (except Cranberry); middles (between rows of trees); strips (in rows of trees); perennial grass suppression (chemical mowing); and wiper/wick applicator equipment.

GENERAL USE

Applications may be made with boom equipment; shielded sprayers; CDA; hand-held and high-volume wands; lances; orchard guns; or wiper/wick application equipment, except as directed in the **GENERAL USE PRECAUTIONS FOR BERRIES, FRUITS, NUTS, VEGETABLES, AND VINES** section. Follow directions listed in the **APPLICATION DIRECTIONS, SPRAY ADDITIVES, and APPLICATION EQUIPMENT AND TECHNIQUES** sections of this label. Refer to the **WEEDS CONTROLLED** section, beginning on Page _____, for application rates and timing.

Multiple applications may be necessary to control certain perennial weeds. For residual weed control, tank mix TOUCHDOWN 5 with residual herbicides as prescribed in the **TANK MIXTURES FOR BERRIES, FRUITS, NUTS, VEGETABLES, AND VINES** section, or make multiple applications.

GENERAL USE PRECAUTIONS FOR BERRIES, FRUITS, NUTS, VEGETABLES, AND VINES

- Do not allow the spray, spray drift, or mist to contact foliage, fruit, shoots, branches, canes, suckers, open wounds or green parts of crops. Contact with any crop part other than mature brown woody bark can result in severe crop injury.
- Avoid contact with stumps as injury to adjacent trees may occur from root grafting.
- For **PEACHES** grown in *Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, and Tennessee*, apply TOUCHDOWN 5 with a shielded applicator which prevents contact with foliage, suckers, or bark of trees. Apply no later than 90 days after first bloom to avoid severe damage. Avoid application to peach trees with recent mechanical injury or pruning wounds. Apply only near trees which have been planted in the orchard for two or more years. **SEVERE INJURY WILL OCCUR IF ANY PORTION OF THE PEACH TREE IS CONTACTED WITH SPRAY OR SPRAY DRIFT.**
- For **APRICOTS, NECTARINES, PEACHES, PLUMS and PRUNES** grown in *Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah, and Washington*, any application equipment listed for these crops may be used.
- For **APRICOTS, NECTARINES, PEACHES, PLUMS, and PRUNES** grown in all other states not previously listed, use only wiper/wick application equipment.
- For **GRAPES** grown in the *Great Lakes and Northeast regions*, apply TOUCHDOWN 5 prior to the end of bloom stage to avoid injury, or apply with shielded equipment.
- For **COFFEE and BANANA**, delay application 3 months after transplanting to allow the new plants to become established.
- Allow at least 13 days from the last application to harvest of citrus, pome, or stone fruit.
- Allow at least 14 days from the last application to harvest of grapes.
- Allow at least 20 days from the last application to harvest of nuts.

TANK MIXTURES WITH RESIDUAL HERBICIDES AND 2,4-D FOR BERRIES, FRUITS, NUTS, VEGETABLES, AND VINES

TOUCHDOWN 5 can be tank mixed with the following herbicides for control or suppression of annual and perennial weeds provided that the tankmix product label allows use of the product. Refer to the **WEEDS CONTROLLED** section, beginning on Page _____, for application rates and timing. Apply TOUCHDOWN 5 at 1.2 to 6.4 pints/A in these tankmixes for control or suppression of annual and perennial weeds. For control or suppression of dense populations or weeds greater than 12 inches in height or weeds under stress, consider use rates at the higher end of the rate range.

DEVIRINOL®	Krovar®	Sim-Trol®
Direx®	Princep® 4L	Cinibar®
Goal®	Princep Caliber 90®	Solicam®
Karmex®	Prowl®	Surflan®
Kerb®	Simazine	2,4-D

Refer to the individual product labels for precautionary statements, restrictions, recommended rates, approved crops, and a list of weeds controlled.

HARD TO CONTROL WEED RECOMMENDATIONS IN CITRUS (FLORIDA AND TEXAS ONLY)

To control or suppress the perennial weeds listed in the following table, apply the recommended rate of TOUCHDOWN 5 in 3 to 30 gallons of water per acre. Use 10-30 gallons per acre if weed foliage is dense. Apply when weeds are actively growing. Refer to the **PERENNIAL WEED CONTROL** section, Table 4, for application timing. If weeds have been mowed or grazed, allow new growth to reach recommended growth stage prior to application.

Weed Species	Rate of TOUCHDOWN 5 (Pints per Acre)			
	1.6	3.2	4.8	6.4
Bermudagrass	B	B	PC	C
Guineagrass				
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	NR	B	C	C
Paragrass	B	C	C	C
Torpedograss	NR	S	PC	PC

B = Burndown

C = Control

NR = Not Recommended

PC = Partial Control

S = Suppression

PERENNIAL GRASS SUPPRESSION (CHEMICAL MOWING) OF ORCHARD FLOORS

BAHIAGRASS

TOUCHDOWN 5 can be used to inhibit seedhead emergence and suppress vegetative growth for approximately 40 to 50 days with a single application. By using a sequential application, suppression of vegetative growth and inhibition of seedhead emergence can be extended to 120 days. Applications must be made prior to seedhead emergence. Apply TOUCHDOWN 5 at 20 to 30 days after complete green-up or after mowing to 4 inches tall. When a single application is planned, use 5 fluid ounces of TOUCHDOWN 5 per acre in 10 to 20 gallons of water. When a sequential application is planned, use 3½ fluid ounces per acre for the first application followed by another application of 3½ fluid ounces per acre 40 to 50 days later.

BERMUDAGRASS

For Suppression Only:

East of the Rocky Mountains:

Apply 5 to 12 fluid ounces of TOUCHDOWN 5 in 3 to 20 gallons of water per acre. Make the application 2 weeks after complete green-up or after 3 to 4 inches of regrowth following mowing. Use 5 to 8 fluid ounces per acre if a lesser degree of suppression is desired. A sequential application can be used when regrowth occurs.

West of the Rocky Mountains:

Apply 12 fluid ounces of TOUCHDOWN 5 in 3 to 10 gallons of water per acre. Make the application 2 weeks after complete green-up or after 3 to 4 inches of regrowth following mowing. A sequential application can be used when regrowth occurs.

For Partial Control and Burndown:

TOUCHDOWN 5 can be used for burndown and partial control of bermudagrass at 1.6 to 3.2 pints in 3 to 20 gallons of water per acre. Use 1.6 pints east of the Rocky Mountains and 3.2 pints west of the Rocky Mountains.

Use this treatment only if reduction of the bermudagrass stand can be tolerated. Allow at least 21 days for complete burndown.

COOL SEASON GRASS COVERS (fine fescue, Kentucky bluegrass, orchardgrass, quackgrass, tall fescue)

For suppression of orchardgrass, fine fescue, tall fescue, and quackgrass, apply 6 fluid ounces of TOUCHDOWN 5 in 10 to 20 gallons of water per acre. Add AMS. See **SPRAY ADDITIVES** section for rates.

For suppression of Kentucky bluegrass, use 5 fluid ounces of TOUCHDOWN 5. Do not add AMS for suppression of Kentucky bluegrass.

CONSERVATION COMPLIANCE/ CONSERVATION RESERVE PROGRAM (CRP)

METHOD OF APPLICATION: Rotating out of CRP, site preparation (sequential herbicide applications), dormant beneficial plant management.

- **Site Preparation:** Prior to application, removal of excessive vegetation by grazing, mowing, burning, etc. may improve control. When annual weeds have been mowed or grazed, wait for 3 to 4 inches of new growth before application. When perennial weeds have been mowed or grazed, allow regrowth to reach recommended stage (see **PERENNIAL WEED CONTROL** section, Table 4 for rates and timing).

Sequential applications of TOUCHDOWN 5 and GRAMOXONE® EXTRA herbicides are effective in controlling established CRP grasses. Refer to the GRAMOXONE EXTRA herbicide label for recommended rates and tankmixes.

TOUCHDOWN 5/GRAMOXONE EXTRA Herbicide Sequential Program: (Spring Application)

Weed Species	Program A	Program B
Fescue Orchardgrass Ryegrass	GRAMOXONE EXTRA at 2-2.5 pints/A followed 7-10 days later with GRAMOXONE EXTRA at 2-2.5 pints/A	TOUCHDOWN 5 at 1.6-2.0 pints/A followed 10-14 days later with GRAMOXONE EXTRA at 2-2.5 pints/A

- **Dormant Beneficial Plant Applications:** Apply 0.6 to 0.8 pints/acre in early spring before desirable species, such as crested and tall wheatgrass, break dormancy. Late fall applications can be made after desirable grasses have reached dormancy. If perennial grasses are not dormant at time of application, stunting can occur.
- TOUCHDOWN 5 may be tank mixed with other herbicides registered for this use such as atrazine, dicamba, and 2,4-D.
- Any crops not listed on this label may be planted back into TOUCHDOWN 5 treated areas 35 days after application provided plant back is not prohibited by labels of other products if used in a tankmix or sequential application.

CORN (Field Corn, Popcorn, and Seed Corn)

METHOD OF APPLICATION: Before, during, or after planting but before crop emergence; spot spray; and postharvest.

Follow the directions listed in the **APPLICATION DIRECTIONS**, **SPRAY ADDITIVES**, and **APPLICATION EQUIPMENT AND TECHNIQUES** sections. Refer to the **WEEDS CONTROLLED** section, beginning on Page _____, for application rates and timing.

GENERAL USE PRECAUTIONS FOR CORN

- Spot application must be made at least 90 days before harvest of corn grain or corn fodder.
- Crop plants contacted by TOUCHDOWN 5 will be injured or killed.
- Do not use on sweet corn.
- Do not apply more than 6.4 pints per acre per year.

TANK MIXTURES FOR CORN

For Control of Annual Weeds in a Residual Herbicide Tankmix : Refer to the **ANNUAL WEEDS CONTROLLED** section, Tables 1 and 2, for application rates and timing. Apply TOUCHDOWN 5 at 0.8 - 4.8 pints/A for the control of annual weeds that are less than 6 inches tall and actively growing. When annual weeds are taller than 6 inches or under stress, use 1.6 - 4.8 pints/A of TOUCHDOWN 5 herbicide.

For Control or Suppression of Perennial Weeds in a Residual Herbicide Tankmix : Refer to the **PERENNIAL WEEDS CONTROLLED** section, Table 4, for application rates and timing. Use rates at the higher end of the rate range when weed populations are dense or plants are under stress. Perennial weeds may require multiple applications for control.

UAN may be used as a carrier at 10 - 70 gallons per acre with 2,4-D, dicamba, or any residual herbicides on the following list. For use with 2,4-D and dicamba on annual and perennial weeds, consult Tables 3 and 4. Reduced weed control may occur on certain weeds as a result of UAN foliar burn which can reduce uptake of TOUCHDOWN 5.

TOUCHDOWN 5 can be tank mixed with the following products:

AMBUSH®	Dicamba	Lasso®
Atrazine	Dual®	Lightning™
Axiom™	Dual II®	Linex®
Balance®	Dual II Magnum™	Lorox®
Basis®	Extrazine® II	Marksman®
Bicep®	Frontier®	Micro-Tech®
Bicep II®	FULTIME™	Prowl
Bicep II Magnum™	Guardzman®	Simazine
Bladex®	Harness®	SURPASS® EC
Broadstrike®	Harness Xtra	SURPASS 100
Bullet®	Hornet™	TOPNOTCH®
Clarity®	KARATE®	WARRIOR®
	Lariat®	2,4-D

Refer to individual product labels for precautionary statements, restrictions, rates, and a list of weeds controlled.

FALLOWLAND AND POSTHARVEST USE

METHOD OF APPLICATION: Chemical fallow; fallow beds; stale seedbeds; aid to tillage and postharvest.

TOUCHDOWN 5 may be applied during the fallow period prior to planting or emergence of any crop listed on this label. For any crop not listed on this label, applications must be made 35 days prior to planting.

Chemical Fallow - Ecofallow

TOUCHDOWN 5 may be used in place of tillage to control annual weeds or volunteer wheat in fallow fields. Repeat applications may be necessary to control weeds emerging after application. Refer to Table 1 and 2 for use rates and timing. Broadcast or spot treatments of TOUCHDOWN 5 will control or suppress perennial weeds. Refer to Table 4 for use rates and timing. Follow directions listed in the **APPLICATION DIRECTIONS, SPRAY ADDITIVES, and APPLICATION EQUIPMENT AND TECHNIQUES** sections of this label.

Tankmixes with 2,4-D or dicamba may be used for additional control of annual weeds listed in Table 3. Tank mixtures with atrazine or cyanazine (e.g. Bladex) may provide residual control of weeds listed on their individual product labels. Cyanazine cannot be used after December 31, 2002.

Postharvest Chemical Fallow for Cereals

TOUCHDOWN 5 may be applied after harvest to control newly emerged weeds, volunteer cereals, or weeds which were present at harvest. Allow sufficient time after harvest for weed regrowth to occur before making application. Refer to Table 1 and 2 for use rates and annual weeds controlled. Higher rates may be required for control of large weeds which were present at the time of harvest. Repeat applications may be necessary for fall germinating weeds. Broadcast or spot treatments of TOUCHDOWN 5 will control or suppress perennial weeds. Refer to Table 4 for use rates and timing. Follow directions listed in the **APPLICATION DIRECTIONS, SPRAY ADDITIVES, and APPLICATION EQUIPMENT AND TECHNIQUES** sections of this label. Do not graze or harvest treated areas for feed.

Tankmixes with 2,4-D or dicamba may be used for additional control of weeds listed in Table 3. Postharvest tankmixes with atrazine or cyanazine may be used if the field will be planted to corn or sorghum or laid fallow the following season. Tankmixes with atrazine or cyanazine may be applied for residual control of certain annual weeds such as common lambsquarters, kochia, mustards, pigweeds, and volunteer wheat. Tankmixes with residual herbicides may result in reduced performance. Cyanazine cannot be used after December 31, 2002.

Aid to Tillage

TOUCHDOWN 5 may be used in conjunction with tillage operations in fallow systems to control cheat, downy brome, foxtails, tansy mustard, and volunteer cereals. Apply 0.4 - 0.6 pints/A of TOUCHDOWN 5 in 3-10 gallons of water per acre. Apply before weeds exceed 6 inches in height. Application must be followed by tillage no later than 15 days after treatment or before weed regrowth. Allow at least one day after application before tillage. Tankmixes with residual herbicides may reduce performance. Follow directions listed in the **APPLICATION DIRECTIONS, SPRAY ADDITIVES, and APPLICATION EQUIPMENT AND TECHNIQUES** sections of this label.

Fallow Beds/Stale Seedbeds

TOUCHDOWN 5 may be used to control weeds in fallow or stale seedbeds. Follow directions listed in the **APPLICATION DIRECTIONS, SPRAY ADDITIVES, and APPLICATION EQUIPMENT AND TECHNIQUES** sections of this label. Refer to **WEEDS CONTROLLED** section, beginning on Page ____, for application rates and timing. TOUCHDOWN 5 can be tank mixed with 2,4-D, dicamba, or Goal herbicide for improved control of certain weeds.

Postharvest Use

TOUCHDOWN 5 may be applied after harvest of any crop to control newly emerged weeds, volunteer crops, or weeds which were present at harvest. Refer to **WEEDS CONTROLLED** Section, beginning on Page ____, for use rates. Repeat applications may be necessary to control weeds emerging after application. Use the higher rate on heavy or sodded infestations.

GENERAL USE PRECAUTIONS FOR FALLOWLAND AND POSTHARVEST USE

- For any crop not listed on this label, TOUCHDOWN 5 applications must be made 35 days prior to planting.
- Allow sufficient time for weed regrowth to occur after harvest before making applications.
- Avoid application after plants have been exposed to a severe frost.
- Refer to the individual labels of all products used in a tankmix for precautionary statements, recropping intervals, restrictions, and a list of weeds controlled.
- TOUCHDOWN 5 will not control volunteer glyphosate-tolerant crops.

FARMSTEAD (NONCROP)

- Applications can be made in noncrop areas on the farm such as:

Barrier Strips	Farmyards
Ditchbanks	Fence Rows
Dry Ditches and Canals	Fuel Storage Areas
Equipment Areas	Rights-of-way
Farm Buildings	Soil Bank Land

- Refer to **WEEDS CONTROLLED** section, beginning on Page ____, for application rates and timing.
- Avoid contact with the foliage of ornamentals or other desirable plants.
- Repeat applications may be necessary.

SOYBEANS

METHOD OF APPLICATION: Before, during, or after planting, but before crop emergence; spot spray; wiper/wick; preharvest; post harvest.

Follow directions listed in the **APPLICATION DIRECTIONS**, **SPRAY ADDITIVES**, and **APPLICATION EQUIPMENT AND TECHNIQUES** sections. Refer to **WEEDS CONTROLLED** section, beginning on Page _____, for application rates and timing.

PREHARVEST

TOUCHDOWN 5 may be applied preharvest as a broadcast spray with ground or aerial equipment as a harvest aid. TOUCHDOWN 5 provides weed control when applied preharvest to soybeans and may aid in crop dry down. Apply to mature soybeans when pods have lost their color.

GENERAL USE PRECAUTIONS FOR SOYBEANS

- Spot application must be made at least 8 weeks before harvest.
- Wiper/wick application must be made at least 7 days before harvest.
- Soybeans, except glyphosate-tolerant varieties, will be injured or killed when contacted with TOUCHDOWN 5.
- Make preharvest applications at least 7 days before harvest with no more than 1.6 pints/acre.
- Do not graze or harvest for hay following harvest aid application.
- Do not apply more than 12.8 pints per acre per year.

TANK MIXTURES FOR SOYBEANS (PREPLANT/PREEMERGENCE)

For Control of Annual Weeds in a Residual Herbicide Tankmix : Refer to the **ANNUAL WEEDS CONTROLLED** section, Tables 1 and 2, for application rates and timing. Apply TOUCHDOWN 5 at 0.8 - 4.8 pints/A for the control of annual weeds that are less than 6 inches tall and actively growing. When annual weeds are taller than 6 inches or under stress, use 1.6 - 4.8 pints/A of TOUCHDOWN 5 herbicide.

For Control or Suppression of Perennial Weeds in a Residual Herbicide Tankmix : Refer to the **PERENNIAL WEEDS CONTROLLED** section, Table 4, for application rates and timing. Use rates at the higher end of the rate range when weed populations are dense or plants are under stress. Perennial weeds may require multiple applications for control.

For use with 2,4-D on perennial weeds, consult Table 4.

TOUCHDOWN 5 can be tank mixed with the following products:

Authority™	Dual II Magnum	Lexone®	REFLEX®
Authority Broadleaf	FirstRate™	Linex	Scepter®
Broadstrike®	FLEXSTAR®	Lorox	Sencor®
Canopy®	Frontier	Lorox Plus	Squadron®
Canopy XL	FUSILADE®	Partner®	Steel™
Command®	FUSION®	Preview®	Turbo®
Cover™	Gemini®	Prowl	WARRIOR
Dual	KARATE	Pursuit	2,4-D
Dual II	Lasso	Pursuit Plus	2,4-DB

Refer to individual product labels for precautionary statements, restrictions, rates, and a list of weeds controlled.

SOYBEANS, GLYPHOSATE-TOLERANT (including ROUNDUP READY® soybeans)

METHOD OF APPLICATION: Before, during, or after planting; postemergence; and preharvest in soybean varieties which have been genetically modified to be tolerant to glyphosate based herbicides. Follow directions listed in the **APPLICATION DIRECTIONS, SPRAY ADDITIVES, and APPLICATION EQUIPMENT & TECHNIQUES** sections. Refer to **WEEDS CONTROLLED** section, beginning on Page _____, for application rates and timing. For best results, use minimum rates of 1.2 pints/acre for annual weeds and 1.6 pints/acre for perennials.

GENERAL USE PRECAUTIONS FOR GLYPHOSATE-TOLERANT SOYBEANS

- Make postemergence applications up to and including the full bloom stage of soybeans at a maximum of 3.2 pints/acre.
- Make preharvest applications at least 7 days before harvest with no more than 1.6 pints per acre.
- Do not graze or harvest for forage or hay.
- Apply no more than 12.8 pints per acre per year.

PREPLANT/PREEMERGENCE

TOUCHDOWN 5 may be used as a broadcast spray to control emerged annual and perennial weeds. Apply before, during, or after planting of soybeans. TOUCHDOWN 5 can be tank mixed with the following products:

Authority	Dual II Magnum	Lexone	REFLEX
Authority Broadleaf	FirstRate	Linex	Scepter
Broadstrike	FLEXSTAR	Lorox	Sencor
Canopy	Frontier	Lorox Plus	Squadron
Canopy XL	FUSILADE	Partner	Steel
Command	FUSION	Preview	Turbo
Cover	Gemini	Prowl	WARRIOR
Dual	KARATE	Pursuit	2,4-D
Dual II	Lasso	Pursuit Plus	2,4-DB

Refer to individual product labels for precautionary statements, restrictions, rates, and a list of weeds controlled.

POSTEMERGENCE

ANNUAL WEEDS

Apply TOUCHDOWN 5 at 0.8 - 3.2 pints/acre from cracking through full flowering of soybean plants to control annual weeds. For best results, make the first application of TOUCHDOWN 5 at 1.6 pints per acre within 30 days after planting on weeds up to 6 inches tall. Refer to Table 1 and 2 for specific rate information. For any single in-crop application, use a minimum of 0.8 pints per acre of TOUCHDOWN 5. TOUCHDOWN 5 will not provide residual control. To control new weed flushes, repeat applications may be required.

PERENNIAL WEEDS

Apply TOUCHDOWN 5 at 1.6-3.2 pints per acre to actively growing perennial grasses, sedges, and broadleaf weeds. Applications in crop on glyphosate-tolerant soybeans normally occur before perennial weeds reach the most desirable growth stage for control. Treatments made prior to the timing designated in Table 4 may require retreatment. Best control will be obtained when perennial broadleaf weeds are treated in the early bud to flowering stage and when perennial grasses are in the boot to seedhead stage. Refer to Table 4 for additional rate and timing information.

TANK MIXTURES

TOUCHDOWN 5 may be tank mixed with one or more of the following products:

Basagran®	Pursuit
Classic®	Raptor™
FirstRate	REFLEX
FLEXSTAR	Reliance™ STS®
FUSILADE	Scepter
FUSION	Synchrony® STS®
KARATE	WARRIOR
Pinnacle®	2,4-DB

Use a minimum of 1.2 pints per acre TOUCHDOWN 5 in mixture with postemergent tankmix herbicides on 3 inch tall weeds. Use a minimum of 1.6 pints per acre TOUCHDOWN 5 in mixture with postemergent tankmix herbicides on 3-6 inch tall weeds. Under certain conditions, the mixture of TOUCHDOWN 5 with one or more of the above mentioned herbicides may result in a reduction of activity. Tankmixes can result in increased crop injury as compared to either product used alone. Refer to individual product labels for precautionary statements, restrictions, rates, and list of weeds controlled.

PREHARVEST

TOUCHDOWN 5 may be applied preharvest as a broadcast spray with ground or aerial equipment as a harvest aid. TOUCHDOWN 5 provides weed control when applied preharvest to soybeans. Apply to mature soybeans when pods have lost their color.

WHEAT

METHOD OF APPLICATION: Before, during, or after planting, but before crop emergence; and as a spot spray.

Follow directions listed in the **APPLICATION DIRECTIONS, SPRAY ADDITIVES,** and **APPLICATION EQUIPMENT AND TECHNIQUES** sections. Refer to **WEEDS CONTROLLED** section, beginning on Page _____, for application rates and timing.

GENERAL USE PRECAUTIONS FOR WHEAT

- Apply up to 14 days before forage and hay harvest; and up to 8 weeks before grain and straw harvest.
- Crop plants contacted by TOUCHDOWN 5 will be injured or killed.
- Do not apply more than 12.8 pints per acre per year.

TANK MIXTURES FOR WHEAT

TOUCHDOWN 5 can be tank mixed with the following herbicides for control or suppression of annual and perennial weeds.

Dicamba
Hoelon®
2,4-D

Refer to individual product labels for precautionary statements, restrictions, rates, and a list of weeds controlled.

WEEDS CONTROLLED

Water volumes of 3 to 40 gallons per acre by ground equipment and 3 - 15 gallons by air are recommended. Use the minimum spray volume that provides adequate coverage.

When tank mixing with residual herbicides, refer to the individual crop section for recommendations.

Apply to actively growing weeds.

MAP POSITIONED HERE

TABLE 1
NORTH/SOUTH REGION
ANNUAL WEED CONTROL - TOUCHDOWN 5 RATES

Use the higher end of the rate range when stressful growing conditions or dense plant populations exist.

WEED SPECIES	SCIENTIFIC NAME	TOUCHDOWN 5 RATES PINTS PER ACRE				
		MAXIMUM WEED (HEIGHT/LENGTH)				
		3"	6"	12"	18"	18"+
Barley	<i>Hordeum vulgare</i>		0.6-1.2	0.6-1.6	0.8-2.0	1.2-2.4
Barnyardgrass	<i>Echinochloa crus-galli</i>	0.8	0.8-1.6	1.6-3.2		
Black nightshade	<i>Solanum nigrum</i>	1.2	0.8-1.6	1.2-3.2		
Bluegrass, annual	<i>Poa annua</i>		0.6-0.8	0.8		
Bluegrass, bulbous	<i>Poa bulbosa</i>			0.8		
Bristly starbur	<i>Acanthospermum hispidum</i>	0.8	0.8-1.2	1.2-1.6		
Brome, downy ¹	<i>Bromus tectorum</i>	0.6	0.6-0.8	1.0		
Brome, Japanese	<i>Bromus japonicus</i>	0.6	0.8	1.0		
Buckwheat, wild ²	<i>Polygonum convolvulus</i>		1.6			
Buffalobur	<i>Solanum rostratum</i>	1.6	1.6-2.4	2.4		
Burcucumber	<i>Sicyos angulatus</i>		1.2	1.6		
Burgherkin	<i>Cucumis anguria</i>	1.2-1.6	1.6-2.4			
Buttercup ³	<i>Ranunculus</i> spp.	1.2-1.6				
Camphorweed	<i>Heterotheca subaxillaris</i>		2.4			
Carolina geranium ⁴	<i>Geranium carolinianum</i>	1.6-2.4	2.4-3.2			
Cheat	<i>Bromus secalinus</i>	0.6	0.8	1.0	1.2	
Chickweed, common	<i>Stellaria media</i>		0.6-0.8	0.8	1.2	
Chickweed, mouseear	<i>Cerastium vulgatum</i>		0.6-0.8	0.8	1.2	
Citronmelon	<i>Citrullus lanatus</i>	1.2-1.6	1.6-2.4			
Cocklebur, common	<i>Xanthium strumarium</i>		0.8	0.8-1.2	1.2-1.6	1.6-2.4
Coffee senna	<i>Cassia occidentalis</i>	1.2-1.6	1.6-2.4			
Corn ⁵	<i>Zea mays</i>			0.8	1.0-1.6	
Cowpea	<i>Vigna unguiculata</i>	1.2-1.6	1.6-2.4			
Crabgrass ⁶	<i>Digitaria</i> spp.		0.8-1.2	0.8-1.6	1.2-1.6	

Crotalaria, showy	<i>Crotalaria spectabilis</i>	0.8-1.2	1.2-1.6	1.6-2.4		
Croton, tropic	<i>Croton glandulosus</i>	1.2-1.6	1.6-2.4			
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	0.8	0.8-1.6	1.6-3.2		
Cutleaf eveningprimrose ⁴	<i>Oenothera laciniata</i>	1.6-2.4	2.4-3.2			
Fall panicum	<i>Panicum dichotomiflorum</i>	0.8-1.2	1.2-1.6	1.2-2.4		
Filaree	<i>Erodium</i> spp.			2.4		
Fleabane, annual	<i>Erigeron annuus</i>		0.8-1.2	1.2-2.4	1.2-2.4	
Fleabane, hairy	<i>Conyza bonariensis</i>		0.8-1.2	1.2-2.4	1.2-2.4	
Fleabane, rough	<i>Erigeron strigosus</i>		0.8-1.2	1.2-2.4	1.2-2.4	
Florida beggarweed	<i>Desmodium tortuosum</i>		0.8-1.2	1.2-1.6		
Foxtails	<i>Setaria</i> spp.		0.6-1.2	0.8-1.6	0.8-1.6	0.8-1.6
Goatgrass, jointed	<i>Aegilops cylindrica</i>		0.8	1.0		
Goosegrass	<i>Eleusine indica</i>	0.8	1.2	1.6-2.4		2.4-4.8
Groundsel, common	<i>Senecio vulgaris</i>		0.8			
Hemp sesbania ⁴	<i>Sesbania exaltata</i>	1.2	1.6-2.4			
Henbit ⁴	<i>Lamium amplexicaule</i>		1.6	2.4		
Hophornbeam copperleaf	<i>Acalypha ostryifolia</i>	1.6	2.4			
Horseweed/Marestail	<i>Conyza canadensis</i>		0.8-1.2	1.2-1.6	1.6	1.6-2.4
Itchgrass	<i>Rottboellia cochinchinensis</i>		0.8	1.2	1.6	
Jimsonweed	<i>Datura stramonium</i>	1.2	1.6	2.4		
Johnsongrass, seedling	<i>Sorghum halepense</i>			1.0-1.2	1.2-2.4	2.4
Kochia ³	<i>Kochia scoparia</i>	0.8	1.2-1.6	1.2-2.4		
Lambsquarters, common	<i>Chenopodium album</i>		1.2-1.6	1.6-2.4	2.4-3.2	
Lettuce, prickly	<i>Lactuca serriola</i>		0.8-1.6	1.2-2.4		1.6-2.4
Morningglory ^{4,7}	<i>Ipomoea</i> spp.	1.2-2.4	1.6-3.2			
Mustard, blue	<i>Chorispora tenella</i>	0.6-0.8	0.6-0.8	1.0		1.2
Mustard, tansy	<i>Descurainia pinnata</i>	0.6-0.8	0.6-1.0			1.2
Mustard, tumble	<i>Sisymbrium altissimum</i>	0.6-0.8	0.6-0.8	1.0		1.2
Mustard, wild	<i>Brassica kaber</i>	0.6-0.8	0.6-0.8	0.8		1.2
Oats	<i>Avena sativa</i>		0.6-1.2	0.6-1.6	0.8-2.0	1.2-2.4

Oats, wild	<i>Avena fatua</i>		0.6-1.2	0.8-1.2		
Panicum, Texas	<i>Panicum texanum</i>		0.8-1.2	1.6-2.4		
Pennycress, field	<i>Thlaspi arvense</i>		0.6-1.2	1.0-1.6		
Pigweed	<i>Amaranthus</i> spp.		0.6-0.8	0.8-1.0		1.6
Poinsettia, wild	<i>Euphorbia heterophylla</i>	0.8-1.2	1.6-2.4			
Prickly sida (Teaweed) ^{4,7}	<i>Sida spinosa</i>	1.2-2.0	2.4-3.2			
Purslane, common	<i>Portulaca oleracea</i>	1.2-1.6	1.6-2.4			
Pusley, Florida	<i>Richardia scabra</i>	1.2-1.6	1.6-2.4	2.4-3.2		
Ragweed, common	<i>Ambrosia artemisiifolia</i>		0.8-1.2	1.2-3.2		
Ragweed, giant	<i>Ambrosia trifida</i>	1.2	1.6	1.6-2.4	1.6-3.2	2.4-3.2
Red rice	<i>Oryza sativa</i>	1.6	2.4-3.2			
Redweed	<i>Melochia corchorifolia</i>	1.2-1.6	1.6-2.4			
Rocket, London	<i>Sisymbrium irio</i>	0.6	0.6-0.8	0.8-1.0		
Rye	<i>Secale cereale</i>		0.6-1.2	0.8-1.6	1.2-2.0	1.6-2.4
Ryegrass, Italian	<i>Lolium multiflorum</i>		1.6	2.4-3.2		
Sandbur, field	<i>Cenchrus incertus</i>			0.6		
Sandbur, southern	<i>Cenchrus echinatus</i>	0.8	1.2-1.6	1.6-2.4		
Shattercane	<i>Sorghum bicolor</i>			0.8-1.0	1.2	1.2
Shepherdspurse	<i>Capsella bursa-pastoris</i>		0.8-1.0	1.2-1.6		
Sicklepod	<i>Cassia obtusifolia</i>	1.6	2.4-3.2			
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>	0.8	1.0-1.6	1.6-3.2		
Smartweed (ladysthumb)	<i>Polygonum persicaria</i>	0.8-1.0	1.2-1.6	1.6-2.4		
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	0.8-1.0	1.2-1.6	1.6-2.4		
Sowthistle, annual	<i>Sonchus oleraceus</i>		1.6	2.4		
Spanishneedles	<i>Bidens bipinnata</i>		1.6	1.6-2.4	2.4	
Sprangletop	<i>Leptochloa</i> spp.		0.8	1.2		1.6
Spurge	<i>Euphorbia</i> spp.		0.8	1.2		1.6
Stinkgrass	<i>Eragrostis cilianensis</i>	0.6-1.0	0.8-1.0			
Sunflower, common	<i>Helianthus annuus</i>		0.8-1.0	1.0-1.2	1.2-1.6	2.4
Thistle, Russian	<i>Salsola iberica</i>	0.8-1.0	1.2-1.6	1.6-2.4		
Velvetleaf ⁷	<i>Abutilon theophrasti</i>	0.8-1.2	1.2-1.6	1.6-2.4	2.0-3.2	
Virginia copperleaf	<i>Acalypha virginica</i>	1.6	2.4			

Waterhemp	<i>Amaranthus</i> spp.	1.2-1.6	1.2-1.6	1.6-2.0	2.4	
Wheat	<i>Triticum aestivum</i>		0.6-1.2	0.6-1.6	0.8-2.0	1.2-2.4
Witchgrass	<i>Panicum capillare</i>			0.8		
Woolly cupgrass	<i>Eriochloa villosa</i>	0.8-1.2	0.8-1.6	1.2-1.6		

- ¹ In no-till systems, use 1 pint/A.
- ² Maximum runner length.
- ³ Control will be reduced at the button stage.
- ⁴ When the predominant weed species include Carolina geranium, cutleaf eveningprimrose, fescue, hemp sesbania, henbit, morningglory, prickly sida, and vetch that are less than 6 inches tall, GRAMOXONE EXTRA should be considered as an alternative.
- ⁵ Will not control glyphosate-tolerant volunteer corn.
- ⁶ Plant diameter.
- ⁷ Multiple applications may be required.

TABLE 2
WESTERN REGION
ANNUAL WEED CONTROL - TOUCHDOWN 5 RATES

Use the higher end of the rate range when stressful growing conditions or dense plant populations exist.

WEED SPECIES	SCIENTIFIC NAME	TOUCHDOWN 5 RATES PINTS PER ACRE			
		MAXIMUM WEED (HEIGHT/LENGTH)			
		3"	6"	12"	18"
Barley	<i>Hordeum vulgare</i>		0.6-1.2	0.6-2.0	0.6-2.4
Barnyardgrass	<i>Echinochloa crus-galli</i>		0.6-1.6		
Bluegrass, annual	<i>Poa annua</i>		0.6-1.6		
Bluegrass, bulbous	<i>Poa bulbosa</i>		0.8-2.0		
Brome, downy ¹	<i>Bromus tectorum</i>		0.6-1.6		
Buttercup	<i>Ranunculus</i> spp.			0.8-2.0	
Canarygrass	<i>Phalaris canariensis</i>		0.8-2.0		
Cheat	<i>Bromus secalinus</i>		0.8-2.0		
Chickweed, common	<i>Stellaria media</i>		0.8-2.0		
Chickweed, mouseear	<i>Cerastium vulgatum</i>		0.8-2.0		
Cocklebur, common	<i>Xanthium strumarium</i>			0.8-2.0	
Corn ²	<i>Zea mays</i>		0.8-2.0		
Crabgrass	<i>Digitaria</i> spp.			0.8-2.0	
Dwarf dandelion, Virginia	<i>Krigia virginica</i>			0.8-2.0	
Fall panicum	<i>Panicum dichotomiflorum</i>			0.8-2.0	
Falseflax, smallseed	<i>Camelina microcarpa</i>			0.8-2.0	
Fiddleneck	<i>Amsinckia</i> spp.		0.8-2.0		
Filaree	<i>Erodium</i> spp.			2.4-6.4	
Fleabane, hairy	<i>Conyza bonariensis</i>		0.8-2.0		
Foxtail	<i>Setaria</i> spp.		0.4-1.2	0.4-2	
Goatgrass, jointed	<i>Aegilops cylindrica</i>		0.8-2.0		
Goosefoot, nettleleaf	<i>Chenopodium murale</i>		0.8-2.0		
Groundcherry	<i>Physalis</i> spp.		1.6-2.4		

Groundsel, common	<i>Senecio vulgaris</i>		0.8-2.0		
Henbit	<i>Lamium amplexicaule</i>		0.8-2.0		
Horseweed/Marestail	<i>Conyza canadensis</i>		0.8-2.0		
Johnsongrass, seedling	<i>Sorghum halepense</i>			0.8-2.0	
Junglerice	<i>Echinochloa colonum</i>		0.8-2.0		
Kochia ³	<i>Kochia scoparia</i>	0.8-1.2	1.2-1.6		
Lambsquarters, common	<i>Chenopodium album</i>		0.8-2.0		
Morningglory ^{4,5}	<i>Ipomoea</i> spp.	0.6-2.0			
Mustard, blue	<i>Chorispora tenella</i>		0.6-1.6		
Mustard, tansy	<i>Descurainia pinnata</i>		0.6-1.6		
Mustard, tumble	<i>Sisymbrium altissimum</i>		0.6-1.6		
Mustard, wild	<i>Sinapis arvensis</i>		0.6-1.6		
Nightshade	<i>Solanum</i> spp.		1.6-2.4		
Oats	<i>Avena sativa</i>		0.6-1.2	0.6-2.0	0.6-2.4
Oats, wild	<i>Avena fatua</i>			0.8-2.0	
Panicum, Texas	<i>Panicum texanum</i>			0.8-2.0	
Pennycress, field	<i>Thlaspi arvense</i>		0.8-2.0		
Pigweed	<i>Amaranthus</i> spp.			0.8-2.0	
Puncturevine	<i>Tribulus terrestris</i>	0.8-1.6	1.2-2.4		
Pusley, Florida	<i>Richardia scabra</i>			1.6-3.2	
Rabbitfootgrass	<i>Polypogon monspeliensis</i>		0.8-2.0		
Rocket, London	<i>Sisymbrium irio</i>		0.8-2.0		
Rockpurslane, Redmaids	<i>Calandrinia caulescens</i>		0.8-2.0		
Rye	<i>Secale cereale</i>		0.6-1.2	0.8-1.6	1.2-2.0
Ryegrass, Italian/annual	<i>Lolium multiflorum</i>		0.8-2.0		
Sandbur, field	<i>Cenchrus incertus</i>			0.6-1.6	
Shattercane	<i>Sorghum bicolor</i>			0.6-1.6	
Shepherdspurse	<i>Capsella bursa-pastoris</i>		0.8-2.0		
Sowthistle, annual	<i>Sonchus oleraceus</i>		0.8-2.0		

Sprangletop	<i>Leptochloa</i> spp.		1.6-2.4		
Spurge	<i>Euphorbia</i> spp.		0.8-2.0		
Spurge, prostrate	<i>Euphorbia supina</i>		1.6-2.4		
Stinkgrass	<i>Eragrostis cilianensis</i>			0.6-1.6	
Sunflower, common	<i>Helianthus annuus</i>		0.8-1.2	1.2-1.6	
Waterhemp	<i>Amaranthus</i> spp.	1.2-1.6	1.2-1.6	1.6-2.0	2.4
Wheat	<i>Triticum aestivum</i>		0.6-1.2	0.6-2.0	0.6-2.4
Witchgrass	<i>Panicum capillare</i>			0.8-2.0	

¹ In no-till systems, use 0.8-1.6 pints/A.

² Will not control glyphosate-tolerant volunteer corn.

³ Do not apply in the button stage.

⁴ Multiple applications may be required.

⁵ When the predominate weed species include morningglory less than six inches tall, GRAMOXONE EXTRA should be considered as an alternative.

TABLE 3
ANNUAL WEED CONTROL - TOUCHDOWN 5 RATES IN A TANKMIX
WITH 0.25 LBS A/A OF DICAMBA OR 0.5 LBS A/A OF 2,4-D

WEED SPECIES	SCIENTIFIC NAME	MAXIMUM HEIGHT/LENGTH	TOUCHDOWN 5 PINTS PER ACRE
Lettuce, prickly Morningglory Ragweed, common Ragweed, giant Smartweed, Pennsylvania Velvetleaf	<i>Lactuca serriola</i> <i>Ipomoea</i> spp. <i>Ambrosia artemisiifolia</i> <i>Ambrosia trifida</i> <i>Polygonum pennsylvanicum</i> <i>Abutilon theophrasti</i>	6"	0.8
Cocklebur, common Fleabane, rough Horseweed/Marestail Kochia Lambsquarters, common Sunflower, common Thistle, Russian	<i>Xanthium strumarium</i> <i>Erigeron strigosus</i> <i>Conyza canadensis</i> <i>Kochia scoparia</i> <i>Chenopodium album</i> <i>Helianthus annuus</i> <i>Salsola iberica</i>	12"	

Read and follow dicamba and 2,4-D labels

TABLE 4
PERENNIAL AND WOODY PLANT WEED CONTROL AND WEED MANAGEMENT
TOUCHDOWN 5 RATES USED ALONE OR IN TANKMIX WITH
0.25 LBS A/A OF DICAMBA OR 0.5 LBS A/A OF 2,4-D

WEED SPECIES	SCIENTIFIC NAMES	SPOT SPRAY % v/v	PINTS PER ACRE	TANKMIX with 2,4-D or Dicamba	APPLICATION TIMING AND REMARKS
Alfalfa	<i>Medicago sativa</i>	1.5	2.4-4.8		At 6 to 8" stage or more after final cutting in fall. Deep till 7 days after treatment.
			0.8-1.6	Yes	At 6 to 8" stage or more for suppression.
Artichoke, Jerusalem ¹	<i>Helianthus tuberosus</i>	1.5	4.8-6.4		At or after flowering.
Balsamapple	<i>Momordica charantia</i>	1.5			For partial control only, apply at or beyond bloom.
Bahiagrass	<i>Paspalum notatum</i>	1.5	4.8-6.4		Early seedhead stage.
Barley, foxtail	<i>Hordeum jubatum</i>	1.5	1.6-3.2		4-6" stage.
Bermudagrass	<i>Cynodon dactylon</i>	1.5	3.2-6.4		Seedheads present, may require retreatment.
Bindweed, field	<i>Convolvulus arvensis</i>	1.5	3.2-6.4		At or after flowering, requires 2 applications.
			0.8-1.6	Yes	At 8-12" stage or more for suppression.
			1.6	Yes	At or after flowering for control, multiple applications may be required.
			1.6-6.4		In California: 12" or greater runner length. Use high end of rate range where dense populations exist. For suppression on land which is irrigated and tilled, use 1.6 pints/A.
Blackberry	<i>Rubus</i> spp.	1.0	4.8-6.4		Apply after full leaf expansion. For best results, apply late summer or fall but before a killing frost.

Bluegrass, Kentucky	<i>Poa pratensis</i>	1.5	3.2-4.8		Boot to early seedhead stage.
			1.6-2.4		For suppression.
Brackenfern	<i>Pteridium aquilinum</i>	1.25	4.8-6.4		Fronks fully expanded and at least 18" long.
Bromegrass, smooth	<i>Bromus inermis</i>	1.5	1.6-3.2		Boot to early seedhead
			0.8		For suppression.
Canarygrass, reed	<i>Phalaris arundinacea</i>	1.5	3.2-4.8		Boot to head.
Cattail	<i>Typha sp.</i>	1.5	4.8-6.4		Early head to early bud.
Clover, red Clover, white	<i>Trifolium pratense</i> <i>Trifolium repens</i>	1.5	4.8-6.4		Early head to early bud. May require retreatment.
			0.8-1.6		Yes
Cogongrass ¹	<i>Imperata cylindrica</i>	1.5	4.8-6.4		Late summer/fall greater than 18" in height. May require retreatment.
Dandelion	<i>Taraxacum officinale</i>	1.5	2.4-4.8		Early bud.
			0.8		Yes
Dayflower ¹	<i>Commelina spp.</i>	1.5	4.8-6.4		Less than 4" in height.
Dock, curly	<i>Rumex crispus</i>	1.5	3.2-4.8		Early bud.
			0.8-1.6		Yes

Dogbane, hemp	<i>Apocynum cannabinum</i>	1.5	3.2-6.4		Late bud to flower. May require retreatment.
			0.8-3.2	Yes	Actively growing at 6-12" stage for suppression.
			1.6	Yes	Actively growing, late bud to flower.
Dogfennel	<i>Eupatorium capillifolium</i>	1.5	3.2-4.8		Actively growing, less than 12" in height.
Fescue ²	<i>Festuca</i> spp.	1.5	4.8-6.4		Boot to early seedhead.
Fescue, tall	<i>Festuca arundinacea</i>	1.5	1.6-4.8		For fall application on 8 to 12" plants, use 1.6-2.4 pints/A. Use 4.8 pints/A on CRP sodded stands at boot or later.
Goatweed	<i>Scoparia dulcis</i>	2.0	4.8-6.4		Less than 7" stage.
Guineagrass	<i>Panicum maximum</i>	1.0	4.8		7-10 leaf stage.
Horsenettle	<i>Solanum carolinense</i>	1.5	3.2-6.4		Late bud to flower.
			1.6-3.2	Yes	Late bud to flower.

Johnsongrass	<i>Sorghum halepense</i>	1.0	1.6-4.8		Apply at boot to head stage and in the fall prior to frost. Use 1.6-3.2 pints/A for annual tillage systems. Use 3.2-4.8 pints/A on no-till areas. Allow 3-7 days before tillage.
			0.8-1.6		For burndown, apply when plants are 10-12" stage and allow 3 days before tillage.
Lantana, largeleaf	<i>Lantana camara</i>	1.0			For partial control, apply at or beyond bloom stage.
Milkweed, common	<i>Asclepias syriaca</i>	1.5	3.2-4.8		Late bud to early flower. May require retreatment.
			1.6	Yes	Late bud to early flower.
			0.8-1.6	Yes	Suppression.
Milkweed, honeyvine	<i>Ampelamus albidus</i>	1.5	3.2-4.8	Yes	Late bud to early flower. May require retreatment.
Muhly, wirestem	<i>Muhlenbergia frondosa</i>	1.5	1.6-3.2		6-8" actively growing. Use high end of rate range in non-tillage areas.
Mullein, common	<i>Verbascum thapsus</i>	1.5	3.2-6.4		Early bud to head.
Nightshade, silverleaf	<i>Solanum eleagnifolium</i>	1.5	1.6-3.2		Apply when 60% of plants have berries. Apply fall treatments before a killing frost.
Nutsedge, purple Nutsedge, yellow	<i>Cyperus rotundus</i> <i>Cyperus esculentus</i>	1.5	1.6-4.8		Treat when most plants are in flower stage. Repeat applications may be required for dense populations of nutsedge and subsequent germination of dormant nuts. Allow 5" of regrowth before retreatment. Use 0.8-1.6 pints/A for suppression. A second application will be required on recovered plants with 5" of new growth.
Orchardgrass	<i>Dactylis glomerata</i>	1.5	2.4-3.2		Apply 3.2 pints/A on plants at early boot to seedhead stage. For spring applications, use 3.2 pints/A and allow a minimum height of 10-12". Fall applications may be made on actively growing plants at the boot to early seedhead stage at 2.4-3.2 pints/A.
Pampasgrass	<i>Erianthus ravennae</i>	1.5			For partial control only; apply at or beyond boot stage.
Paragrass	<i>Brachiaria mutica</i>	1.5	2.4-3.2		Early seedhead stage.
Phaseybean ¹	<i>Phaseolus lathyroides</i>	1.5	4.8-6.4		Less than 8" tall.

Poison ivy ¹	<i>Toxicodendron radicans</i>				Apply after full leaf expansion. For best results, apply late summer or fall but before leaves lose green color.
Poison oak ¹	<i>Toxicodendron toxicarium</i>	1.5	6.4		
Quackgrass	<i>Agropyron repens</i>	1.5	1.6-3.2		Apply at 6-10" stage in annual cropping systems followed by tillage. Use the higher rate on dense established stands. Allow 3 days before tillage.
			3.2-4.8		Apply at 6-10" stage if tillage is not performed following application.
Redvine ¹	<i>Brunnichia ovata</i>	1.5	0.8-2.4		Greater than 18" tall in September/October; retreatment will be required at rates less than 1.6 pint/A. Allow 12" of regrowth before retreatment.
Ryegrass, perennial	<i>Lolium perenne</i>	1.0	1.6-4.8		Boot to head or prior to frost; use higher end of rate range if no tillage will be performed following application.
Smallflowered Alexandergrass	<i>Brachiaria subquadrifera</i>	1.5	3.2-4.8		Less than 4" in height, actively growing.

Smartweed, swamp	<i>Polygonum coccineum</i>	1.5	2.4-6.4		Early bud, 12" stage
			0.8	Yes	Early bud, 12" stage.
Spurge, leafy	<i>Euphorbia esula</i>	1.5	3.2		Greater than 12" tall in late summer or fall.
			0.8	Yes	Greater than 12" tall in late summer.
Sumac, dwarf ¹ Sumac, poison ¹ Sumac, smooth ¹	<i>Rhus copallina</i> <i>Toxicodendron vernix</i> <i>Rhus glabra</i>	1.5	3.2-6.4		Apply after full leaf expansion. For best results, apply late summer or fall but before leaves lose green color.
Sweet potato, wild	<i>Ipomea pandurata</i>	1.5			For partial control only, apply at or beyond flowering stage.
Switchgrass	<i>Panicum virgatum</i>	1.5	2.4-3.2		Boot to head stage.
Thistle, Canada	<i>Cirsium arvense</i>	1.5	3.2-4.8		Apply at bud to early flower stage. For fall applications or following mowing, allow a minimum of 6-8" rosette development.
Timothy	<i>Phleum pratense</i>	1.5	3.2-4.8		Boot to head; wait 3 days before tillage.
Torpedograss ¹	<i>Panicum repens</i>	1.5	6.4		At or beyond seedhead for suppression.
Trumpet creeper ¹	<i>Campsis radicans</i>	1.5	3.2-4.8		Late September/October applications on actively growing plants, retreatment may be required.
Vaseygrass	<i>Paspalum urvillei</i>	1.5	4.8-6.4		Apply at early head stage.
Vetch ²	<i>Vicia</i> spp.	1.5	3.2-4.8		Boot to head.
Virginia creeper	<i>Parthenocissus quinquefolia</i>	1.5	6.4		Full leaf expansion.
Wheatgrass, Western	<i>Agropyron smithii</i>	1.5	3.2-4.8		Boot to head.

¹ Partial control.² When the predominant weed species include Carolina geranium, cutleaf eveningprimrose, fescue, hemp sesbania, henbit, morningglory, prickly sida, and vetch that are less than 6 inches tall, GRAMOXONE EXTRA should be considered as an alternative.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified. Country: United States of America. U.S. Patent No. 4,315,765.

AMBUSH®, BICEP®, BICEP II®, BICEP II MAGNUM™, DEVRINOL®, DUAL®, DUAL II®, DUAL II MAGNUM™, FLEXSTAR®, FULTIME™, FUSILADE®, FUSION®, GRAMOXONE®, KARATE®, PRINCEP®, PRINCEP® CALIBER 90®, REFLEX®, SOLICAM®, TOUCHDOWN®, and WARRIOR® Registered trademarks of a Syngenta Group Company.

Authority™, and Command® Registered trademarks of FMC Corporation.

Axiom™, Sencor®, and Turbo® Registered trademarks of Bayer.

Balance® Registered trademark of Rhone-Poulenc Ag Company.

Basagran®, Clarity®, Frontier®, Guardsman®, and Marksman® Registered trademarks of BASF Corporation.

Basis®, Bladex®, Canopy®, Classic®, Cover™, Extrazine® II, Gemini®, Karmex®, Krovar®, Lexone®, Lorox®, Pinnacle®, Preview®,

Reliance™, Sinbar®, and Synchrony® Registered trademarks of E. I. du Pont de Nemours & Co., Inc.

Broadstrike®, FirstRate™, Hornet™, Surflan®, Surpass®, and Topnotch® Registered trademarks of Dow AgroSciences.

Bullet®, Harness®, Lariat®, Lasso®, Micro-Tech®, Partner®, and ROUNDUP READY® Registered trademarks of Monsanto Company.

Direx® and Linex® Registered trademarks of Griffin Corporation.

Goal® and Kerb® Registered trademarks of Rohm and Haas Company.

Hoelon® Registered trademark of AgrEvo USA Company.

Lightning™, Prowl®, Pursuit®, Raptor™, Scepter®, Squadron®, and Steel™ Registered trademarks of American Cyanamid Company.

Sim-Trok® Registered trademark of Oxon Italia.