

	United States	<input type="checkbox"/> Registration	OPP Identifier Number NOTIFICATION
	Environmental Protection Agency	<input type="checkbox"/> Amendment	
	Washington, DC 20460	<input checked="" type="checkbox"/> Other	

Application for Pesticide - Section I

1. Company/Product Number 100-1112	2. EPA Product Manager George LaRocca	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Warrior Insecticide with Zeon Technology	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) Syngenta Crop Protection, Inc. P. O. Box 18300 Greensboro, NC 27419 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

NOTIFICATION

DEC 20 2001

Explanation: Use additional page(s) if necessary. (For Section I and Section II.).
Submission of Final Printed Labels Per PR-Notice 98-10

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. The following changes are being made via this notification: 1) Company name and address have been updated to reflect Syngenta Crop Protection, Inc. 2) The Conditions of Sale and Warranty statement has been changed to reflect the name change. Because Syngenta has been formed by the merger of Novartis Crop Protection, Inc. and Zeneca Ag Products, we have chosen to use the former Zeneca warranty statement as the Syngenta warranty statement. No other changes occur in the statement other than the name change. 3) The copyright date reflects Syngenta. 4) Trademark statements have been updated to reflect Syngenta for those products for which Syngenta holds the trademark. 5) The Internet address has been changed to reflect Syngenta. 6) Other places in the label which referring to the company name have been updated.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No			
*Certification must be submitted			If "Yes" No. per Unit Packaging wgt. Container		
			If "Yes" No. per Unit Packaging wgt. container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					


Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Fred J. Pearson	Title NAFTA Insecticide Team Leader	Telephone No. (Include Area Code) 336-632-2365	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Data Application Received (Stamped)
2. Signature 	3. Title NAFTA Insecticide Team Leader		
4. Typed Name Fred J. Pearson	5. Date December 6, 2001		

2/23

PULL HERE TO OPEN ►

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS
 FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR
 DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED
 APPLICATOR'S CERTIFICATION.

WARRIOR®


Insecticide

Active Ingredient:
 Lambda-cyhalothrin:
 [1 α (S*), 3 α (Z)]-(+)-cyano-(3-phenoxyphenyl)methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-
 2,2-dimethylcyclopropanecarboxylate 11.4%

Other Ingredients: 88.6%

Total: 100.0%

Warrior Insecticide with Zeon Technology contains 1 lb. of active ingredient per gallon
 and is a capsule suspension.

KEEP OUT OF REACH OF CHILDREN.
SHAKE WELL BEFORE USING.
WARNING/AVISO


*Si usted no entiende la etiqueta, busque a alguien para que se la explique
 a usted en detalle. (If you do not understand the label, find
 someone to explain it to you in detail.)*

EPA Reg. No. 100-1112 EPA Est. 070989-AR-001

Product of the United Kingdom
 Formulated in the USA

SCP 1112A-L1 0801

1 gallon
 U.S. Standard
 Measure



NOTIFICATION

DEC 20 2001

12/05/01

Warrior® with Zeon™ Technology

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a physician or poison control center. • Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. • If person is unconscious, do not give anything by mouth and do not induce vomiting.
If on skin	<ul style="list-style-type: none"> • Get medical attention if irritation persists. • Wash with plenty of soap and water.
If in eyes	<ul style="list-style-type: none"> • Hold eyelids open and flush with steady, gentle stream of water. • Remove any contact lenses and repeat flushing procedure for several minutes. • Get medical attention if irritation persists.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals****WARNING**

May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. May cause allergic skin reactions. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2-30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber or viton > 14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitations of Warranty and of Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE**Restricted Use Pesticide**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber or viton > 14 mils
- Shoes plus socks

STORAGE AND DISPOSAL**Prohibitions**

Do not contaminate water, food, or feed by storage and disposal.

Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area. DO NOT ALLOW PRODUCT TO FREEZE.

Warrior® with Zeon™ Technology

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent): then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

For Bulk, Mini-Bulk, EZ Handler® and Boomerang Container Disposal

Return container to Syngenta for reuse with seal intact and in salable condition.

Container Precautions

Before refilling RETURNABLE CONTAINERS, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices.

REFILL ONLY WITH WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY: The contents of RETURNABLE CONTAINERS cannot be completely removed by cleaning. Refilling with materials other than Warrior® Insecticide with Zeon™ Technology will result in contamination and may weaken container.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals./A by air or 10 gals./A by ground unless otherwise specified in this label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, Warrior® Insecticide with Zeon Technology may be applied before, during, or after planting. For soil incorporated applications, use higher rates for improved control.

Resistance

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

SPRAY DRIFT PRECAUTIONS

Observe the Following Precautions When Spraying in the Vicinity of Aquatic Areas Such as Lakes; Reservoirs; Rivers; Permanent Streams, Marshes or Natural Ponds; Estuaries and Commercial Fish Farm Ponds:

- Do not apply by ground within 25 ft., or by air within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, or natural ponds; estuaries and commercial fish farm ponds. Increase the buffer zone to 450 ft. when ultra-low volume (ULV) application is made.
- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.
- Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 ft. above the crop canopy should be avoided.
- Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3 – 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.
- Risk of exposure to aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
- Do not cultivate within 10 ft. of the aquatic area so as to allow growth of a vegetative filter strip.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

6/23

Warrior® with Zeon™ Technology

- Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

In the State of New York, a 25 ft. vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for both aerial or ground application. For aerial applications, the 25 ft. vegetated non-cropped buffer strip for runoff protection would be part of the larger 150 ft. buffer strip (or 450 ft. buffer strip for ULV application) required for spray drift.

TANK MIX APPLICATION

When tank mixing with any other agricultural products, always add Warrior Insecticide with Zeon Technology last. Fill the tank with one half to two thirds volume of the mixing diluent. Make sure all other products are fully dispersed in the mixing diluent before adding the recommended rate of Warrior Insecticide with Zeon Technology to the tank. Add the remainder of the mixing diluent volume. It is recommended that mixing and spray equipment have continuous agitation for best results. Follow the precautions and limitations of the most restricted product in the tank mixture.

While Warrior Insecticide with Zeon Technology has good flexibility for tank mixing with other agricultural products, a jar test for physical compatibility is recommended for untried mixtures using proper ratios and mixing sequences of all ingredients to be included in the mixture.

Warrior Insecticide with Zeon Technology is an aqueous based formulation. It is recommended that no type of non-emulsifiable oils be used in combination with Warrior Insecticide with Zeon Technology. If adjuvants are used, use only:

Nonionic Surfactant (NIS) containing at least 75% surface agent or Non-phytotoxic Crop Oil Concentrate (COC), including once refined Vegetable Oil Concentrate (VOC), or Methylated Sunflower Oils (MSO) containing a minimum of 17% emulsifier.

Adjuvants other than NIS or COC may be used providing the product meets the following criteria:

1. Contains only EPA exempt ingredients.
2. Is non-phytotoxic to the target crop.
3. Is compatible in mixture. (May be established through a jar test.)
4. Is supported locally for use with Warrior Insecticide with Zeon Technology on the target crop through proven field trials and through university and extension recommendations.

In addition, the following may be used as diluents:

Crop Oil Concentrate
Methylated Sunflower Oils
Urea-Ammonium Nitrate

It is recommended that the following not be used in combination with Warrior Insecticide with Zeon Technology as diluents or adjuvants:

Non-emulsifiable oils
Diesel Fuel
Straight Mineral Oil

CHEMIGATION

Sprinkler Irrigation Application

Apply Warrior Insecticide with Zeon Technology at rates and timing described elsewhere in this label. As local recommendations differ, consult your local State Extension Service or other local experts for recommendations on adjuvant or diluent types, (see Tank Mix Application) rates and mixing instructions. These recommendations should be proven, through university and extension field trials, to be effective with Warrior Insecticide with Zeon Technology applied by chemigation.

Check the irrigation system to insure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period.

Apply by injecting the recommended rate of Warrior Insecticide with Zeon Technology into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1-0.2 A/in. of water. In general, use the least amount of water required for proper distribution and coverage. It is recommended that the product be injected into the main irrigation line ahead of a right angle turn in the line to insure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the recommended rate of Warrior Insecticide with Zeon Technology for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

It is not recommended that Warrior Insecticide with Zeon Technology be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Warrior® with Zeon™ Technology**Use Precautions - Sprinkler Irrigation Application**

- A. Apply this product only through (sprinkler including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Do not apply through chemigation systems connected to public water systems.

8/23

Warrior® with Zeon™ Technology

Apply Warrior Insecticide with Zeon Technology as shown in the following charts:

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Alfalfa, Alfalfa grown for seed	Alfalfa Caterpillar Cutworm Spp. Green Cloverworm Looper Spp. Velvetbean Caterpillar Webworm Spp. Leafhopper Spp. Threecornered Alfalfa Hopper	0.015–0.025	1.92–3.20	<ul style="list-style-type: none"> • Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals./A by air or 10 gals./A by ground. When foliage is dense and/or pest populations are high 5–10 gals./A by air or 20 gals./A by ground and higher use rates are recommended. Use higher rates for increased residual control. • Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2–3 days following application. Avoid direct application to bee shelters. • Do not apply more than 0.03 lb. a.i. (0.24 pts.)/A per cutting. Do not apply more than 0.12 lb. a.i. (0.96 pts.)/A per season. Do not apply within 1 day of harvest for forage or within 7 days of harvest for hay.
	Armyworm Corn Earworm Fall Armyworm ¹ Western Yellow-striped Armyworm Yellow-striped Armyworm Alfalfa Weevil Bean Leaf Beetle (Adult) Blister Beetle Spp. Clover Leaf Weevil Spp. Clover Root Borer (Adult) Clover Root Curculio Spp. (Adult) Clover Stem Borer (Adult) Cowpea Curculio (Adult) Cowpea Weevil (Adult) Cucumber Beetle Spp. (Adult) Egyptian Alfalfa Weevil Grape Colaspis (Adult) Green June Beetle (Adult) Japanese Beetle (Adult) Mexican Bean Beetle Pea Weevil (Adult) Sweet Clover Weevil (Adult) Whitefringed Beetle Spp. (Adult) Meadow Spittlebug Plant Bug Spp. including Lygus Spp. ³ Stink Bug Spp. Alfalfa Seed Chalcid (Adult) Blue Alfalfa Aphid Cowpea Aphid Green Peach Aphid ³ Pea Aphid Spotted Alfalfa Aphid Thrips Spp. Grasshopper Spp.	0.02–0.03	2.56–3.84	
	Beet Armyworm ^{1,3} Blotch Leafminer ³ Spider Mites ²	0.03	3.84	

¹For control of first and second instar only.

²Suppression only.

³See resistance statement under General Directions for Use.

Warrior® with Zeon™ Technology

9/23

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Broccoli	Alfalfa Looper	0.015–0.025	1.92–3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
Brussels Sprouts	Cabbage Looper			
Cabbage	Imported Cabbageworm			
Cavalo	Southern Cabbageworm			
Broccoli	Cutworm Spp.	0.02–0.03	2.56–3.84	<ul style="list-style-type: none"> • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A. • Do not apply within 1 day of harvest. • Do not apply more than 0.24 lb. a.i. (1.92 pts./A) per season.
Cauliflower	Cabbage Webworm			
Chinese Broccoli (gai lon)	Diamondback Moth ³			
Chinese Armyworm	Beet Armyworm ^{1,3}			
Cabbage (napa),	Fall Armyworm ¹			
Chinese Yellow-striped	Armyworm			
Mustard	Corn Earworm			
Cabbage (gai choy)	Flea Beetle Spp.			
Kohlrabi	Japanese Beetle (Adult)			
	Vegetable Weevil (Adult)			
	Grasshopper Spp.			
	Leafhopper Spp.			
	Plant Bug Spp. including Lygus Spp. ³			
	Stink Bug Spp.			
	Meadow Spittlebug			
	Aphid Spp. ^{2,3}			
	Whitefly Spp. ^{2,3}			
	Thrips Spp. ²			
	Spider Mite Spp. ²			

¹For control of first and second instar only.

²Suppression only.

³See resistance statement under General Directions for Use.

Warrior® with Zeon™ Technology

10/23

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Corn Field Corn Popcorn Seed Corn	Cutworm Spp. Western Bean Cutworm ¹ Corn Earworm ¹ Green Cloverworm Meadow Spittlebug	0.015–0.025	1.92–3.20	<ul style="list-style-type: none"> • Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods. • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gals. of water/A. • For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3- to 5-day intervals if needed. Warrior Insecticide with Zeon Technology may only suppress heavy infestations and/or subsequent migrations. • For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program, use a minimum of 3.84 fl. oz./A (0.03 lb. a.i./A). • Do not apply within 21 days of harvest. • Do not apply more than 0.12 lb. a.i. (0.96 pts.)/A per season. Do not apply more than 0.06 lb. a.i. (0.48 pts.) after silk initiation. Do not apply more than 0.03 lb. a.i. (0.24 pts.) after corn has reached the milk stage (yellow kernels with milky fluid).
	Tobacco Budworm ^{1,4} European Corn Borer ¹ Southwestern Corn Borer ¹ Stalk Borer ¹ Hop Vine Borer ¹ Armyworm ² Fall Armyworm ² Yellow-striped Armyworm ² Webworm Spp. Flea Beetle Spp. Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Bean Leaf Beetle Cereal Leaf Beetle Japanese Beetle (Adult) Sap Beetle (Adult) Stink Bug Spp. Grasshopper Spp. Corn Leaf Aphid ³ Bird Cherry-Oat Aphid ³ English Grain Aphid ³	0.02–0.03	2.56–3.84	
	Beet Armyworm ^{2,4} Chinch Bug Green Bug ^{3,4}	0.03	3.84	<p>¹For control before the larva bores into the plant stalk or ear.</p> <p>²For control of first and second instar only.</p> <p>³Suppression only.</p> <p>⁴See resistance statement under General Directions for Use.</p>

Warrior® with Zeon™ Technology

11/23

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Sweet Corn	Corn Earworm Fall Armyworm ¹ Southern Armyworm ¹ Beet Armyworm ^{1,3} Yellow-Striped Armyworm ¹ Cutworm Spp. Western Bean Cutworm Webworm Spp. European Corn Borer Southwestern Corn Borer Common Cornstalk Borer Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Japanese Beetle (Adult) Sap Beetle (Adult) Flea Beetle Spp. Tarnished Plant Bug Stink Bug Spp. Chinch Bug Aster Leafhopper Grasshopper Spp. Aphid Spp. ^{2,3} Spider Mite Spp. ²	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 4 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods and should be targeted for control before insects enter the stalk or ear. • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage and ears (if present). When applying by air, apply in a minimum of 2 gals. of water/A. • For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program use a minimum of 3.2 fl. oz./A (0.025 lb. a.i./A). • Do not apply within 1 day of harvest. • Do not apply more than 0.48 lb. a.i. (3.84 pts.)/A per season.
	Corn Silkfly (Adult) ²	0.03	3.84	¹ For control of first and second instar only. ² Suppression only. ³ See resistance statement under General Directions for Use.

12/23

Warrior® with Zeon™ Technology

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Cotton	Cutworm Spp. Tobacco Thrips Soybean Thrips	0.015-0.02	1.92-2.56	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5-7 days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. • Applications may also be made with equipment adapted and calibrated for ULV sprays. Warrior Insecticide with Zeon Technology may be mixed with once-refined vegetable oil and applied in a minimum of at least one quart of finished spray per acre. • Under light bollworm/budworm infestation levels, 0.02 lb. a.i./A may be applied in conjunction with intense field monitoring. • For boll weevil control spray on a 3-5 day schedule. • When applied according to label directions for control of cotton bollworm and tobacco budworm, Warrior Insecticide with Zeon Technology also provides ovicidal control of unhatched <i>Heliothis</i> spp. eggs. • Do not apply within 21 days of harvest. • Do not graze livestock in treated areas. • Do not apply more than 1.6 pts. (0.2 lb. a.i./A) per season. • Do not make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush® insecticide, Ammo® insecticide, Asana® XL insecticide, Baythroid® emulsifiable pyrethroid insecticide, Capture® insecticide/miticide, Danitol® 2.4 EC Spray insecticide/miticide, Decis® insecticide, Fury™ insecticide, Karate® insecticide, Mustang® insecticide, Pounce®, Scout X-TRA® insecticide, SynerGin™ insecticide and Warrior insecticide with Zeon Technology.
	Lygus Bug Spp. ³ Pink Bollworm Cabbage Looper Cotton Leafperforator Saltmarsh Caterpillar Cotton Leafworm Cotton Fleahopper	0.02-0.03	2.56-3.84	
	Cotton Bollworm Tobacco Budworm ³ Boll Weevil Fall Armyworm Beet Armyworm ^{1,3} European Corn Borer Brown Stink Bug Green Stink Bug Southern Green Stink Bug Twospotted Spider Mite ² Cotton Aphid ^{2,3} Bandedwing Whitefly ^{2,3} Sweetpotato Whitefly ^{2,3}	0.025-0.04	3.20-5.12	

¹For control of first and second instar only.
²Suppression only.
³See resistance statement under General Directions for Use.

13/23

Warrior® with Zeon™ Technology

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Lettuce (Head and Leaf)	Alfalfa Looper Cabbage Looper Imported Cabbageworm Cutworm Spp. Saltmarsh Caterpillar Green Cloverworm	0.015–0.025	1.92–3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A. • Do not apply within 1 day of harvest. • Do not apply more than 0.3 lb. a.i. (2.4 pts.)/A per season.
	Diamondback Moth ³ Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Southern Armyworm Corn Earworm Tobacco Budworm ³ European Corn Borer Flea Beetle Spp. Japanese Beetle (Adult) Vegetable Weevil (Adult) Grasshopper Spp. Leafhopper Spp. Plant Bug Spp. including Lygus Spp. ³ Stink Bug Spp. Meadow Spittlebug Aphid Spp. ^{2,3} Whitefly Spp. ^{2,3} Spider Mite Spp. ²	0.02–0.03	2.56–3.84	
Onion (Bulb and Garlic)	Cutworm Spp. Seedcorn Maggot (Adult) Onion Maggot (Adult) Leafminer Spp. (Adult)	0.015–0.025	1.92–3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Use the higher label rates as thrips population increases and avoid rescue situations. • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A. • For thrips, control by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v, or a silicone adjuvant (follow manufacturer's use directions) may enhance the deposition of the spray and increase plant coverage. • Do not apply within 14 days of harvest. Do not apply more than 0.24 lb. a.i. (1.92 pts.)/A per season.
	Armyworm Spp. ¹ Onion Thrips Tobacco Thrips Western Flower Thrips ^{2,3} Flower Thrips ² Aphid Spp. ² Plant Bug Spp. Stink Bug Spp.	0.02–0.03	2.56–3.84	

¹For control of the first and second instar only.
²Suppression only.
³See resistance statement under General Directions for Use.

14/23

Warrior® with Zeon™ Technology

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Peanut	Cutworm Spp.	0.015–0.025	1.92–3.20	<ul style="list-style-type: none">• Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.• Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.• Do not apply within 14 days of harvest.• Do not apply more than 0.12 lb. a.i. (0.96 pts.)/A per season.• Do not graze livestock in treated areas. Do not use treated vines or hay for animal feed.
	Green Cloverworm			
	Velvetbean Caterpillar			
	Red-necked Peanut Worm			
	Potato Leafhopper			
	Corn Earworm	0.02–0.03	2.56–3.84	
	Fall Armyworm ¹			
	Bean Leaf Beetle			
	Southern Corn Rootworm (Adult)			
	Vegetable Weevil			
	Whitefringed Beetle (Adult)			
	Stink Bug Spp.	0.03	3.84	
	Tobacco Thrips			
	Grasshopper Spp.			
	Beet Armyworm ^{1,3}			
	Soybean Looper ^{2,3}			
	Lesser Cornstalk Borer ²			
	Spider Mite Spp. ²			
	Aphid Spp. ²			
¹ For control of the first and second instar only.				
² Suppression only.				
³ See resistance statement under General Directions for Use.				

¹For control of the first and second instar only.
²Suppression only.
³See resistance statement under General Directions for Use.

15/23

Warrior® with Zeon™ Technology

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Rice	True Armyworm Fall Armyworm Yellow-striped Armyworm Rice Water Weevil (Adult) Rice Stink Bug Chinch Bug Grasshopper Spp. Leafhopper Spp. Oat Birdcherry Aphid Green Bug	0.025-0.04	3.20-5.10	<ul style="list-style-type: none"> • Apply as required by scouting. Timing and frequency of application should be based upon insect populations reaching locally determined economic thresholds. Determine the need for repeat applications, usually at intervals of 5-7 days, by scouting. • Warrior Insecticide with Zeon Technology can be safely used when propanil products are being used for weed control. • Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 5 gals. of water/A. • For control of rice water weevil in dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. Do not exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations. • For control of rice water weevil in water seeded rice, make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars usually when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations.

continued...

Warrior® with Zeon™ Technology

16/23

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Rice (continued)	True Armyworm Fall Armyworm Yellow-striped Armyworm Rice Water Weevil (Adult) Rice Stink Bug Chinch Bug Grasshopper Spp. Leafhopper Spp. Oat Birdcherry Aphid Green Bug	0.025-0.04	3.20-5.10	<ul style="list-style-type: none"> • California: In addition to above directions for control of rice water weevil in water seeded rice, Warrior Insecticide with Zeon Technology may be applied at the 1-3 leaf growth stage, with the majority at the 2 leaf growth stage. Adults are vulnerable on levees and in the water. Larvae are vulnerable while feeding on the leaf prior to entering the soil. Monitor for adults, based upon field history and density of population. Monitor field edges and levee areas for adults. Treat in the following manner: a) spray the inside perimeter of the field, or b) spray the entire field. • Green bug is known to have many biotypes. Warrior Insecticide with Zeon Technology may only provide suppression. If satisfactory control is not achieved with the first application of Warrior Insecticide with Zeon Technology, a resistant biotype may be present. Use alternate chemistry for control. • Do not release flood water within 7 days of an application. • Do not apply more than 0.12 lb. a.i. (0.96 pts.)/A per season. Do not apply more than 0.08 lb. a.i. (0.64 pts.)/A within 28 days of harvest or more than 0.04 lb. a.i. (0.32 pts.)/A within 21 days of harvest. • Do not apply within 21 days of harvest. • Do not use treated rice fields for the aquaculture of edible fish and crustacea. • Do not apply as an ultra-low volume (ULV) spray.

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Sorghum (Grain)	Cutworm Spp. Sorghum Midge	0.015-0.02	1.92-2.56	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gals. of water/A. • For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 5-day intervals if needed. • For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of sorghum plants. Repeat applications at 3- to 5-day intervals if needed. Warrior Insecticide with Zeon Technology may only suppress heavy infestations and/or subsequent migrations. • Do not apply within 30 days of harvest. Do not apply more than 0.08 lb. a.i. (0.64 pts.)/A per season. Do not apply more than 0.06 lb. a.i. (0.48 pts.)/A per season after crop emergence. Do not apply more than 0.02 lb. a.i. (0.16 pts.)/A per season once crop is in soft dough stage. • Do not graze livestock in treated areas or harvest for fodder, silage or hay.
	Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Yellow-striped Armyworm ¹ Corn Earworm Webworm Spp. European Corn Borer ² Southwestern Corn Borer ² Lesser Cornstalk Borer ² Flea Beetle Spp. Stink Bug Spp. Grasshopper Spp.	0.02-0.03	2.56-3.84	
	Chinch Bug	0.03	3.84	¹ For control of the first and second instars only. ² For control before the larva bores into the plant stalk. ³ See resistance statement under General Directions for Use.

³See resistance statement under General Directions for Use.

18/23

Warrior® with Zeon™ Technology

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A.	
Soybean	Corn Earworm Velvetbean Caterpillar Green Cloverworm Cabbage Looper Painted Lady (Thistle) Caterpillar Saltmarsh Caterpillar Woollybear Caterpillar Cutworm Spp. Bean Leaf Beetle Mexican Bean Beetle Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Three-Cornered Alfalfa Hopper Potato Leafhopper Thrips Spp. Chinese Aphid ⁴	0.015–0.025	1.92–3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Do not graze or harvest treated soybean forage, straw or hay for livestock feed. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A. • For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program, use a minimum of 2.56 fl. oz./A (0.02 lb. a.i./A). • Do not apply within 45 days of harvest. Do not apply more than 0.06 lb. a.i. (0.48 pts.)/A per season.
	Armyworm ¹ Fall Armyworm ¹ Yellow-striped Armyworm ¹ Tobacco Budworm ³ Webworm Spp. European Corn Borer Silverspotted Skipper Japanese Beetle (Adult) Blister Beetle Spp. Stink Bug Spp. Plant Bug Spp. Grasshopper Spp. Beet Armyworm ^{1,3} Soybean Looper ^{2,3} Lesser Cornstalk Borer ² Spider Mite Spp. ²	0.025–0.03	3.20–3.84	¹ For control of the first and second instar only. ² Suppression only. ³ See resistance statement under General Directions for Use. ⁴ Use lower rates for early season applications and/or lighter populations.
		0.03	3.84	

Crop	Target Pests	Rate		Remarks	
		lb. a.i./A	fl. oz./A.		
Sunflower	Sunflower Beetle	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of sunflower heads and/or foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. • Do not apply within 45 days of harvest. • Do not apply more than 0.12 lb. a.i. (0.96 pts.)/A per season. Do not apply more than 0.09 lb. a.i. (0.72 pts.)/A per season after bloom initiation. Do not apply as an ultra-low volume (ULV) spray. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.</p>	
	Cutworm Spp.				
	Sunflower Moth	0.02-0.03	2.56-3.84		
	Banded Sunflower Moth				
	Fall Armyworm ¹				
	Woollybear Caterpillar				
	Spotted Cabbage Looper				
	Painted Lady (Thistle) Caterpillar				
	Seed Weevil (Adult)				
	Stem Weevil (Adult)				
	Head-Clipper Weevil (Adult)				
	Japanese Beetle (Adult)				
Tomato and Tomatillo	Sunflower Maggot (Adult)	0.03	3.84		
	Leafhopper Spp.				
	Meadow Spittlebug	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A. • Do not apply within 5 days of harvest. • Do not apply more than 0.36 lb. a.i. (2.88 pts.)/A per season. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.</p>	
	Stink Bug Spp.				
	Grasshopper Spp.				
	Beet Armyworm ^{1,3}				
	Spider Mite Spp. ²				
	Cabbage Looper				
	Cutworm Spp.				
	Hornworm Spp.				
	Tomato Fruitworm	0.02-0.03	2.56-3.84		
	Tobacco Budworm ³				
Tomato and Tomatillo	Tomato Pinworm	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A. • Do not apply within 5 days of harvest. • Do not apply more than 0.36 lb. a.i. (2.88 pts.)/A per season. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.</p>	
	Beet Armyworm ^{1,3}				
	Southern Armyworm ¹				
	Yellow-striped Armyworm ¹				
	Fall Armyworm ¹				
	European Corn Borer				
	Leafminer Spp. ²				
	Colorado Potato Beetle ³				
	Flea Beetle Spp.				
	Grasshopper Spp.				
Tomato and Tomatillo	Leafhopper Spp.	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A. • Do not apply within 5 days of harvest. • Do not apply more than 0.36 lb. a.i. (2.88 pts.)/A per season. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.</p>	
	Aphid Spp. ^{2,3}				
	Whitefly Spp. ^{2,3}	0.015-0.025	1.92-3.20		
	Meadow Spittlebug				
	Stink Bug Spp.				
	Plant Bug Spp.				
	Cabbage Looper				
	Cutworm Spp.				
	Hornworm Spp.				
	Tomato Fruitworm				
	Tobacco Budworm ³				
	Tomato Pinworm				

¹For control of first and second instar only.
²Suppression only.
³See resistance statement under General Directions for Use.

20/23

Warrior® with Zeon™ Technology

SPRAY RECOMMENDATIONS				
Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
Wheat, Wheat Hay, and Triticale	Cutworm Spp.	0.015–0.025	1.92–3.20	<ul style="list-style-type: none"> • Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. • Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A. • For chinch bug control, repeat applications at 3-5-day intervals if needed. Warrior Insecticide with Zeon Technology may only suppress heavy infestations and/or migrations. • Greenbug is known to have many biotypes. Warrior Insecticide with Zeon Technology may provide suppression only. In this situation, a second application using an alternative chemistry may be needed. • Do not apply within 30 days of harvest. • Do not apply more than 0.06 lb. a.i. (0.48 pts.)/A per season. <p>¹Best control is obtained before insects begin to roll leaves. Once wheat has started to boot, Warrior Insecticide with Zeon Technology may provide suppression only. Higher rates and increased coverage will be necessary.</p> <p>²See resistance statement under General Directions for Use.</p>
	Army Cutworm			
	Armyworm	0.02–0.03	2.56–3.84	
	Fall Armyworm			
	Yellow-striped Armyworm			
	Flea Beetle Spp.			
	Cereal Leaf Beetle			
	Stink Bug Spp.			
	English Grain Aphid ¹			
	Russian Wheat Aphid ¹			
	Bird Cherry-Oat Aphid ¹			
	Grasshopper Spp.			
	Grass Sawfly	0.025–0.03	3.20–3.84	
	Chinch Bug	0.03	3.84	
	Greenbug ^{1,2}			

Rate Conversion Chart

lb. a.i./A	fl. oz./A	pts./A	Treated Acres/gal.
0.015	1.92	0.12	66
0.02	2.56	0.16	50
0.025	3.20	0.20	40
0.03	3.84	0.24	33
0.04	5.12	0.32	25

21/23

Warrior® with Zeon™ Technology

Ambush®, Warrior®, E-Z Handler®, Zeon™, and the Syngenta logo are trademarks of a Syngenta Group Company

Ammo®, Capture®, Fury™, Mustang®, and Pounce® are trademarks of FMC Corporation

Asana® is a trademark of E. I. du Pont de Nemours & Co. (Inc.)

Baythroid® is a trademark of Bayer AG

Danitol® is a trademark of Valent USA Corporation

DECIS® and SCOUT-XTRA® are registered trademarks of the Aventis Group

SynerGin™ is a trademark of the Aventis Group

© 2001 Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.
--

Product of the United Kingdom
Formulated in the USA

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com
SCP 1112A-L1 0801

RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR BY PERSONS UNDER THEIR DIRECT SUPERVISION SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

**Insecticide**

Active ingredient:	
Lambda-cyhalothrin: [1 α (S*),	
3 α (Z)]-(s)-cyano-(3-phenoxyphenyl)	
methyl-3-(2-chloro-3,3,3-trifluoro-	
1-propenyl)-2,2-dimethylcyclo	
propanecarboxylate	11.4%
Other Ingredients:	88.6%
Total:	100.0%

Warrior Insecticide with Zeon Technology contains 1 lb. of active ingredient per gallon and is a capsule suspension.

See directions for use in attached booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1112

EPA Est. 070989-AR-001

Warrior®, E-Z Handler® and the Syngenta logo are trademarks of a Syngenta Group Company

©2001 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1112A-L1 0801

1 gallon

U.S. Standard Measure

KEEP OUT OF REACH OF CHILDREN.

SHAKE WELL BEFORE USING.

WARNING/AVISO

Si usted no entiende le etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If swallowed: Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

If on skin: Get medical attention if irritation persists. Wash with plenty of soap and water.

If in eyes: Hold eyelids open and flush with steady, gentle stream of water. Remove any contact lenses and repeat flushing procedure for several minutes. Get medical attention if irritation persists.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER: For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372.

syngenta

**KEEP OUT OF REACH OF
CHILDREN.
SHAKE WELL BEFORE USING.
WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING

May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. May cause allergic skin reactions. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2-30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Container Disposal

Triple rinse (or equivalent): then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

©2001 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1112A-L8 0801