

**RESTRICTED USE PESTICIDE
Due to Toxicity to Fish and Aquatic Organisms**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**WARRIOR® Insecticide with Zeon™ Technology
Insecticide**

Active Ingredient:

Lambda-cyhalothrin

[1α(S*),3α(Z)]-(±)-cyano-(3-phenoxyphenyl)methyl-
3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-

dimethylcyclopropanecarboxylate.....11.4%

Other Ingredients: 88.6%

Total: 100.0%

Warrior Insecticide with Zeon Technology contains 1 lb. of active ingredient per gal. and is a capsule suspension.

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)


See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1112
EPA Est. XXXX

Product of the United Kingdom
Formulated in USA

SCP 1112-M (DRAFT 2.1)

U.S. Standard Measure



2827

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. May cause allergic skin reactions. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hours after exposure and may last 2-30 hours, without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, or Viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

3 7 27

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

4 3 27

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, or Viton \geq 14 mils
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

STORAGE AND DISPOSAL

Prohibitions

Do not contaminate water, food, or feed by storage and disposal.

Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area. **DO NOT ALLOW PRODUCT TO FREEZE.**

6727

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

For Bulk, Mini-Bulk, and Ez Handler® Disposal

Return container to Syngenta for reuse with seal intact and in salable condition.

Container Precautions

Before refilling RETURNABLE CONTAINERS, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices.

REFILL ONLY WITH WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY. The contents of RETURNABLE CONTAINERS cannot be completely removed by cleaning. Refilling with materials other than Warrior Insecticide with Zeon Technology will result in contamination and may weaken container.

After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals./A by air or 10 gals./A by ground unless otherwise specified in this label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, Warrior Insecticide with Zeon Technology may be applied before, during, or after planting. For soil incorporated applications, use higher rates for improved control.

Resistance

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Spray Drift Precautions

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

- Do not apply by ground within 25 ft., or by air within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 ft. when ultra-low volume (ULV) application is made.
- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.
- Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 ft. above the crop canopy should be avoided.
- Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3–10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.
- Risk of exposure to aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
- Do not cultivate within 10 ft. of the aquatic area so as to allow growth of a vegetative filter strip.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.
- Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

In the State of New York, a 25 ft. vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for both aerial or ground application. For aerial applications, the 25 ft. vegetated non-cropped buffer strip for runoff protection would be part of the larger 150 ft. buffer strip (or 450 ft. buffer strip for ULV application) required for spray drift.

Tank Mix Application

When tank mixing with any other agricultural products, always add Warrior Insecticide with Zeon Technology last. Fill the tank with one half to two thirds volume of the mixing diluent. Make sure all other products are fully dispersed in the mixing diluent before adding the recommended rate of Warrior Insecticide with Zeon Technology to the tank. Add the remainder of the mixing diluent volume. It is recommended that mixing and spray equipment have continuous agitation for best results. Follow the precautions and limitations of the most restricted product in the tank mixture.

While Warrior Insecticide with Zeon Technology has good flexibility for tank mixing with other agricultural products, a jar test for physical compatibility is recommended for untried mixtures using proper ratios and mixing sequences of all ingredients to be included in the mixture.

SPECIFIC USE DIRECTIONS

AGRICULTURAL USES

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
ALFALFA AND ALFALFA GROWN FOR SEED				
	Alfalfa Caterpillar Cutworm Spp. Green Cloverworm Looper Spp. Velvetbean Caterpillar Webworm Spp. Leafhopper Spp. Threecornered Alfalfa Hopper	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply only to fields planted to pure strands of alfalfa. Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals. per acre by air or 10 gals./A by ground. When foliage is dense and/or pest populations are high 5-10 gals./A by air or 20 gals./A by ground and higher use rates are recommended. Use higher rates for increased residual control. Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2-3 days following application. Avoid direct application to bee shelters. Do not apply more than 0.03 lb. a.i. (0.24 pts.) per acre per cutting. Do not apply more than 0.12 lb. a.i. (0.96 pts.) per acre per season. Do not apply within 1 day of harvest for forage or within 7 days of harvest for hay. <p>¹ For control of first and second instar only. ² Suppression only. ³ See resistance statement under General Directions for Use.</p>
	Amyworm Corn Earworm Fall Armyworm ¹ Western Yellow-striped Armyworm Yellow-striped Armyworm Alfalfa Weevil Bean Leaf Beetle (Adult) Blister Beetle Spp. Clover Leaf Weevil Spp. Clover Root Borer (Adult) Clover Root Curculio Spp. (Adult) Clover Stem Borer (Adult) Cowpea Curculio (Adult) Cowpea Weevil (Adult) Cucumber Beetle Spp. (Adult) Egyptian Alfalfa Weevil Grape Colaspis (Adult) Green June Beetle (Adult) Japanese Beetle (Adult) Mexican Bean Beetle Pea Weevil (Adult) Sweet Clover Weevil (Adult) Whitefringed Beetle Spp. (Adult) Meadow Spittlebug Plant Bug Spp. including Lygus Spp. ³ Stink Bug Spp. Alfalfa Seed Chalcid (Adult) Blue Alfalfa Aphid Cowpea Aphid Green Peach Aphid ³ Pea Aphid Spotted Alfalfa Aphid Thrips Spp. Grasshopper Spp.	0.02-0.03	2.56-3.84	
	Beet Armyworm ^{1,2} Blotch Leafminer ² Spider Mites ²	0.03	3.84	

It is not recommended that Warrior Insecticide with Zeon Technology be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Use Precautions: Sprinkler Irrigation Application

- A. Apply this product only through (sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Do not apply through chemigation systems connected to public water systems.

Warrior Insecticide with Zeon Technology is an aqueous based formulation. It is recommended that no type of non-emulsifiable oils be used in combination with Warrior Insecticide with Zeon Technology. If adjuvants are used, use only:

- Nonionic Surfactant (NIS) containing at least 75% surface agent, or
- Non-phytotoxic Crop Oil Concentrate (COC), including once refined Vegetable Oil Concentrate (VOC),
- Methylated Sunflower Oils (MSO) containing a minimum of 17% emulsifier.

Adjuvants other than NIS or COC may be used providing the product meets the following criteria:

1. Contains only EPA exempt ingredients.
2. Is non-phytotoxic to the target crop.
3. Is compatible in mixture. (May be established through a jar test.)
4. Is supported locally for use with Warrior Insecticide with Zeon Technology on the target crop through proven field trials and through university and extension recommendations.

In addition, the following may be used as diluents:

- Crop Oil Concentrate
- Methylated Sunflower Oils
- Urea-Ammonium Nitrate

It is recommended that the following not be used in combination with Warrior Insecticide with Zeon Technology as diluents or adjuvants:

- Non-emulsifiable oils
- Diesel Fuel
- Straight Mineral Oil

CHEMIGATION

Sprinkler Irrigation Application

Apply Warrior Insecticide with Zeon Technology at rates and timing described elsewhere in this label. As local recommendations differ, consult your local State Extension Service or other local experts for recommendations on adjuvant or diluent types, (see **Tank Mix Application**) rates and mixing instructions. These recommendations should be proven, through university and extension field trials, to be effective with Warrior Insecticide with Zeon Technology applied by chemigation.

Check the irrigation system to insure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period.

Apply by injecting the recommended rate of Warrior Insecticide with Zeon Technology into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1–0.2 acre-inch of water. In general, use the least amount of water required for proper distribution and coverage. It is recommended that the product be injected into the main irrigation line ahead of a right angle turn in the line to insure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the recommended rate of Warrior Insecticide with Zeon Technology for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

Crop	Target Pests	Rate		Remarks																												
		lb. a.i./A	fl. oz./A																													
CANOLA																																
	Cutworm Spp. Armyworm Spp. Diamondback Moth Flea Beetle Cabbage Seedpod Weevil Lygus Bug Grasshoppers	0.015-0.03	1.92-3.84	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply a minimum of 2 gallons of water per acre. Do not apply within 7 days of harvest. 																												
	Cabbage Aphid	0.03	3.84	<ul style="list-style-type: none"> Do not apply more than 0.09 lb. ai (0.72 pints) per acre per year. 																												
CEREAL GRAINS																																
Corn (At Plant): Field Corn Popcorn Seed Corn Sweet Corn	Corn Rootworm Larvae Western Northern Southern Mexican Cutworm Spp. Seedcorn Maggot Seedcorn Beetle Lesser Cornstalk Borer White Grub Spp. Wireworm Spp. Billbug ¹ Red Imported Fire Ant ¹	0.05	0.66	<ul style="list-style-type: none"> Banded Applications – Apply at planting as a 5–7 inch T-band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel. In-Furrow Applications – Apply into the seed furrow through spray nozzles or microtubes, behind the planter furrow openers and in front of the press wheel. Apply a minimum of 3 gallons finished spray per acre. Do not harvest or graze livestock or cut treated crops for feed within 21 days of at plant application. Do not apply more than 0.09 lb. ai (0.72 pints) per acre per crop at plant. For field corn, popcorn, and seed corn do not apply more than 0.12 lb. ai per acre per crop from at plant and foliar applications. For sweet corn do not apply more than 0.48 lb. ai per acre per crop from at plant and foliar applications. <p>¹Suppression only.</p> <p>Lbs. ai and fl. oz./A of Warrior Insecticide With Zeon Technology Applied at 0.66 fl. oz./1000 ft. of Row for Various Row Spacings</p> <table border="1"> <thead> <tr> <th>Row Spacing</th> <th>40"</th> <th>38"</th> <th>36"</th> <th>34"</th> <th>32"</th> <th>30"</th> </tr> </thead> <tbody> <tr> <td>Linear Ft./A</td> <td>13,068</td> <td>13,756</td> <td>14,520</td> <td>15,374</td> <td>16,335</td> <td>17,424</td> </tr> <tr> <td>Lbs. ai/A</td> <td>0.067</td> <td>0.07</td> <td>0.075</td> <td>0.079</td> <td>0.084</td> <td>0.09</td> </tr> <tr> <td>Fl. oz./A</td> <td>8.6</td> <td>9.1</td> <td>9.6</td> <td>10.1</td> <td>10.8</td> <td>11.5</td> </tr> </tbody> </table>	Row Spacing	40"	38"	36"	34"	32"	30"	Linear Ft./A	13,068	13,756	14,520	15,374	16,335	17,424	Lbs. ai/A	0.067	0.07	0.075	0.079	0.084	0.09	Fl. oz./A	8.6	9.1	9.6	10.1	10.8	11.5
Row Spacing	40"	38"	36"	34"	32"	30"																										
Linear Ft./A	13,068	13,756	14,520	15,374	16,335	17,424																										
Lbs. ai/A	0.067	0.07	0.075	0.079	0.084	0.09																										
Fl. oz./A	8.6	9.1	9.6	10.1	10.8	11.5																										

12 8 27

WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY

WARZEON1112MAS-0902

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CEREAL GRAINS				
Corn (Foliar) Field Corn Popcorn Seed Corn	Cutworm Spp. Western Bean Cutworm ¹ Corn Earworm ¹ Green Cloverworm Meadow Spittlebug	0.015–0.025	1.92–3.20	<ul style="list-style-type: none"> Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods. Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gals. of water per acre. For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3-5-day intervals if needed. Warrior Insecticide with Zeon Technology may only suppress heavy infestations and/or subsequent migrations. For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program use a minimum of 3.84 fl. oz./A (0.03 lb. a.i./A). Do not apply within 21 days of harvest. Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment. Do not apply more than 0.12 lb. a.i. (0.96 pts.) per acre per season. Do not apply more than 0.06 lb. a.i. (0.48 pts.) after silk initiation. Do not apply more than 0.03 lb. a.i. (0.24 pts.) after corn has reached the milk stage (yellow kernels with milky fluid). <p>¹For control before the larva bores into the plant stalk or ear. ²For control of first and second instar only. ³Suppression only. ⁴See resistance statement under General Directions for Use.</p>
	Tobacco Budworm ^{1,4} European Corn Borer ¹ Southwestern Corn Borer ¹ Stalk Borer ¹ Hop Vine Borer ¹ Armyworm ² Fall Armyworm ² Yellow-striped Armyworm ² Webworm Spp. Flea Beetle Spp. Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Bean Leaf Beetle Cereal Leaf Beetle Japanese Beetle (Adult) Sap Beetle (Adult) Stink Bug Spp. Grasshopper Spp. Corn Leaf Aphid ³ Bird Cherry-Oat Aphid ³ English Grain Aphid ³	0.02–0.03	2.56–3.84	
	Beet Armyworm ^{2,4} Chinch Bug ^{3,4} Green Bug ^{3,4}	0.03	3.84	

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CEREAL GRAINS				
Sweet Corn (Foliar)	Corn Earworm Fall Armyworm ¹ Southern Armyworm ¹ Beet Armyworm ^{1,2} Yellow-Striped Armyworm ¹ Cutworm Spp. Western Bean Cutworm Webworm Spp. European Corn Borer Southwestern Corn Borer Common Cornstalk Borer Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Japanese Beetle (Adult) Sap Beetle (Adult) Flea Beetle Spp. Tarnished Plant Bug Stink Bug Spp. Chinch Bug Aster Leafhopper Grasshopper Spp. Aphid Spp. ^{2,3} Spider Mite Spp. ²	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 4 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods and should be targeted for control before insects enter the stalk or ear. Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage and ears (if present). When applying by air, apply in a minimum of 2 gals. of water per acre. For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program use a minimum of 3.2 fl. oz./A (0.025 lb. a.i./A). Do not apply within 1 day of harvest. Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment. Do not apply more than 0.48 lb. a.i. (3.84 pts.) per acre per season. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.</p>
	Corn Silkfly (Adult) ²	0.03	3.84	

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CEREAL GRAINS				
Rice	True Armyworm Fall Armyworm Yellow-striped Armyworm Rice Water Weevil (Adult) Rice Stink Bug Chinch Bug Grasshopper Spp. Leafhopper Spp. Oat Birdcherry Aphid Green Bug	0.025-0.04	3.20-5.10	<ul style="list-style-type: none"> Apply as required by scouting. Timing and frequency of application should be based upon insect populations reaching locally determined economic thresholds. Determine the need for repeat applications, usually at intervals of 5-7 days, by scouting. Warrior Insecticide with Zeon Technology can be safely used when propanil products are being used for weed control. Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals of water (or a total carrier volume) per acre but ensure sufficient volume is used to provide adequate coverage. In addition, adding an emulsifiable crop oil (e.g., 1 pint per acre) when lower aerial application volumes are used as recommended to help improve coverage, reduce evaporation, and improve efficacy. For control of rice water weevil in dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. Do not exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations. For control of rice water weevil in water seeded rice, make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars usually when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations. California: In addition to above directions for control of rice water weevil in water seeded rice, Warrior Insecticide with Zeon Technology may be applied at the 1-3 leaf growth stage, with the majority at the 2 leaf growth stage. Adults are vulnerable on levees and in the water. Larvae are vulnerable while feeding on the leaf prior to entering the soil. Monitor for adults, based upon field history and density of population. Monitor field edges and levee areas for adults. Treat in the following manner: a) spray the inside perimeter of the field, or b) spray the entire field. Green bug is known to have many biotypes. Warrior Insecticide with Zeon Technology may only provide suppression. If satisfactory control is not achieved with the first application of Warrior Insecticide with Zeon Technology, a resistant biotype may be present. Use alternate chemistry for control. Do not release flood water within 7 days of an application. Do not apply more than 0.12 lb. a.i. (0.96 pts.) per acre per season. Do not apply more than 0.08 lb. a.i. (0.64 pts.) per acre within 28 days of harvest or more than 0.04 lb. a.i. (0.32 pts.) per acre within 21 days of harvest. Do not apply within 21 days of harvest. Do not use treated rice fields for the aquaculture of edible fish and crustacea. Do not apply as an ultra-low volume (ULV) spray.

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CEREAL GRAINS				
Sorghum (Grain)	Cutworm Spp. Sorghum Midge	0.015-0.02	1.92-2.56	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
	Armyworm Beet Armyworm ^{1,2} Fall Armyworm ¹ Yellow-striped Armyworm ¹ Corn Earworm Webworm Spp. European Corn Borer ² Southwestern Corn Borer ² Lesser Cornstalk Borer ² Flea Beetle Spp. Stink Bug Spp. Grasshopper Spp.	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gals. of water per acre. For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 5-day intervals if needed. For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of sorghum plants. Repeat applications at 3-5-day intervals if needed. Warrior Insecticide with Zeon Technology may only suppress heavy infestations and/or subsequent migrations. Do not apply within 30 days of harvest. Do not apply more than 0.08 lb. a.i. (0.64 pts.) per acre per season. Do not apply more than 0.06 lb. a.i. (0.48 pts.) per acre per season after crop emergence. Do not apply more than 0.02 lb. a.i. (0.16 pts.) per acre per season once crop is in soft dough stage.
	Chinch Bug	0.03	3.84	¹ For control of the first and second instars only. ² For control before the larva bores into the plant stalk. ³ See resistance statement under General Directions for Use.

16 3 27

WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY
 WARZEON1112MAS-0902
 Page 16

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CEREAL GRAINS				
Wheat, Wheat Hay, and Triticale	Cutworm Spp. Army Cutworm	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. For chinch bug control, repeat applications at 3- to 5-day intervals if needed. Warrior Insecticide with Zeon Technology may only suppress heavy infestations and/or migrations. Greenbug is known to have many biotypes. Warrior Insecticide with Zeon Technology may provide suppression only. In this situation, a second application using an alternative chemistry may be needed. Do not apply within 30 days of harvest. Do not apply more than 0.06 lb. a.i. (0.48 pts.) per acre per season. <p>¹Best control is obtained before insects begin to roll leaves. Once wheat has started to boot, Warrior Insecticide with Zeon Technology may provide suppression only. Higher rates and increased coverage will be necessary. ²See resistance statement under General Directions for Use.</p>
	Amyworm Fall Armyworm Yellow-striped Armyworm Flea Beetle Spp. Cereal Leaf Beetle Stink Bug Spp. English Grain Aphid ¹ Russian Wheat Aphid ¹ Bird Cherry-Oat Aphid ¹ Grasshopper Spp. Orange Blossom Wheat Midge	0.02-0.03	2.56-3.84	
	Grass Sawfly	0.025-0.03	3.20-3.84	
	Chinch Bug Greenbug ² Corn Leaf Aphid ² Mite Spp. ²	0.03	3.84	

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
COLE CROPS				
Broccoli Brussels Sprouts Cabbage Cavalo Broccoli Cauliflower Chinese Broccoli (gai lon) Chinese Cabbage (napa), Chinese Mustard Cabbage (gai choy) Kohlrabi	Alfalfa Looper Cabbage Looper Imported Cabbageworm Southern Cabbageworm Cutworm Spp. Cabbage Webworm	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. Do not apply within 1 day of harvest. Do not apply more than 0.24 lb. a.i. (1.92 pts.) per acre per season. ¹ For control of first and second instar only. ² Suppression only. ³ See resistance statement under General Directions for Use.
	Diamondback Moth ³ Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Yellow-striped Armyworm Corn Earworm Flea Beetle Spp. Japanese Beetle (Adult) Vegetable Weevil (Adult) Grasshopper Spp. Leafhopper Spp. Plant Bug Spp. including Lygus Spp. ³ Stink Bug Spp. Meadow Spittlebug Aphid Spp. ^{2,3} Whitefly Spp. ^{2,3} Thrips Spp. ² Spider Mite Spp. ²	0.02-0.03	2.56-3.84	

WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY
 WARZEON1112MAS-0902
 Page 18

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
COTTON				
	Cutworm Spp. Tobacco Thrips Soybean Thrips	0.015-0.02	1.92-2.56	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 to 7 days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Applications may also be made with equipment adapted and calibrated for ULV sprays. Warrior Insecticide with Zeon Technology may be mixed with once-refined vegetable oil and applied in a minimum of at least one quart of finished spray per acre. Under light bollworm/budworm infestation levels, 0.02 lb. a.i./A may be applied in conjunction with intense field monitoring. For boll weevil control spray on a 3 to 5 day schedule. When applied according to label directions for control of cotton bollworm and tobacco budworm, Warrior Insecticide with Zeon Technology also provides ovicidal control of unhatched <i>Heliothis</i> spp. eggs. Do not apply within 21 days of harvest. Do not graze livestock in treated areas. Do not apply more than 1.6 pts. (0.2 lb. a.i.) per acre per season. Do not make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ammo® insecticide, Asana® XL insecticide, Baythroid® emulsifiable pyrethroid insecticide, Capture® Insecticide/miticide, Danitol® 2.4 EC Spray insecticide/miticide, Decis® Insecticide, Fury™ Insecticide, Karate® insecticide, Karate® insecticide with Zeon™ technology, Mustang® insecticide, Scout X-TRA® insecticide, SynerGin™ Insecticide and Warrior insecticide with Zeon™ technology. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.</p>
	Lygus Bug Spp. ³ Pink Bollworm Cabbage Looper Cotton Leafperforator Saltmarsh Caterpillar Cotton Leafworm Cotton Fleahopper	0.02-0.03	2.56-3.84	
	Cotton Bollworm Tobacco Budworm ³ Boll Weevil Fall Armyworm Beet Armyworm ^{1,3} European Corn Borer Brown Stink Bug Green Stink Bug Southern Green Stink Bug Twospotted Spider Mite ² Cotton Aphid ^{2,3} Bandedwing Whitefly ^{2,3} Sweetpotato Whitefly ^{2,3}	0.025-0.04	3.20-5.12	

19 3 27

WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY
 WARZEON1112MAS-0902
 Page 19

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
FRUITING VEGETABLES (EXCEPT CUCURBITS)				
Tomato and Tomatillo Peppers (bell and nonbell), Eggplant, Ground cherry, Pepino	Cabbage Looper Cutworm Spp. Homworm Spp.	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. Do not apply within 5 days of harvest. Do not apply more than 0.36 lb. a.i. (2.88 pts.) per acre per season. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use. ⁴For control before the larva bores into the plant stalk or fruit.</p>
	Tomato Fruitworm Tobacco Budworm ³ Tomato Pinworm Beet Armyworm ^{1,3} Southern Armyworm ¹ Yellow-striped Armyworm ¹ Fall Armyworm ¹ European Com Borer ⁴ Leafminer Spp. ² Colorado Potato Beetle ³ Flea Beetle Spp. Grasshopper Spp. Leafhopper Spp. Aphid Spp. ^{2,3} Whitefly Spp. ^{2,3} Meadow Spittlebug Stink Bug Spp. Plant Bug Spp. Stalk Borer ⁴ Blister Beetle Spp. Japanese Beetle (Adult) Pepper Weevil (Adult) ² Vegetable Weevil (Adult) Tomato Psyllid ^{2,3} Spider Mite Spp. ² Thrips ³ Cucumber Beetle Spp. (Adult)	0.02-0.03	2.56-3.84	

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
LEGUME VEGETABLES				
Edible Podded (Only) <i>Canavalia gladiata</i> – sword bean <i>Canavalia ensiformis</i> – jackbean Glycine max – Soybean (immature seed)	Cutworm Spp. Green Cloverworm Imported Cabbageworm Saltmarsh Caterpillar Velvetleaf Caterpillar Mexican Bean Beetle	0.015–0.025	1.92–3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gallons of water per acre. For edible podded and succulent shelled legume vegetables, do not apply within 7 days of harvest. For dried shelled legume vegetables, do not apply within 21 days of harvest. Do not apply more than 0.12 lb. ai (0.96 pints) per acre per season. For succulent and dried shelled peas and beans, do not graze livestock in treated areas or harvest vines for forage or hay. <p>¹For control before the larva bores into the plant stalk or pods. ²For control of the first and second instar only. ³For suppression only. ⁴See resistance statement under General Directions for Use.</p>
Edible Podded, Succulent Shelled or Dried Shelled <i>Phaseolus</i> spp. – includes: field, kidney, lima, navy, pinto, runner, snap, tepary and wax beans <i>Vigna</i> spp. – includes: adzuki, asparagus, moth, mung, rice, urd and yardlong beans, black-eye pea, catjang, Chinese longbean, cowpea, Crowder pea, and Southern pea <i>Pisum</i> spp. – includes: dwarf, edible-pod, English, field, garden, green, snow and sugar snap peas <i>Cajanus cajan</i> – Pigeon pea	Corn Earworm Painted Lady Butterfly (Larva) European Corn Borer ¹ Looper Spp. Western Bean Cutworm Tobacco Budworm ⁴ Armyworm ² Fall Armyworm ² Yellow-Striped Armyworm ² Western Yellow-Striped Armyworm ² Bean Leafskeltonizer Webworm Spp. Leafminer Spp. Alfalfa Caterpillar Stalk Borer ¹ Cucumber Beetle Spp. (Adult) Corn Rootworm Beetle Spp. (Adult) Flea Beetle Spp. (Adult) Curculio and Weevil Spp. ¹ (foliage and pod feeding adults and larvae) Blister Beetle Spp. Bean Leaf Beetle Japanese Beetle (Adult) Leafhopper Spp. Flea Hopper Spp. Three-Cornered Alfalfa Hopper Meadow Spittlebug Stink Bug Spp. Plant Bug Spp. Including Lygus Spp. ⁴ Grasshopper Spp. Thrips Spp. ⁴ Aphid Spp. ⁴	0.02–0.03	2.56–3.84	
Succulent Shelled or Dried Shelled <i>Vicia faba</i> . – broadbean (favabean)	Beet Armyworm ^{2,3,4} Soybean Looper ^{3,4} Lesser Cornstalk Borer ³ Leafminer Spp. ^{3,4} Whitefly Spp. ^{3,4} Spider Mite Spp. ³	0.03	3.84	
Dried Shelled (Only) <i>Lupinus</i> spp. – includes: grain, sweet, white and sweet white lupines <i>Cicer arietinum</i> – chickpea (garbanzo bean) <i>Cyamopsis tetragonoloba</i> – guar <i>Labiab pupureus</i> – Lablab bean (hyacinth bean) <i>Lens esculata</i> – Lentils				

WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY

WARZEON1112MAS-0902

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
LEGUME VEGETABLES (PEAS AND BEANS)				
Soybean	Corn Earworm Velvetbean Caterpillar Green Cloverworm Cabbage Looper Painted Lady (Thistle) Caterpillar Saltmarsh Caterpillar Woollybear Caterpillar Cutworm Spp. Bean Leaf Beetle Mexican Bean Beetle Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Three-Cornered Alfalfa Hopper Potato Leafhopper Thrips Spp. Chinese Aphid ⁴	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Do not graze or harvest treated soybean forage, straw or hay for livestock feed. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. For control of adult corn rootworm beetles (<i>Diabrotica</i> species) as part of an aerial applied corn rootworm control program use a minimum of 2.56 fl. oz./A (0.02 lb. a.i./A). Do not apply within 45 days of harvest. Do not apply more than 0.06 lb. a.i. (0.48 pts.) per acre per season. <p>¹For control of the first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use. ⁴Use lower rates for early season applications and/or lighter populations.</p>
	Armyworm ¹ Fall Armyworm ¹ Yellow-striped Armyworm ¹ Tobacco Budworm ³ Webworm Spp. European Corn Borer Silverspotted Skipper Japanese Beetle (Adult) Blister Beetle Spp. Stink Bug Spp. Plant Bug Spp. Grasshopper Spp.	0.025-0.03	3.20-3.84	
	Beet Armyworm ^{1,3} Soybean Looper ^{2,3} Lesser Cornstalk Borer ² Spider Mite Spp. ²	0.03	3.84	

WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY

WARZEON1112MAS-0902

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
LETTUCE (HEAD AND LEAF)				
	Alfalfa Looper Cabbage Looper Imported Cabbageworm Cutworm Spp. Saltmarsh Caterpillar Green Cloverworm	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. Do not apply within 1 day of harvest.
	Diamondback Moth ³ Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Southern Armyworm Corn Earworm Tobacco Budworm ³ European Corn Borer Flea Beetle Spp. Japanese Beetle (Adult) Vegetable Weevil (Adult) Grasshopper Spp. Leafhopper Spp. Plant Bug Spp. including Lygus Spp. ³ Stink Bug Spp. Meadow Spittlebug Aphid Spp. ^{2,3} Whitefly Spp. ^{2,3} Spider Mite Spp. ²	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> Do not apply more than 0.3 lb. a.i. (2.4 pts.) per acre per season. ¹For control of first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.
ONION (BULB) AND GARLIC				
	Cutworm Spp. Seedcorn Maggot (Adult) Onion Maggot (Adult) Leafminer Spp. (Adult)	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
	Armyworm Spp. ¹ Onion Thrips Tobacco Thrips Western Flower Thrips ^{2,3} Flower Thrips ² Aphid Spp. ² Plant Bug Spp. Stink Bug Spp.	0.02-0.03	2.56-3.84	<ul style="list-style-type: none"> Use the higher label rates as thrips population increases and avoid rescue situations. Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. For thrips control by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v or a silicone adjuvant (follow manufacturers use directions) may enhance the deposition of the spray and increase plant coverage. Do not apply within 14 days of harvest. Do not apply more than 0.24 lb. a.i. (1.92 pts.) per acre per season. <p>¹For control of the first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.</p>

WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY
 WARZEON1112MAS-0902
 Page 23

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
PEANUT				
	Cutworm Spp. Green Cloverworm Velvetbean Caterpillar Red-necked Peanut Worm Potato Leafhopper	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. Do not apply within 14 days of harvest. Do not apply more than 0.12 lb. a.i. (0.96 pts.) per acre per season. <p>¹For control of the first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.</p>
	Corn Earworm Fall Armyworm ¹ Bean Leaf Beetle Southern Corn Rootworm (Adult) Vegetable Weevil Whitefringed Beetle (Adult) Stink Bug Spp. Tobacco Thrips Grasshopper Spp.	0.02-0.03	2.56-3.84	
	Beet Armyworm ^{1,3} Soybean Looper ^{2,3} Lesser Comstalk Borer ² Spider Mite Spp. ² Aphid Spp. ²	0.03	3.84	
POME FRUITS				
Apple Crabapple Loquat Mayhaw Oriental Pear Pear Quince	Learoller Spp. Codling Moth Tufted Apple Budworm Oriental Fruit Moth Lesser Appleworm Green Fruitworm Tent Caterpillar Spp. Tentiform Leaf Miner Spp. Apple Maggot (Adult) Cherry Fruit Fly Spp. (Adult) Pear Sawfly Plum Curculio Japanese Beetle Plant Bug Spp. Stink Bug Spp. Leafhopper Spp. Periodical Cicada Apple Aphid Rosy Apple Aphid Pear Psylla San Jose Scale (fruit infestations only)	0.02-0.04	2.56-5.12	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply in a minimum of 5 gallons of water per acre. Do not apply within 21 days of harvest. Do not apply more than 0.2 lb. ai (1.6 pints) per acre per year. Do not apply more than 0.16 lb. ai (1.28 pints) per acre per year post bloom.

WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY

WARZEON1112MAS-0902

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
STONE FRUITS				
Apricot Sweet and Tart Cherry Nectarine Peach Plum Chickasaw Plum Damson Plum Japanese Plum Plumcot Prune	Leafroller Spp. Peach Twig Borer Oriental Fruit Moth Peachtree Borer Spp. Green Fruitworm Tent Caterpillar Spp. American Plum Borer Cherry Fruit Fly Spp. (Adult) Plum Curculio Rose Chafer Japanese Beetle Plant Bug Spp. Stink Bug Spp. Leafhopper Spp. Periodical Cicada Black Cherry Aphid	0.02-0.04	2.56-5.12	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 5 gallons of water per acre. Do not apply within 14 days of harvest. Do not apply more than 0.2 lb. ai (1.6 pints) per acre per year. Do not apply more than 0.16 lb. ai (1.28 pints) per acre per year post bloom.
SUGARCANE				
	Sugarcane Borer ¹ Rice Borer ¹ Sugarcane Beetle (Adult) ² Yellow Sugarcane Aphid ³	0.025-0.04	3.20-5.12	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 2 gallons of water per acre. Do not apply within 21 days of harvest. Do not apply more than 0.16 lb. ai (1.28 pints) per acre per season. <p>¹For control before the larva bores into the plant stalk. ²Suppression only of beetles active above ground. ³See resistance statement under General Directions for Use.</p>
SUNFLOWER				
	Sunflower Beetle Cutworm Spp.	0.015-0.025	1.92-3.20	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground or air equipment using sufficient water to obtain full coverage of sunflower heads and/or foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. Do not apply within 45 days of harvest. Do not apply more than 0.12 lb. a.i. (0.96 pts.) per acre per season. Do not apply more than 0.09 lb. a.i. (0.72 pts.) per acre per season after bloom initiation. Do not apply as an ultra-low volume (ULV) spray. <p>¹For control of first and second instar only. ²Suppression only. ³See resistance statement under General Directions for Use.</p>
	Sunflower Moth Banded Sunflower Moth Fall Armyworm ¹ Woollybear Caterpillar Spotted Cabbage Looper Painted Lady (Thistle) Caterpillar Seed Weevil (Adult) Stem Weevil (Adult) Head-Clipper Weevil (Adult) Japanese Beetle (Adult) Sunflower Maggot (Adult) Leafhopper Spp. Meadow Spittlebug Stink Bug Spp. Grasshopper Spp.	0.02-0.03	2.56-3.84	
	Beet Armyworm ^{1,3} Spider Mite Spp. ²	0.03	3.84	

WARRIOR INSECTICIDE WITH ZEON TECHNOLOGY

WARZEON1112MAS-0902

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
TOBACCO (AIR DRIED) BURLEY TOBACCO AND FLUE-CURED TOBACCO				
	Tobacco Budworm Tobacco Homworm Cabbage Looper Corn Earworm Salt Marsh Caterpillar Armyworm Spp. ¹ Cutworm Spp. Webworm Spp. Tobacco Flea Beetle (Adult) Cucumber Beetle Spp. (Adult) Blister Beetle Spp. Vegetable Weevil (Adult) Japanese Beetle (Adult) Grasshopper Spp. Tree Cricket Spp. Katydid Spp. Plant Bug Spp. ³ Stinkbug Spp. ² Thrips Spp. ² Aphid Spp. ^{2,3}	0.015-0.03	1.92-3.84	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage. When applying by air, apply in a minimum of 2 gals. of water per acre. Do not apply within 40 days of harvest. Do not apply more than 0.09 lb. a.i. (0.72 pts.) per acre per year. <p>¹For control of first and second instars only. ²Suppression only. ³See resistance statement under "GENERAL DIRECTIONS FOR USE."</p>
TREE NUTS				
Almond Beech Nut Brazil Nut Butternut Cashew Chestnut Chinquapin Filbert (Hazlenut) Hickory Nut Macadamia Nut (Bush Nut) Walnut, Black Walnut, English (Persian)	Leafroller Spp. Navel Orangeworm Codling Moth Filbertworm Peach Twig Borer Walnut Husk Fly Spp. (Adult) Ants Plant Bug Spp. Stink Bug Spp. Chinch Bug Leaffooted Bug Walnut Aphid	0.02-0.04	2.56-5.12	<ul style="list-style-type: none"> Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold. Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply in a minimum of 5 gals. of water per acre. Do not apply within 14 days of harvest. Do not apply more than 0.16 lb. ai (1.28 pts.) per acre per year. Do not apply more than 0.12 lb. ai (0.96 pts) per acre per year post bloom.
Pecan	Hickory Shuckworm Pecan Casebearer Spp. Pecan Weevil Pecan Aphid Spp. Pecan Spittlebug Pecan Phylloxera Spp.	0.02-0.04	2.56-5.12	

NON-AGRICULTURAL USES

Crop	Target Pests	Rate		Remarks
		lb. a.i./A	fl. oz./A	
CONIFER AND DECIDUOUS TREES				
Plantations Nurseries Seed Orchards	Pine Tip Moth Spp. Spruce Budworm Bagworm Tent Caterpillar Spp. Leafroller Spp. Gypsy Moth Webworm Spp. Tussock Moth Spp. Pine Sawfly Spp. Sawfly Spp. Pine Chafer Japanese Beetle May Beetle Spp. June Beetle Spp. Pine Colaspis Beetle Leaf Beetle Spp. Pales Weevil Pine Weevil Spp. Pine Conelet Bug Spittlebug Spp. Pine Leaf Chermid Balsam Woolly Aphid Balsam Twig Aphid	0.02-0.04	2.56-5.12	<ul style="list-style-type: none"> To control exposed foliage, flower, cone, seed and bark feeding insects, apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Apply with ground equipment using sufficient water to obtain full coverage of target site. When applying by air, apply a minimum of 2 gals. of water per acre. Do not apply more than 0.24 lb. ai (1.92 pts.) per acre per year.
	Coneworm Spp. Seed Bug Spp.	See Remarks	See Remarks	<ul style="list-style-type: none"> For high volume sprayers, dilute 5.12 fl. oz. per 100 gallons of water and apply 5 to 10 gallons of finished spray per tree. For low volume sprayers, dilute 20 fl. oz. per 100 gallons of water and apply 100 gallons of finished spray per acre. For aerial applications, apply 15 fl. oz. per acre in a minimum of 10 gallons finish spray per acre. Do not apply more than 0.5 lb. ai (4 pts.) per acre per year.
NON-CROPLAND (EXCLUDING PUBLIC LAND)				
	See Crop Outlets	See Crop Outlets	See Crop Outlets	<ul style="list-style-type: none"> Spray non-cropland adjacent to agricultural areas to control migratory insects, which may threaten crops. Follow General Use Directions, rates and spray recommendations found elsewhere in this label for the adjacent crop out and target pests. Use highest labeled rates for dense/large foliage, high insect populations and larger larval stages. Repeat as necessary to maintain control. Do not exceed 0.2 lb. ai (1.6 pints) per acre per year. Do not graze livestock in treated areas.

RATE CONVERSION CHART

Lb. A.I. Per Acre	Fl. Oz. Per Acre	Pts. Per Acre	Treated Acres Per Gal.
0.015	1.92	0.12	66
0.02	2.56	0.16	50
0.025	3.20	0.20	40
0.03	3.84	0.24	33
0.04	5.12	0.32	25

Karate® Insecticide, Karate® Insecticide With Zeon™ Technology, Warrior® Insecticide with Zeon™ Technology, E-Z Handler®, and the Syngenta logo are trademarks of a Syngenta Group Company

Ammo®, Capture®, Fury™, and Mustang® are trademarks of FMC Corporation

Asana® is a trademark of E. I. du Pont de Nemours & Co. (Inc.)

Baythroid® is a trademark of Mobay Corporation

Danitol® is a trademark of Sumitomo Chemical Co., LTD

DECIS® and SCOUT-XTRA® are registered trademarks of Hoechst Schering AgrEvo S.A.de C.V

SynerGin™ is a trademark of AgrEvo USA Company

©2002 Syngenta

<p>For non-emergency (e.g. current product information) call Syngenta Crop Protection at 1-800-334-9481.</p>
--

Product of the United Kingdom
Formulated in USA

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1112-M (Draft2.1)

WAR 1112 MAS (DRAFT 2.1CLEAN) – LG – 9-26-02