

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Dr. Jerry Wells Senior Regulatory Product Manager Regulatory Affairs, Herbicides Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, NC 27419-8300

NOV 2 2 2013

Subject: Adding Aerial Application for Corn and Sugarcane and Other Label Revisions

Approving Master Label + Two Supplemental Labels

PRIA Code: R350

Name: Callisto Herbicide Active Ingredient: Mesotrione

EPA Registration Number: 100-1131 Submission Date: January 8, 2013

Decision Number: 474131

Dear Dr. Wells:

The proposed amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) is <u>acceptable</u>. This action includes one approved master label and two approved supplemental labels.

This action registers aerial application to corn and sugarcane, changes perennial ryegrass grown for seed use to include both annual and perennial ryegrass grown for seed, revises the format for the rotational crop information, increases the number of applications and rate for grasses grown for seed, adds chemigation for cranberry, changes the Bush and Caneberries crop group to 13-07A and 13-07B, and modifies grain sorghum application timing by adding post-directed application for this mesotrione product.

Please note that while no additional data is being requested at this time, any marketing claims made on the pesticide label must be substantiated by data maintained in your files. If data supporting marketing claims made on the product label is not available then those claims must be removed.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on

Page 2 of 2

Adding Aerial Application for Corn and Sugarcane and Other Label Revisions

Name: Callisto Herbicide Active Ingredient: Mesotrione EPA Registration Number: 100-1131 Submission Date: January 8, 2013

Decision Number: 474131

the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Products released for shipment after 18 months from the date on this notice or the next printing of the label, whichever occurs first, must bear the new revised label must bear the new label language. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of this product constitutes acceptance of these conditions. As required, you must submit one copy of the final printed label before the product is released for shipment. If you have any questions please contact Michael Walsh by phone at (703) 308-2972 or via email at "walsh.michael@epa.gov".

Sincerely,

Kathryn V. Montague Product Manager (23)

Herbicide Branch

Registration Division (7505P)

them & Monk

Enclosure

[Booklet]

GROUP 27 HERBICIDE

Callisto® Herbicide

For Control of Annual Broadleaf Weeds in Field Corn, Seed Corn, Yellow Popcorn, Soybean, Sweet Corn, and Other Listed Crops

Active Ingredient:

Mesotrione: (CAS No. 104206	5-82-8)	40.0%
Other Ingredients:		60.0%
Total:		100.0%

Contains 4 lbs. of active ingredient mesotrione per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1131

EPA Est.

Callisto Plant Technology

SCP 1131

NOV 22 2013

100-1131

1 gallon 15 gallons ____ gallons Net Contents

FIRST AID			
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.		
·	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.		
	Call a poison control center or doctor for treatment advice.		
If on skin or	Take off contaminated clothing.		
clothing	Rinse skin immediately with plenty of water for 15-20 minutes.		
	Call a poison control center or doctor for treatment advice.		
If inhaled	Move person to fresh air.		
	 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to mouth, if possible. Call a poison control center or doctor for further treatment advice. 		
If swallowed	Call a poison control center or doctor immediately for treatment advice.		
•	 Have person sip a glass of water if able to swallow. 		
	 Do not induce vomiting unless told to by the poison control center or doctor. 		
	Do not give anything to an unconscious person.		
Have the product container or label with you when calling a poison control center or			
doctor, or going for treatment.			
_	HOTLINE NUMBER		
For 24-Hour Medical Emergency Assistance (Human or Animal),			
or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident)			
Call			
1-800-888-8372			

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes, or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.



Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves Category A (e.g. barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Physical and Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves Category A (e.g., barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils, or Viton ≥14 mils)

PRODUCT INFORMATION

Callisto is a systemic preemergence and postemergence herbicide for the selective contact and residual control of broadleaf weeds in field corn, seed corn, yellow popcorn, sweet corn, and other listed crops. When used preemergence, weeds take up the product through the soil during emergence. Dry conditions following application may reduce the preemergence activity of Callisto. If an activating rain (0.25 inches) is not received within 7-10 days after a preemergence application, where appropriate, rotary

hoeing is suggested to activate the herbicide. When used postemergence, susceptible weeds take up the herbicide through the treated foliage and cease growth soon after application. Complete death of the weeds may take up to 2 weeks. The product is absorbed through the soil and/or by the foliage of emerged weeds.

Callisto is not effective for the control of most grass weeds. Preemergence grass herbicides or postemergence grass herbicides can be tank mixed with Callisto to provide broad spectrum weed control in corn (see appropriate section of label for this information). Callisto can be applied postemergence following a preemergence grass herbicide application. Callisto can also be used in combination with a burndown herbicide, prior to planting, to provide added burndown and residual weed control in field corn, seed corn, yellow popcorn, and sweet corn.

RESISTANCE MANAGEMENT

Callisto is a **Group 27 Herbicide** (contains the active ingredient mesotrione).

Naturally occurring biotypes of certain broadleaf weed species with resistance to triazines, glyphosate, PPO, HPPD and ALS inhibiting herbicides are known to exist. Performance of Callisto is not affected by the presence of biotypes resistant to triazines, glyphosate, PPO or ALS inhibiting herbicides.

To prevent the risk of weeds developing resistance to Callisto in corn, always use full labeled rates. If applying Callisto postemergence after a mesotrione-containing preemergence herbicide, always add atrazine as a tank mix partner. No more than 0.24 lb. of mesotrione active ingredient must be applied per acre of corn per year (equivalent of 7.7 fl. oz. per acre per year of Callisto). If additional herbicide must be applied, it is recommended that a different mode of action be used, i.e., other than an HPPD inhibitor (Group 27 Herbicide). Callisto must be applied at full label rates to help prevent selection for, or population shifts toward, marginally tolerant weed species and/or species biotypes.

INTEGRATED PEST (WEED) MANAGEMENT

Callisto should be integrated into an overall weed and pest management strategy whenever the use of a herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional IPM strategies established for your area.

USE RESTRICTIONS AND PRECAUTIONS

Do not apply Callisto to white popcorn or ornamental (Indian) corn.

Severe corn injury resulting in yield loss may occur if Callisto is applied postemergence to corn that was treated with Counter® or Lorsban®.

Severe corn injury resulting in yield loss may occur if Callisto is applied foliar postemergence to corn in a tank mix with any organophosphate or carbamate insecticide.

Severe corn injury resulting in yield loss may occur if any organophosphate or carbamate insecticide is applied foliar postemergence within 7 days before or 7 days after Callisto application.

Do not cultivate corn within 7 days before or after a Callisto application as weed control from the Callisto application may be reduced.

When weeds are stressed due to drought, heat, lack of fertility, flooding, or prolonged cool temperatures, control can be reduced or delayed since the weeds are not actively growing. Weed escapes or regrowth may occur when application is made under prolonged stress conditions. Optimum weed control will be obtained if an application of Callisto is made following label directions when weeds are actively growing.

Do not apply this product through any type of irrigation system unless specified otherwise under the specific crop section on the label.

Do not apply this product with suspension fertilizers as the carrier.

Do not apply Callisto postemergence in a tank mix with emulsifiable concentrate grass herbicides, unless specifically addressed under one of the tank mix sections of this label, or injury may occur.

Do not use aerial application to apply Callisto unless specified otherwise under the specific crop section on the label.

Callisto may be applied with pyrethroid type insecticides (e.g., Warrior®).

SPRAY DRIFT PRECAUTIONS

Avoid drift onto adjacent crops and other nontarget areas.

RESTRICTION: For aerial application use only nozzles producing coarse-ultra coarse droplets. Do not use nozzles producing fine-medium size droplets.

Do not apply when weather conditions may cause drift to nontarget areas. Drift may result in injury to adjacent crops and vegetation. To avoid spray drift, DO NOT apply when wind speed is greater than 10 mph or during periods of temperature inversions. Use of larger droplet sizes will also reduce spray drift.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making a decision.

Information on Droplet Size

The most effective way to reduce spray drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions. Refer to the Aerial Application section for specific instructions regarding droplet size.

Controlling Droplet Size

- Volume Use high flow rate nozzles to apply the highest practical spray volume.
 Nozzles with higher rated flows produce larger droplets.
- Pressure Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher rate nozzles instead of increasing pressure.
- Number of Nozzles Use the minimum number of nozzles that provide uniform coverage.

Sensitive Areas

The pesticide must only be applied when the potential for drift to adjacent sensitive areas, (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

ADDITIONAL SPRAY DRIFT PRECAUTIONS FOR AERIAL APPLICATIONS

The distance of the outer-most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be observed.

Spray must be released at the lowest height consistent with effective weed control and flight safety.

For best results, each specific aerial application vehicle used should be quantifiably

pattern tested for aerial application of Callisto initially and every year thereafter.

RESTRICTION: For aerial application use only nozzles producing coarse-ultra coarse droplets. Do not use nozzles producing fine-medium size droplets.

For some use patterns, reducing the effective boom length to less than ¾ of the wingspan or rotor length may further reduce drift without reducing swath width.

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

APPLICATION INFORMATION

PREEMERGENCE GROUND APPLICATION

Apply Callisto preemergence with a carrier volume of 10-60 gals./A.

Spray nozzles must be uniformly spaced, the same size and type, and must provide accurate and uniform application. Use spray nozzles that provide medium to coarse droplet size to provide good coverage and avoid drift. Apply in a spray volume of 10-60 gals./A using water or liquid fertilizer (excluding suspension fertilizers) as the carrier. Use a pump that can maintain a pressure of at least 35-40 psi at the nozzles and provide proper agitation within the tank to keep the product dispersed. Lower pressures may be used with extended range or drift reduction nozzles.

Always ensure that agitation is maintained until spraying is completed, even if stopped for brief periods of time. If the agitation is stopped for more than 5 minutes, resuspend the spray solution by running on full agitation prior to spraying.

POSTEMERGENCE GROUND APPLICATION

Spray nozzles must be uniformly spaced, the same size and type, and must provide accurate and uniform application. Use spray nozzles that provide medium to coarse droplet size to provide good coverage and avoid drift. Good weed coverage is essential for optimum weed control. Boom height for broadcast over-the-top applications must be based on the height of the crop – at least 15 inches above the crop canopy.

Apply in a spray volume of 10-30 gals./A using water as a carrier. Use a pump that can maintain a pressure of at least 35-40 psi at the nozzles and provide proper agitation within the tank to keep the product dispersed. Lower pressures may be used with extended range or drift reduction nozzles. When weed foliage is dense, use a minimum of 20 gals.

Flat fan nozzles of 80° or 110° are recommended for optimum postemergence coverage. Do not use floodjet nozzles or controlled droplet application equipment for postemergence applications.

Nozzles may be angled forward 45° to enhance penetration of the crop and provide better coverage. Ensure that all in-line strainer and nozzle screens in the sprayer are 50-mesh or coarser.

Always ensure that agitation is maintained until spraying is completed, even if stopped for brief periods of time. If the agitation is stopped for more than 5 minutes, resuspend the spray solution by running on full agitation prior to spraying.

Aerial Application

RESTRICTION: Callisto can be applied aerially only to corn and sugarcane.

RESTRICTION: For aerial application use only nozzles producing coarse-ultra coarse droplets. Do not use nozzles producing fine-medium size droplets.

Callisto may be applied aerially for preemergence or postemergence weed control in corn only in the following states: Alabama, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Îdaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Nebraska, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin and Wyoming.

Callisto may be applied aerially for preemergence or postemergence weed control in sugarcane only in the following states: Florida, Louisiana and Texas.

Applications must be made in a minimum of 2 gallons of water per acre.

SPRAY ADDITIVES

POSTEMERGENCE ADJUVANTS

When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is recommended.

The following adjuvant recommendations are intended primarily for Callisto use in corn. Refer to the use directions section of each crop section for specific adjuvant recommendations.

POSTEMERGENCE APPLICATIONS TO FIELD CORN AND SEED CORN

For postemergence applications made after the crop has emerged, add crop oil concentrate (COC) to the spray solution at the rate of 1.0 gal./100 gals. of water (1.0% v/v). The use of a nonionic surfactant (NIS) at 1 qt./100 gallons of water (0.25% v/v) instead of COC is allowed, but the weed control achieved with COC is consistently better than NIS. The use of methylated seed oil (MSO) adjuvants or MSO blend adjuvants for postemergence applications of Callisto may cause severe crop injury to occur. Do not use MSO adjuvants for postemergence use unless directed for a specific tank mix under the CALLISTO TANK MIXTURES FOR CORN section of this label, or unless permitted by a supplemental Callisto label. In addition to COC, always add spray grade UAN (e.g., 28-0-0) to the spray solution at a

rate of 2.5% (v/v) or AMS at 8.5 lb./100 gals. of spray solution, except if precluded elsewhere on this label or by a supplemental Callisto label.

POSTEMERGENCE APPLICATIONS TO SWEET CORN AND YELLOW POPCORN

Do not add UAN or AMS when making postemergence applications of Callisto to yellow popcorn or sweet corn, or severe crop injury may occur.

For postemergence applications to yellow popcorn and sweet corn, the use of a nonionic surfactant (NIS) instead of a crop oil concentrate (COC) is recommended, so as to minimize the risk of crop injury. A COC may be used, and will increase the level of weed control achieved, especially under dry growing conditions, but the risk of crop injury is increased significantly under lush growing conditions. For optimum control, the addition of atrazine is recommended wherever rotational or local atrazine restrictions allow.

PREEMERGENCE ADJUVANTS

For Callisto preplant or preemergence applications, and where weeds are present, the use of any adjuvant for agricultural use is permitted. In these situations, MSO type adjuvants are typically better than COC type adjuvants, which are typically better than NIS type adjuvants for enhancing weed control. UAN or AMS can be added and typically provides better weed control than not adding one of these. If Callisto is being tank mixed with another registered herbicide in this situation, refer to the tank mix partner label for adjuvant precautions and restrictions.

SPRAY EQUIPMENT

Cleaning Equipment After Callisto Application

Special attention must be given to cleaning equipment before spraying a crop other than corn. Mix only as much spray solution as needed.

- 1. Flush tank, hoses, boom, and nozzles with clean water.
- 2. Prepare a cleaning solution of 1 gal. of household ammonia per 25 gals. of water. Many commercial spray tank cleaners may be used.
- 3. Use a pressure washer to clean the inside of the spray tank with this solution. Take care to wash all parts of the tank, including the inside top surface. If a pressure washer is not available, completely fill the sprayer with the cleaning solution to ensure contact of the cleaning solution with all internal surfaces of the tank and plumbing. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.

- 4. Flush hoses, spray lines, and nozzles for at least 1 minute with the cleaning solution.
- 5. Dispose of rinsate from steps 1-3 in an appropriate manner.
- 6. Repeat steps 2-5.
- 7. Remove nozzles, screens, and strainers and clean separately in the ammonia solution after completing the above procedures.
- 8. Rinse the complete spraying system with clean water.

MIXING PROCEDURES

Refer to the Crop Use Directions sections of this label for recommended tank mixes.

Always refer to labels of other pesticide products for mixing directions and precautions which may differ from those outlined here. Use in accordance with the most restrictive of label limitations and precautions. No label dosage rates may be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Do not tank mix Callisto with any other insecticide, fungicide, fertilizer solution, or adjuvant not recommended on the label without testing compatibility, as poor mixing may result. It is recommended that the compatibility of any tank mix combination be tested on a small scale such as a jar test before actual tank mixing.

Follow the mixing instructions for adding Callisto to the spray tank:

- 1. Only use sprayers in good running condition with good agitation. Ensure the sprayer is cleaned according to instructions on label of the product used prior to Callisto. For postemergence applications, use only clean water for the spray solution. Ensure that all in-line strainer and nozzle screens in the sprayer are 50-mesh or coarser. Screens finer than 50-mesh should not be used.
- 2. Liquid fertilizer (excluding suspension fertilizers) may be used as the carrier for preemergence applications.
- 3. Begin to fill sprayer tank or premix tank with clean water and engage agitator. Agitation must be continued throughout the entire mixing and spraying procedure.
- 4. When the sprayer or premix tank is half full of water, add AMS and agitate until completely dispersed.
- 5. Next add Callisto slowly and agitate until completely dissolved. Wait at least 1 minute after the last of the Callisto has been added to the tank to allow for

complete dispersion. A longer agitation period may be required to disperse Callisto when using cold water from sources such as deep drilled wells.

- 6. If tank mixing, add the tank mix product next.
- 7. Finally, add adjuvant and UAN, if needed, and then continue to fill tank to desired level with water.

WEEDS CONTROLLED

Callisto applied as directed in this label will control or partially control the weeds listed in Tables 1 and 2.

Where reference is made to weeds partially controlled, partial control can either mean erratic control (good to poor) or consistent control at a level below that generally considered acceptable for commercial weed control.

For best postemergence results, apply Callisto to actively growing weeds. Dry weather following preemergence application of Callisto may reduce residual weed control effectiveness. If irrigation is available, apply ½ to 1 inch of water after preemergence application. If irrigation is not available, a uniform shallow cultivation is recommended as soon as weeds emerge.

Callisto applied alone or in mixture with atrazine will not provide consistent or effective control of weeds identified as resistant to postemergence HPPD inhibiting herbicides.

Refer to the crop sections on this label for specific rates and use directions.

Table 1. Weeds Controlled With Postemergence Applications of Callisto

NA/	Manad	Callisto	Callisto 2.5-3.0 fl.
Weed Common Name	Weed Scientific Name	3 fl. oz./A	oz./A + Atrazine ¹ eeds <5 Inches Tall ²
Amaranth, palmer	Amaranthus palmeri	PC ³	C ³
Amaranth, powell	Amaranthus powellii	C	C
Amaranth, spiny	Amaranthus spinosus	C	C
Atriplex	Chenopodium orach	C	C
Broadleaf signalgrass	Urochloa platyphylla	$\frac{C_3}{C_3}$	C ³
Buckwheat, wild	Polygonum convolvulus	PC	PC
Buffalobur	Solanum rostratium	C	C
Burcucumber	Sicyos angulatus	PC	C ³
Carpetweed	Mollugo verticillata	C	C
Carrot, wild	Daucus carota	PC	C
Chickweed, common	Stellaria media	C	C
Cocklebur, common	Xanthium strumarium	C	C
Crabgrass, large	Digitaria sanguinalis	C_3	C ³
Dandelion	Taraxacum officinale	NC	PC
Dock, curly	Rumex crispus	PC	PC
Galinsoga	Galinsoga parviflora	C	C
Hemp	Cannabis sativa	C	C
Horsenettle	Solanum carolinense	PC	C
Jimsonweed	Datura stramonium	C	C
Horseweed (marestail)	Conyza canadensis	PC	C
Knotweed, prostrate	Polygonum aviculare	PC	PC
Kochia	Kochia scoparia	PC ³	C ³
Lambsquarters, common	Chenopodium album	C	C
Mallow, Venice	Hibiscus trionum	NC NC	C
Morningglory, entireleaf	Ipomoea hederacea	PC	C
Morningglory, ivyleaf	Ipomoea hederacea	PC	C
Morningglory, pitted	Ipomoea lacunosa	PC	C
Mustard, wild	Brassica kaber	C	C
Nightshade, black	Solanum nigrum	C	C
	 	C	
Nightshade, Eastern black	Solanum ptycanthum		C
Nightshade, hairy	Solanum sarrachoides	C	С
Nutsedge, yellow	Cyperus esculentus	PC	PC
Pigweed, redroot	Amaranthus retroflexus	C	С
Pigweed, smooth	Amaranthus hybridus	С	С

Weed	Weed	Callisto 3 fl. oz./A	Callisto 2.5-3.0 fl. oz./A + Atrazine ¹
Common Name	Scientific Name	Apply to Weeds <5 Inches Tall ²	
Pigweed, tumble	Amaranthus albus	С	С
Pokeweed, common	Phytolacca americana	PC	PC
Potatoes, volunteer	Solanum spp.	С	C
Pusley, Florida	Richardia scabra	C ³	C ³
Ragweed, common	Ambrosia artemisiifolia	PC	С
Ragweed, giant	Ambrosia trifida	C ³	С
Sesbania, hemp	Sesbania exaltata	С	С
Sida, prickly (teaweed)	Sida spinosa	NC	C ³
Smartweed, ladysthumb	Polygonum persicaria	C ₃	С
Smartweed, pale	Polygonum lapathifolium	C ₃	С
Smartweed, Pennsylvania	Polygonum pensylvanicum	C ³	С
Sunflower, common	Helianthus annuus	С	С
Thistle, Canada	Circium arvense	NC	PC
Velvetleaf	Abutilon theophrasti	С	С
Waterhemp, common	Amaranthus rudis	C ³	С
Waterhemp, tall	Amaranthus tuberculatus	C ³	С

¹Callisto tank mixture with atrazine is approved only for use on corn and sugarcane.

C = Control

PC = Partial Control

NC = Not Controlled

²Under certain situations weeds can be controlled at larger than listed sizes, however to protect crop yield, manage weed resistance and provide consistent control, treat weeds before they exceed 5 inches in height.

³Apply before weed exceeds 3 inches in height.

Table 2. Weeds Controlled With Preemergence Applications of Callisto

Common Name	Scientific Name	Callisto Applied Alone	Callisto + Atrazine ¹
Amaranth, palmer	Amaranthus palmeri	С	С
Amarath, powell	Amaranthus powellii	С	С
Amaranth, spiny	Amaranthus spinosus	C	С
Broadleaf signalgrass	Urochloa platyphylla	PC	PC
Buffalobur	Solanum rostratum	С	С
Carpetweed	Mollugo verticillata	С	С
Chickweed, common	Stellaria media	С	С
Cocklebur, common	Xanthium strumarium	PC	С
Crabgrass, large	Digitaria sanguinalis	PC	PC
Galinsoga	Galinsoga parviflora	С	С
Jimsonweed	Datura stramonium	С	С
Kochia	Kochia scoparia	PC	С
Lambsquarters, common	Chenopodium album	С	C
Morningglory, entireleaf	Ipomoea hederacea	PC	C .
Morningglory, ivyleaf	Ipomoea hederacea	PC	С
Morningglory, pitted	Ipomoea lacunosa	PC	С
Nightshade, eastern black	Solanum ptycanthum	С	С
Nightshade, hairy	Solanum sarrachoides	С	С
Pigweed, redroot	Amaranthus retroflexus	С	Ç
Pigweed, smooth	Amaranthus hybridus	С	С
Pigweed, tumble	Amaranthus albus	С	С
Ragweed, common	Ambrosia artemisiifolia	С	С
Ragweed, giant	Ambrosia trifida	PC	С
Smartweed, ladysthumb	Polygonum persicaria	С	С
Smartweed, pale	Polygonum lapathifolium	· C	С
Smartweed, Pennsylvania	Polygonum pensylvanicum	С	С
Sunflower, common	Helianthus annuus	PC	. C
Velvetleaf	Abutilon theophrasti	С	С
Waterhemp, common	Amaranthus rudis	С	С
Waterhemp, tall	Amaranthus tuberculatus	С	С

¹Callisto tank mixture with atrazine is approved only for use on corn grain sorghum and sugarcane. Refer to the crop sections on this label for specific use directions.

PC = Partial Control

C = Control

ROTATIONAL CROPS

When Callisto is applied as directed on this label, follow the crop rotation intervals in Table 3. If Callisto is tank mixed with other products, follow the most restrictive product's crop rotation interval.

Table 3. Time Interval Between Callisto Application and Replanting or Planting of Rotational Crop

Crop	Replant/Rotational Interval
Asparagus	
Corn (all types)	
Cranberry	
Flax	
Kentucky bluegrass grown for seed	
Millet, pearl	Anytima
Oats	Anytime
Rhubarb	
Ryegrass (perennial and annual) grown for seed	
Sorghum (grain and sweet)	
Sugarcane	•
Tall fescue grown for seed	
Small grain cereals including wheat, barley and rye	4 Months
Alfalfa	
Blueberry	
Canola	
Cotton	·
Currant	
Lingonberry	
Okra	
Peanuts	10 Months
Peas ^{1,2}	
Potato	
Rice	
Snap beans ^{1,2}	
Soybeans	
Sunflowers	
Tobacco	· ·
Cucurbits	
Dry beans	
Red clover	18 Months
Sugar beets	
All other rotational crops	

¹Plant these rotational crops only if the following criteria below have been met. If all criteria are not met, plant peas and snap beans a minimum of 18 months following Callisto application.

- A minimum of 20" of rainfall plus irrigation has been received between application and planting of the rotational crop.
- Soil pH is 6.0 or greater.
- Application of Callisto at 3 fl. oz./A or less applied no later than June 30th the year preceding rotational crop planting.
- No other HPPD herbicides (e.g., Callisto® Xtra, Halex® GT, Lexar® EZ, Lumax® EZ, Zemax®, Armezon™, Balance® Flexx, Capreno®, Corvus®, Impact®, or Laudis®) were applied the year prior to planting peas and snap beans.

²Do not plant peas or snap beans on sand, sandy loam or loamy sand soils in Minnesota or Wisconsin.

CROP USE DIRECTIONS

CORN

Callisto may be applied by ground for preemergence or postemergence weed control in field corn, seed corn, yellow popcorn, and sweet corn.

Callisto may also be applied aerially for preemergence or postemergence weed control only in the following states: Alabama, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin and Wyoming.

Refer to seed company recommendations for use on field corn inbred lines. Special adjuvant restrictions must be followed for postemergence applications of Callisto in yellow popcorn or sweet corn (see the **SPRAY ADDITIVES** section of this label). Do not apply Callisto to white popcorn or ornamental (Indian) corn.

Postemergence applications (after crop emergence) of Callisto may cause crop bleaching in some yellow popcorn and sweet corn hybrids. Crop bleaching is typically transitory and has no effect on final yield or quality. However, herbicide sensitivity in yellow popcorn and sweet corn varies widely, and all yellow popcorn and sweet corn hybrids have not been tested. Contact your popcorn or sweet corn company, Fieldman, or University Specialist about hybrid recommendations before making a postemergence application of Callisto to yellow popcorn or sweet corn. Do not include nitrogen based adjuvants (UAN or AMS) when making postemergence applications of Callisto to yellow popcorn or sweet corn.

Temporary crop response (transient bleaching) from postemergence applications to field corn may occur under extreme weather conditions or when the crop is suffering from stress. Field corn quickly outgrows these effects and develops normally.

Do not apply more than a total of 7.7 fl. oz. (0.24 lb. mesotrione active ingredient) of Callisto per acre per season. Do not make more than 2 applications of Callisto per season. Do not exceed 3.0 fl. oz. (0.094 lb. a.i./A) in a single postemergence application. Do not make the second application of Callisto within 14 days of the first application.

Apply Callisto for the control of broadleaf and grass weeds listed in Tables 1 and 2. Corn may be treated up to 30 inches tall or up to the 8-leaf stage of corn growth. Do not feed or harvest forage, grain, or stover within 45 days after application.

CALLISTO USED ALONE - POSTEMERGENCE

Apply Callisto at 3.0 fl. oz./A per application. Always add an appropriate adjuvant to the spray tank (see the **SPRAY ADDITIVES** section of this label).

For best results, apply Callisto to actively growing weeds. For a list of weeds controlled see Table 1. Susceptible weeds which emerge soon after application of Callisto may be controlled after they absorb the herbicide from the soil. Callisto will not control most grass weeds.

Two postemergence applications of Callisto may be made with the following restrictions.

- Only one postemergence application may be made if Callisto has been applied preemergence. Do not exceed a total of two applications per season. Do not exceed a total of 7.7 fl. oz./A (0.24 lb. a.i./A) of Callisto per season.
- Do not make the second application within 14 days of the first application.
- Application of Callisto at rates less than 3.0 fl. oz./A (0.094 lb. a.i./A) postemergence may result in incomplete weed control and loss of residual control.
- Do not exceed a total of 6.0 fl. oz./A (0.19 lb. a.i./A) for the two postemergence applications.
- If Callisto is applied postemergence to ground that received a preemergence application of a mesotrione-containing herbicide, atrazine must be tank mixed with Callisto.
- If atrazine is mixed with Callisto, do not apply to corn that is more than 12 inches in height.
- Corn may be treated up to 30 inches tall or up to the 8-leaf stage of corn growth. Do

not harvest forage, grain, or stover within 45 days after application.

CALLISTO USED ALONE - PREEMERGENCE

Apply Callisto alone at 6.0-7.7 fl. oz./A (0.188-0.24 lb. a.i,/A) by ground sprayers in a spray volume of 10-30 gals. of water (up to 80 gals. if applied with liquid fertilizers) per acre for broadleaf weed control. For a list of weeds controlled, refer to Table 2. Callisto may be tank mixed with preemergence grass herbicides for grass control. Refer to the tank mix section for a list of partners.

CALLISTO TANK MIXTURES FOR CORN

Callisto may be tank mixed with other registered herbicides for improved spectrum of weed control in burndown, preemergence or postemergence applications. Additionally these tank mixtures can be used to include a different mode of action herbicide to help control or manage the development of resistant weed biotypes.

Burndown Tank Mixtures in Corn

Callisto may be applied in tank mixture with other registered herbicides for burndown plus residual weed control.

For improved broadleaf weed control with limited residual control prior to planting corn and before corn emergence, apply Callisto at 3.0 fl. oz./A in tank mixes with Gramoxone® brands, Roundup® brands, Touchdown® brands, dicamba brands (e.g. Banvel®) and/or 2,4-D. For greater residual control, use 6.0-7.7 fl. oz./A of Callisto (see Table 2) with the above products. Use the adjuvant system recommended by the burndown herbicide. Refer to individual product labels for precautionary statements, restrictions, rates, approved uses, and a list of weeds controlled.

Preemergence Tank Mixtures in Corn

Callisto may be applied at a rate of 5.3-7.7 fl. oz./A in tank mixture with other registered herbicides (Table 4) for preemergence residual weed control. Refer to Table 2 for a list of weeds controlled by Callisto and Callisto plus AAtrex® applied preemergence.

Table 4. Callisto Tank Mixtures for Preemergence Application in Corn¹

AAtrex	Degree Xtra®	Harness Xtra® 5.6L
Bicep Lite II Magnum®	Dual II Magnum®	Keystone®
Bicep II Magnum®	Expert®	Keystone® LA
Cinch®	Fultime®	Outlook®
Cinch® ATZ	Guardsman Max®	Prowl®
Cinch® ATZ Lite	Harness®	Surpass® EC
Degree®	Harness Xtra®	TopNotch®

¹Refer to individual product labels for precautionary statements, restrictions, rates, approved uses, and a list of weeds controlled.

Postemergence Tank Mixtures in Corn

The tank mixtures with Callisto identified in Table 5 may be applied postemergence to corn (i.e., after corn has emerged). Unless specified otherwise on this label or a Syngenta supplemental label, do not apply Callisto at less than 3.0 fl. oz./A. Application of Callisto at rates less than 3.0 fl. oz. (0.094 lb. ai/A) postemergence may result in a loss of residual control.

Always add an appropriate adjuvant to the spray tank (see the **SPRAY ADDITIVES** section of this label). Refer to individual product labels for precautionary statements, restrictions, rates, approved uses, and a list of weeds controlled. Not all of the tank mix pesticides listed are registered for field corn, yellow popcorn, or sweet corn.

Table 5. Callisto Tank Mixtures for Postemergence Application in Corn

Tank-Mix Partners ¹	Directions
AAtrex® 4L AAtrex® Nine-O®	Refer to Table 1 on this label for application rates and weeds controlled.
Accent® Accent® Q	Use this mixture for additional grass control. Refer to product label for list of weeds controlled.
Basagran®	Use this mixture for additional broadleaf weed control. Refer to product label for list of weeds controlled.
Basis® Basis Gold®	Use this mixture for additional weed control. Refer to product label for list of weeds controlled.

Tank-Mix Partners ¹	Directions	
Bicep II Magnum Bicep Lite II Magnum	When using these tank mixtures, it is recommended to leave the nitrogen based adjuvant (UAN or AMS) out of the mixture or apply as a post-directed spray to minimize contact with crop foliage. To further reduce the risk of crop injury, the user may also leave out the crop oil concentrate (COC), or replace it with a nonionic surfactant (NIS). In all cases, the control of emerged weeds may be reduced somewhat due to less than optimum adjuvant effect or weed coverage.	
Buctril®	Use this mixture for additional broadleaf weed control.	
Moxy®	 Add Buctril (2 lb./gal.) or Moxy (2 lb./gal.) at a rate up to 6 fl. oz./A. Add Buctril (4 lb./gal.) at a rate up to 3 fl. oz./A. 	
Expert	 For use only in glyphosate tolerant corn (e.g. Agrisure® GT, Roundup Ready®). Application of this mixture to a corn hybrid that is not 	
	 glyphosate tolerant will result in crop death. Do not add urea ammonium nitrate (UAN) or methylated seed oil (MSO) type adjuvants to this tank mixture or crop injury may occur. 	
Ignite® Ignite® 280 SL	Use this tank mixture only on corn designated as LibertyLink® or warranted as being tolerant to glufosinate. Application of this mixture to a corn hybrid that is not glufosinate tolerant will result in severe crop injury or death. Do not use crop oil concentrate (COC) as an adjuvant for this mixture or severe crop injury may occur.	
Lightning®	 For use only on corn designated as Clearfield® corn or warranted by BASF as being tolerant to Lightning Herbicide. Application of this mixture to a corn hybrid that is not Lightning tolerant will result in severe crop injury or death. Do not use a Methylated Seed Oil (MSO), or an MSO blend with this mixture or severe crop injury may result. 	
Northstar®	Use this mixture for additional weed control. Refer to product label for list of weeds controlled.	
Peak®	Use this mixture for additional weed control. Refer to product label for list of weeds controlled.	
Spirit®	Use this mixture for additional weed control. Refer to product label for list of weeds controlled.	
Steadfast® Steadfast® ATZ Steadfast® Q	Use this mixture for additional weed control. Refer to product label for list of weeds controlled.	

Tank-Mix Partners ¹	Directions
Stout®	Use this mixture for additional weed control. Refer to product label for list of weeds controlled.
Touchdown Roundup Solo glyphosate products	 For use only in glyphosate tolerant corn (e.g. Agrisure GT, Roundup Ready). Application of this mixture to a corn hybrid that is not glyphosate tolerant will result in crop death. Add spray-grade ammonium sulfate (AMS) at a rate that delivers 8.5-17.0 lbs. of AMS/100 gallons of water. If the glyphosate product label calls for an adjuvant in addition to AMS, add a non-ionic surfactant (NIS) at 0.25-0.5% v/v (1-2 quart/100 gallons). Do not add urea ammonium nitrate (UAN), crop oil concentrate (COC), or methylated seed oil (MSO) type adjuvants to this tank mixture or crop injury may occur.

¹Refer to individual product labels for precautionary statements, restrictions, rates, approved uses, and a list of weeds controlled.

ASPARAGUS

Callisto can be applied broadcast or banded at a rate of 3.0-7.7 fl. oz./A to asparagus as a spring application prior to spear emergence, as a post-harvest application (after final harvest), or both.

Use the 3.0 fl. oz./A rate for postemergence control or partial control of the emerged weeds listed in Table 1. Use the 6.0-7.7 fl. oz./A rate for preemergence control or partial control of the weeds listed in Table 2. For banded applications, the application must be made to account for band width, i.e. to deliver 3.0-7.7 fl. oz. per treated acre. For the best preemergence weed control with spring applications, Callisto must be applied after fern mowing, disking or other tillage operation but prior to asparagus spear emergence.

When making post-harvest applications, the rate applied preemergence in the spring must be taken into account so as not to exceed the 7.7 fl. oz./A/year rate limit. Post-harvest applications must be made in a way that minimizes contact with any standing asparagus spears or ferns and maximizes contact with the weeds and/or soil, e.g. by using a directed or semi-directed type application, or crop injury may occur. With post-harvest applications, the use of an adjuvant will increase the risk of crop injury.

If weeds are emerged at the time of the Callisto application, the addition of a crop oil concentrate (COC) type adjuvant at the rate of 1% v/v or a nonionic surfactant (NIS) at the rate of 0.25% v/v is recommended. In addition to COC or NIS, a spray grade UAN (e.g. 28-0-0) at the rate of 2.5% v/v or ammonium sulfate (AMS) at the rate of 8.5 lb./100 gallons of spray solution may be added for improved burndown of emerged weeds. If weeds have not yet emerged, no adjuvant is recommended.

Restrictions:

- 1. Do not apply more than 7.7 fl. oz./A of Callisto per year.
- 2. Do not make more than two Callisto applications per year.

BLUEGRASS, RYEGRASS (ANNUAL AND PERENNIAL) AND TALL FESCUE GROWN FOR SEED

Callisto can be applied to bluegrass, annual ryegrass, perennial ryegrass, or tall fescue which is grown for seed. Callisto can be applied as a preemergence application to bare soil (new seeding) or as a postemergence application to an emerged grass crop.

Preemergence Application: Apply Callisto as a broadcast, surface spray at a rate of 6.0 fl. oz./A to a newly seeded crop. The Callisto application must be made prior to crop and weed emergence. Rainfall or irrigation as the newly seeded grass crop emerges from the soil may increase the risk of injury from Callisto. Grass crop injury symptoms include temporary bleaching of newly emerged leaves, or in extreme conditions, stunting. For a list of preemergence weeds controlled or partially controlled see Table 2. In addition to the weeds listed in Table 2, Callisto applied preemergence will control mannagrass.

Postemergence Application: Apply Callisto as a broadcast postemergence spray at a rate of 3.0-6.0 fl. oz./A to emerged bluegrass, perennial ryegrass or tall fescue grown for seed. Use the 3.0 fl. oz./A rate for postemergence control or partial control of the weeds listed in Table 1. In addition to the weeds listed in Table 2, Callisto applied postemergence will control mannagrass (up to 3 tillers).

Use the 6.0 fl. oz./A rate for postemergence weed control plus extended residual weed control (see Table 2). The addition of a crop oil concentrate type adjuvant at 1% v/v or a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v is recommended. Postemergence applications of Callisto may result in temporary bleaching of the grass crop.

In addition to COC or NIS, a spray grade UAN (e.g. 28-0-0) at the rate of 2.5% v/v or ammonium sulfate (AMS) at the rate of 8.5 lb./100 gallons of spray solution may also be added for improved control of emerged weeds. The addition of UAN or AMS will improve consistency of postemergence weed control but will also increase the risk of grass crop injury, especially at Callisto rates greater than 3.0 fl. oz./A. If grass crop injury is a concern, do not add UAN or AMS to the spray solution.

Tank mixing other pesticides with Callisto postemergence may increase the risk of crop injury. Avoid adding pesticides with emulsifiable concentrate (EC) type formulations to Callisto for applications made postemergence to the crop.

Restrictions:

- 1. Do not harvest the grass crop for seed or straw within 60 days following the application of Callisto.
- 2. Do not graze or feed forage from treated areas within 14 days following harvest of seed or straw and at least 74 days after application of Callisto.
- 3. Do not make more than two applications of Callisto per year.
- 4. Do not apply more than 6 fl. oz./A in a single application and not more than 9 fl. oz./A of Callisto per year.
- 5. Applications of Callisto to grasses grown for seed species not listed on this label may result in severe injury.

BUSH AND CANEBERRIES (CROP GROUP 13-07A and 13-07B)

Note: Not all cultivars and types of berries that are included within the Environmental Protection Agencies definition of bush and caneberries (Crop Subgroups 13-07A and 13-07B) have been tested and shown to have adequate crop safety to Callisto Herbicide. Those that have been tested, and are believed to be reasonably fit, are listed below along with use directions for that crop. If Callisto is used on bush or caneberries not listed below, severe crop injury may occur.

Callisto may be applied as a pre-bloom post-directed spray in high bush blueberry, lingonberry, red currant, black currant, black raspberry, red raspberry, and blackberry. For a list of weeds controlled see Tables 1 and 2. Callisto may be applied in bush or caneberries at a rate up to 6 fl. oz./A. If a split application weed control program is desired, 3 fl. oz./A followed by 3 fl. oz./A may be used, but no more than two applications per crop per year are allowed and not more than 6 fl. oz./A in total per year. If two applications are made, they must be made no closer than 14 days apart. The use of a crop oil concentrate (COC) type adjuvant at the rate of 1% v/v is recommended, but avoid using COC adjuvants that are injurious to bush or caneberry leaves. Do not apply Callisto to bush or caneberries after the onset of the bloom stage or illegal residues may occur.

In low bush blueberries, Callisto may only be applied in the non-bearing year. This application may be a broadcast application. Up to 6 fl. oz./A of Callisto may be applied in a single application, or 3 fl. oz./A followed by 3 fl. oz./A if used in a split application program. No more than two applications per year are allowed and not more than 6 fl. oz./A in total per year. If two applications are made, they must be made no closer than 14 days apart. The use of a crop oil concentrate (COC) type adjuvant at 1% v/v is recommended. Applications of Callisto during dry weather conditions and/or temperatures above 85° can cause injury to Lowbush blueberries. Applications of

Callisto can cause yellowing or necrosis of leaves and under severe conditions, leaf drop may occur especially on "Sourtop" variety blueberries.

CRANBERRY

Callisto may be applied to bearing or non-bearing cranberry beds for control or suppression of bog St. John's wort (*Hypericum boreala*), rushes (*Juncus canadensis*, *J. effuses*, *J. bufonlus*, *J. tenuis*), sedges spp. (*Carex* spp.), yellow loosestrife (*Lysimachia terrestris*) and silverleaf (*Potentilla pacifica*) in addition to the weeds listed in Tables 1 and 2. Callisto may be applied in cranberries at a rate up to 8 fl. oz./A. Apply no more than two applications per crop per year and not more than 16 fl. oz./A in total per year. If two applications are made, they must be made no closer than 14 days apart. The use of a crop oil concentrate (COC) type adjuvant at 1% v/v or non-ionic surfactant (NIS) at 0.25% v/v is recommended. Avoid using COC adjuvants that are injurious to cranberry leaves. In non-bearing cranberries, make the Callisto application(s) after the bud break stage, but not less than 45 days before flooding in fall or winter. In bearing cranberries, make the Callisto application(s) after the bud break stage, but not less than 45 days prior to flooding or harvest.

Callisto may be applied through irrigation systems (chemigation) including center pivot or solid set.

Chemigation – Sprinkler Irrigation Application for Cranberry Only

Check the irrigation system to ensure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period. Apply by injecting the recommended rate of Callisto herbicide into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target areas in 0.1-0.2 acre-inch of water. In general, use the least amount of water in this range required for proper distribution and coverage.

Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system. In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the recommended rate of Callisto herbicide for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

Chemigation Use Precautions – Sprinkler Irrigation Application

- 1. Apply this product only through sprinkler irrigation systems including center pivot or solid set. Do not apply this product through any other type of irrigation system.
- 2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

30/55

- 3. If you have any questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.
- 4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- 5. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.
- 6. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.
- 7. The pesticide injection pipeline must contain a functional, automatic, quickclosing check valve to prevent the flow of fluid back toward the injection pump.
- 8. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when pressure decreases to the point where pesticide distribution is adversely affected.
- 11. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and are capable of being fitted with a system interlock.
- 12. Any alternatives to the above required safety devices must conform to the list of EPA approved alternative devices.
- 13. Do not apply when wind speed favors drift beyond the area intended for treatment or nonuniform distribution of treated water.

Additional Restrictions: 1) Do not apply directly to water or areas where surface water is present outside the bog system. 2) Do not contaminate water when disposing of equipment wash water or rinsate. 3) Do not apply within 10 feet of surface water outside the bog system. 4) Do not spray to runoff.

FLAX

Callisto may be applied preemergence in flax, i.e. after planting but before crop emergence, at a rate up to 6 fl. oz./A. For a list of weeds controlled see Tables 1 and 2. Do not apply more than one application, and not more than 6 fl. oz./A, per crop or per year in flax. If weeds are emerged at the time of application, the use of a crop oil concentrate (COC) type adjuvant at the rate of 1% v/v is recommended. In addition, a spray grade UAN (e.g., 28-0-0) at the rate of 2.5% (v/v) or AMS at the rate of 8.5 lb./100 gals. of spray solution may be added to improve the burndown of existing weeds. Applications of Callisto to emerged flax can result in severe crop injury.

OATS

Callisto can be applied preemergence or postemergence (but not both) for weed control in oats.

For preemergence control or partial control of the weeds listed in Table 2, apply Callisto broadcast at a rate of 6.0 fl. oz./A prior to oat emergence. For best preemergence weed control, the Callisto application must be made prior to weed emergence.

For postemergence (after oat emergence) control or partial control of the weeds listed in Table 1, apply Callisto at a rate of 3.0 fl. oz./A. For best results, Callisto must be applied to emerged weeds that are less than 5" tall. Postemergence applications of Callisto may result in temporary injury of the oat crop. Injury symptoms may include leaf bleaching, leaf burn and in extreme conditions, stunting.

If emerged weeds are present at the time of the Callisto application, the addition of a crop oil concentrate (COC) type adjuvant at a rate of 1% v/v or a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v is recommended. In addition to COC or NIS, a spray grade UAN (e.g. 28-0-0) at the rate of 2.5% v/v or ammonium sulfate (AMS) at the rate of 8.5 lb./100 gallons of spray solution may be added for improved weed control. If emerged weeds are not present at the time of the Callisto application, no additives are recommended. If oat injury is a concern, eliminating the use of UAN or AMS will reduce the risk for postemergence crop injury. Additionally, the use of NIS instead of COC will also reduce the oat injury risk. However, weed control is also reduced if UAN or AMS is eliminated and when switching from COC to NIS.

Tank mixing other pesticides with Callisto postemergence may increase the risk of injury. Avoid adding pesticides with emulsifiable concentrate (EC) type formulations to Callisto for applications made postemergence to the crop.

Restrictions:

- 1. Do not graze or feed forage from treated areas within 30 days following an application of Callisto.
- 2. Do not harvest oats within 50 days following the application of Callisto.
- 3. Do not make more than one application of Callisto per year.
- 4. Do not apply Callisto preemergence (prior to oat emergence) at more than 6.0 fl. oz./A/year.
- 5. Do not apply Callisto postemergence at more than 3.0 fl. oz./A/year.
- 6. If the oat crop treated with Callisto is lost or destroyed, oats may be replanted immediately. If Callisto was applied to the lost oat crop, no additional Callisto can be applied to the replanted oat crop.

OKRA

Callisto can be applied as a row-middle or a hooded post-direct treatment (but not both) for weed control in okra.

Preemergence row-middle application: Apply Callisto at a rate of 6.0 fl. oz./A as a banded application to the row middles prior to weed emergence. For this banded application, leave one foot of untreated area over the okra row or 6" to each side of the planted row. For banded applications, the application must be made to account for band width, i.e. to deliver 6.0 fl. oz. per treated acre. Do not apply Callisto directly over the planted okra row or severe crop injury may occur. Injury risk is greatest on coarse textured soils (sand, sandy loam or loamy sand).

Postemergence hooded application: Apply Callisto at a rate of 3.0 fl. oz./A as a postemergence directed application using a hooded sprayer for control or partial control of the weeds listed in Table 1. Okra must be at least 3" tall at the time of this application. It is recommended that a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v be added to the spray solution. For postemergence hooded applications, the spray equipment must be set up to minimize the amount of Callisto that contacts the okra foliage or crop injury will occur. For best postemergence results, Callisto must be applied to actively growing weeds.

Restrictions:

- 1. Do not harvest okra within 28 days following the application of Callisto.
- 2. Do not make more than one application of Callisto per okra crop.

- 3. Do not apply Callisto as a row-middle application at more than 6.0 fl. oz. per treated acre per year.
- 4. Do not apply Callisto as a post-directed application at more than 3.0 fl. oz. per acre per year.
- 5. Do not apply Callisto as a broadcast preemergence or broadcast postemergence application to okra or severe injury will occur.
- 6. If the okra crop treated with Callisto is lost or destroyed, okra can be replanted only in the soil band that was not treated with Callisto.

PEARL MILLET

Callisto may be applied preemergence in pearl millet, i.e. after planting but before crop emergence, at a rate up to 6 fl. oz./A. For a list of weeds controlled see Table 2. Do not apply more than one application, and not more than 6 fl. oz./A per crop or per year in pearl millet. If weeds are emerged at the time of application, the use of a crop oil concentrate (COC) type adjuvant at the rate of 1% v/v is recommended. In addition, a spray grade UAN (e.g., 28-0-0) at the rate of 2.5% (v/v) or AMS at the rate of 8.5 lb./100 gals. of spray solution may be added to improve the burndown of existing weeds. Applications of Callisto to emerged pearl millet can result in severe crop injury.

RHUBARB

Callisto can be applied prior to crop emergence for weed control in established rhubarb.

Apply Callisto at a rate of 6.0 fl. oz./A to dormant (prior to any spring green-up) rhubarb for control or partial control of the weeds listed in Table 2. If weeds are emerged at the time of application, it is recommended that a crop oil concentrate (COC) type adjuvant at 1% v/v or a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v be added to the spray solution. Applications of Callisto to rhubarb that is not dormant may result in a temporary bleaching symptomology. Rainfall or irrigation after the Callisto application may increase the risk of injury to emerging rhubarb.

Restrictions:

- 1. Do not harvest rhubarb within 21 days following the application of Callisto.
- 2. Do not make more than one application of Callisto per year.
- 3. Do not apply Callisto at more than 6.0 fl. oz./A/year.

SORGHUM (GRAIN AND SWEET)

Preemergence Application: Callisto can be applied preemergence or preplant non-incorporated up to 21 days before planting sorghum for control or partial control of the weeds listed in Table 2.

Apply Callisto preemergence at a rate of 6.0-6.4 fl. oz./A as a broadcast non-incorporated application prior to sorghum emergence. Applying Callisto less than 7 days before sorghum planting will increase the risk of crop injury, especially if irrigation or rainfall is received following the application. Injury symptoms include temporary bleaching of newly emerging sorghum leaves. Applying Callisto more than 7 days (but not more than 21) prior to planting will reduce the risk of crop injury.

If Callisto is applied prior to planting, minimize disturbance of the herbicide treated soil barrier during the planting process in order to lessen the potential for weed emergence.

If emerged weeds are present at the time of the preemergence application, it is recommended that a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v or a crop oil concentrate (COC) type adjuvant at a rate of 1% v/v be added to the spray solution. In addition to COC or NIS, a spray grade UAN at a rate of 2.5% v/v or ammonium sulfate (AMS) at a rate of 8.5 lb./100 gallons of spray solution can be added to the spray solution.

Preemergence Application Restrictions:

- 1. Do not apply more than 6.4 fl. oz./A of Callisto per year.
- 2. Do not apply Callisto to emerged sorghum or severe crop injury may occur.
- 3. Do not use Callisto in the production of forage sorghum, sudangrass, sorghum-sudangrass hybrids, or dual purpose sorghum.
- 4. Do not apply Callisto to sorghum that is grown on coarse textured soils (e.g. sandy loam, loamy sand, sand).
- 5. In the State of Texas, do not apply Callisto to sorghum grown south of Interstate 20 (I-20) or east of Highway 277.

Post-Directed: Callisto can be applied post-directed to grain sorghum for control or partial control of the weeds listed in Table 1. For best results, apply Callisto to actively growing weeds.

35 55

Apply Callisto at a rate of 3 fl. oz./A as a post-directed application when the grain sorghum is a minimum of 8 inches tall. Make the application by directing the spray between the crop rows and towards the base of the grain sorghum plant. Direct application of Callisto onto grain sorghum foliage can result in crop injury including temporary bleaching. If crop injury does occur, newly emerging leaves following application are typically unaffected.

It is recommended that a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v or a crop oil concentrate (COC) type adjuvant at a rate of 1% v/v be added to the spray solution. In addition to COC or NIS, a spray grade Urea Ammonium Nitrate (UAN) at a rate of 2.5% v/v or ammonium sulfate (AMS) at a rate of 8.5 lb./100 gallons of spray solution can be added to the spray solution.

Callisto may be tank mixed with other herbicides registered for grain sorghum for improved spectrum of weed control. Additionally, these tank mixtures can be used to include a herbicide with a different mode of action to help control or manage the development of resistant weed biotypes.

Post-Directed Restrictions:

- 1. Do not apply more than one post-directed application of Callisto.
- 2. Do not apply more than 3.0 fl. oz./A of Callisto post-directed and not more than 6.4 fl. oz./A of Callisto per grain sorghum crop year.
- 3. Do not apply Callisto broadcast over-the-top to emerged sorghum or severe crop injury may occur.
- 4. Do not harvest grain sorghum for forage for 30 days following application.
- 5. Do not harvest for grain or stover for 60 days following application.
- 6. Do not apply Callisto after the sorghum seedhead has begun to emerge.
- 7. Do not use Callisto in the production of forage sorghum, sudangrass, or sorghum-sudangrass hybrids.

SOYBEAN

Callisto can be applied preemergence to soybeans that are identified as mesotrione tolerant. Applications to soybeans that are not mesotrione tolerant will result in significant crop injury. For a list of mesotrione tolerant soybean varieties, contact a Syngenta Technical Representative.

Preemergence Application: For preemergence control of the weeds listed in Table 2, apply Callisto prior to soybean emergence at a rate of 6.0 fl. oz./A. Apply the higher

36/55

rate for longer residual control. Callisto may be tank mixed with other registered soybean herbicides such as Dual Magnum®, Dual II Magnum, and Prefix®. Refer to the tank mix partner label and follow all precautions and restrictions.

If weeds are emerged at the time of application, add either a non-ionic surfactant (NIS) at 1 qt./100 gallons (0.25% v/v) or a crop oil concentrate (COC) at 1 gallon/100 gallons (1% v/v). In addition to NIS or COC, it is also recommended to add either ammonium sulfate (AMS) at 8.5-17 lbs./100 gallon (or equivalent).

Restrictions:

- 1. Apply no more than 6.0 fl. oz./A per soybean crop per year.
- 2. Do not apply Callisto to emerged soybeans.
- 3. Do not graze or feed soybean forage or hay to livestock.

SUGARCANE

Callisto can be applied by ground for preemergence, postemergence over-the-top or postemergence directed weed control in sugarcane.

Callisto may also be applied aerially for preemergence or postemergence weed control only in the following states: Florida, Louisiana and Texas.

Preemergence Applications: Apply Callisto for preemergence weed control at 6.0-7.7 fl. oz./A after the planting of plant-cane or after harvest of ratoon-cane. For a list of weeds controlled preemergence, refer to Table 2. If some weeds are already emerged at the time of application, add a crop oil concentrate (COC) type adjuvant at a rate of 1% v/v or a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v to the spray solution. In addition to COC or NIS, a spray grade UAN at a rate of 2.5% v/v or ammonium sulfate (AMS) at a rate of 8.5 lb./100 gallons of spray solution can be added to the spray solution. For improved preemergence weed control, AAtrex or Evik® can be tank mixed with Callisto. Refer to the tank mix partner label for specific rates and use directions.

Postemergence Applications: Apply Callisto postemergence at 3.0 fl. oz./A for control of the weeds listed in Table 1. Postemergence applications may be made as a post-over-the-top or as a post-directed spray to the base of the sugarcane. If a preemergence application was made earlier in the season, only one postemergence application can be made. If no preemergence application was made earlier in the season, both a post-over-the-top and a post-directed application can be made. For best results, Callisto must be applied to actively growing weeds.

For postemergence applications, it is recommended that a crop oil concentrate (COC) type adjuvant at a rate of 1% v/v or a nonionic surfactant (NIS) type adjuvant be added

37/55

to the spray solution. In addition to COC or NIS, the use of a spray grade UAN (e.g. 28-0-0) at 2.5% v/v <u>or</u> ammonium sulfate (AMS) at a rate of 8.5 lb./100 gallons of spray solution can be added for improved control of weeds.

For additional postemergence weed control, Callisto can be tank mixed with atrazine, Asulox® and/or Envoke®. Refer to the tank mix product labels for specific rates and use directions.

Restrictions:

- 1. Do not apply more than 7.7 fl. oz./A of Callisto as a preemergence application.
- 2. Do not apply more than 3.0 fl. oz./A of Callisto in a postemergence application.
- 3. Do not make more than two applications of Callisto per year. If a preemergence application of Callisto is made, only one postemergence application is allowed.
- 4. Do not make two Callisto applications less than 14 days apart.
- 5. Do not apply more than 10.7 fl. oz./A of Callisto per year.
- 6. Do not harvest sugarcane within 114 days following a post-over-the-top application of Callisto (114 day PHI).
- 7. Do not harvest sugarcane within 100 days following a post-directed application of Callisto (100 day PHI).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Keep container tightly closed when not in use. Do not store near seed, fertilizers, or foodstuffs. Can be stored at temperatures as low as -20°F. Keep away from heat and flame.

Pesticide Disposal

Open dumping is prohibited. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling [Less Than or Equal to 5 Gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining

contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If

Container Handling [Greater Than 5 Gallons]

burned, stay out of smoke.

Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR

IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

AAtrex®, AAtrex® Nine-O®, Agrisure®, Bicep II Magnum®, Bicep Lite II Magnum®, Callisto®, Callisto® Xtra, Callisto Plant Technology®, Dual Magnum®, Dual II Magnum®, Envoke®, Evik®, Expert®, Gramoxone®, Halex® GT, Lexar® EZ, Lumax® EZ, Northstar®, Peak®, Prefix®, Spirit®, Touchdown®, Warrior®, Zemax®, and the SYNGENTA Logo are Trademarks of a Syngenta Group Company

Accent®, Accent® Q, Basis®, Basis® Gold, Cinch®, Cinch® ATZ, Cinch® ATZ Lite, Steadfast®, Steadfast® ATZ, Steadfast® Q, Stout®, and Viton® are trademarks of E. I. du Pont de Nemours and Company

Asulox® is a trademark of United Phosphorous, Inc.

Balance® Flexx, Buctril®, Capreno®, Corvus®, Ignite®, Ignite® 280 SL, Laudis®, and LibertyLink® are trademarks of Bayer CropScience

Armezon™, Banvel®, Basagran®, Clearfield®, Counter®, Guardsman Max®, Lightning®, Outlook®, and Prowl® are trademarks of BASF Corporation

Degree®, Degree Xtra®, Harness®, Harness® Xtra, Harness® Xtra 5.6L, Roundup®, and Roundup Ready® are trademarks of Monsanto Company

Fultime®, Keystone®, Keystone® LA, Lorsban®, Surpass® EC, and TopNotch® are trademarks of Dow AgroSciences

Impact® is a trademark of Amvac Chemical Corporation

Moxy® is a trademark of Winfield Solutions, LLC

©201X Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for: Syngenta Crop Protection, LLC P. O. Box 18300 Greensboro, North Carolina 27419-8300

SCP 1131A

[Base Label]

GROUP 27 HERBICIDE

Callisto® Herbicide

For Control of Annual Broadleaf Weeds in Field Corn, Seed Corn, Yellow Popcorn, Soybean, Sweet Corn, and Other Listed Crops

Active Ingredient:

 Mesotrione: (CAS No. 104206-82-8)
 40.0%

 Other Ingredients:
 60.0%

 Total:
 100.0%

Contains 4 lbs. of active ingredient mesotrione per gallon.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1131

EPA Est.



SCP 1131A-

1 gallon Net Contents

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements, pesticide storage and disposal statements, and directions for use inside booklet.

Precautionary Statement Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes, or clothing.

FIRST AID	
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	Call a poison control center or doctor for treatment advice.
If on skin or	Take off contaminated clothing.
clothing	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air.
·	 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
	 Call a poison control center or doctor for further treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment
	advice.
	Have person sip a glass of water if able to swallow.
	 Do not induce vomiting unless told to by the poison control center or doctor.
	Do not give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or	
doctor, or going for treatment.	
HOTLINE NUMBER	
For 24-Hour Medical Emergency Assistance (Human or Animal),	
or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident)	
Call	
1-800-888-8372	

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Physical and Chemical Hazards

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Keep container tightly closed when not in use. Do not store near seed, fertilizers, or foodstuffs. Can be stored at temperatures as low as -20°F. Keep away from heat and flame.

Pesticide Disposal

Open dumping is prohibited. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling [Less Than or Equal to 5 Gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Handling [Greater Than 5 Gallons]

Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Bulk]

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. To THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY,

CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE)
RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE
RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION
OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Callisto®, Callisto Plant Technology®, and the Syngenta logo are trademarks of a Syngenta Group Company

©201X Syngenta

Manufactured for: Syngenta Crop Protection, LLC P. O. Box 18300 Greensboro, North Carolina 27419-8300

SCP 1131A

CAL 1131 MAS 1211 AMEND-E 0113 - bb - 11-8-13

000100.01131.20130108E.CALLISTO-AMEND-JAN2013.pdf

SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

GROUP 27 HERBICIDE

Callisto®

Herbicide

This supplemental label expires on October 30, 2016 and must not be used or distributed after this date.

Active Ingredient:

Other Ingredients:

60.0%

Total:

100.0%

Contains 4 lbs. of active ingredient mesotrione per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1131

NOV 22 2013

100-1131

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Callisto Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

ADDITIONAL SPRAY DRIFT PRECAUTIONS FOR AERIAL APPLICATIONS

The distance of the outer-most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be observed.

Spray must be released at the lowest height consistent with effective weed control and flight safety.

For best results, each specific aerial application vehicle used should be quantifiably pattern tested for aerial application of Callisto initially and every year thereafter.

RESTRICTION: For aerial application use only nozzles producing coarse-ultra coarse droplets. Do not use nozzles producing fine-medium size droplets.

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable

directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

APPLICATION INFORMATION

Aerial Application

RESTRICTION: Callisto can be applied aerially only to corn and sugarcane.

RESTRICTION: For aerial application use only nozzles producing coarse-ultra coarse droplets. Do not use nozzles producing fine-medium size droplets.

Callisto may be applied aerially for preemergence or postemergence weed control in corn only in the following states: Alabama, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Nebraska, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin and Wyoming.

Callisto may be applied aerially for preemergence or postemergence weed control in sugarcane only in the following states: Florida, Louisiana and Texas.

Applications must be made in a minimum of 2 gallons of water per acre.

BLUEGRASS, RYEGRASS (ANNUAL AND PERENNIAL) AND TALL FESCUE GROWN FOR SEED

Callisto can be applied to bluegrass, annual ryegrass, perennial ryegrass, or tall fescue which is grown for seed. Callisto can be applied as a preemergence application to bare soil (new seeding) or as a postemergence application to an emerged grass crop.

Preemergence Application: Apply Callisto as a broadcast, surface spray at a rate of 6.0 fl. oz./A to a newly seeded crop. The Callisto application must be made prior to crop and weed emergence. Rainfall or irrigation as the newly seeded grass crop emerges from the soil may increase the risk of injury from Callisto. Grass crop injury symptoms include temporary bleaching of newly emerged leaves, or in extreme conditions, stunting. For a list of preemergence weeds controlled or partially controlled see Table 2. In addition to the weeds listed in Table 2, Callisto applied preemergence will control mannagrass.

Postemergence Application: Apply Callisto as a broadcast postemergence spray at a rate of 3.0-6.0 fl. oz./A to emerged bluegrass, perennial ryegrass or tall fescue grown for seed. Use the 3.0 fl. oz./A rate for postemergence control or partial control of the weeds listed in Table 1. In addition to the weeds listed in Table 2, Callisto applied postemergence will control mannagrass (up to 3 tillers).

Use the 6.0 fl. oz./A rate for postemergence weed control plus extended residual weed control (see Table 2). The addition of a crop oil concentrate type adjuvant at 1% v/v or a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v is recommended. Postemergence applications of Callisto may result in temporary bleaching of the grass crop.

In addition to COC or NIS, a spray grade UAN (e.g. 28-0-0) at the rate of 2.5% v/v or ammonium sulfate (AMS) at the rate of 8.5 lb./100 gallons of spray solution may also be added for improved control of emerged weeds. The addition of UAN or AMS will improve consistency of postemergence weed control but will also increase the risk of grass crop injury, especially at Callisto rates greater than 3.0 fl. oz./A. If grass crop injury is a concern, do not add UAN or AMS to the spray solution.

Tank mixing other pesticides with Callisto postemergence may increase the risk of crop injury. Avoid adding pesticides with emulsifiable concentrate (EC) type formulations to Callisto for applications made postemergence to the crop.

Restrictions:

- 1. Do not harvest the grass crop for seed or straw within 60 days following the application of Callisto.
- 2. Do not graze or feed forage from treated areas within 14 days following harvest of seed or straw and at least 74 days after application of Callisto.
- 3. Do not make more than two applications of Callisto per year.
- 4. Do not apply more than 6 fl. oz./A in a single application and not more than 9 fl. oz./A of Callisto per year.

50/55

5. Applications of Callisto to grasses grown for seed species not listed on this label may result in severe injury.

BUSH AND CANEBERRIES (CROP GROUP 13-07A and 13-07B)

Note: Not all cultivars and types of berries that are included within the Environmental Protection Agencies definition of bush and caneberries (Crop Subgroups 13-07A and 13-07B) have been tested and shown to have adequate crop safety to Callisto Herbicide. Those that have been tested, and are believed to be reasonably fit, are listed below along with use directions for that crop. If Callisto is used on bush or caneberries not listed below, severe crop injury may occur.

Callisto may be applied as a pre-bloom post-directed spray in high bush blueberry, lingonberry, red currant, black currant, black raspberry, red raspberry, and blackberry. For a list of weeds controlled see Tables 1 and 2. Callisto may be applied in bush or caneberries at a rate up to 6 fl. oz./A. If a split application weed control program is desired, 3 fl. oz./A followed by 3 fl. oz./A may be used, but no more than two applications per crop per year are allowed and not more than 6 fl. oz./A in total per year. If two applications are made, they must be made no closer than 14 days apart. The use of a crop oil concentrate (COC) type adjuvant at the rate of 1% v/v is recommended, but avoid using COC adjuvants that are injurious to bush or caneberry leaves. Do not apply Callisto to bush or caneberries after the onset of the bloom stage or illegal residues may occur.

In low bush blueberries, Callisto may only be applied in the non-bearing year. This application may be a broadcast application. Up to 6 fl. oz./A of Callisto may be applied in a single application, or 3 fl. oz./A followed by 3 fl. oz./A if used in a split application program. No more than two applications per year are allowed and not more than 6 fl. oz./A in total per year. If two applications are made, they must be made no closer than 14 days apart. The use of a crop oil concentrate (COC) type adjuvant at 1% v/v is recommended. Applications of Callisto during dry weather conditions and/or temperatures above 85° can cause injury to Lowbush blueberries. Applications of Callisto can cause yellowing or necrosis of leaves and under severe conditions, leaf drop may occur especially on "Sourtop" variety blueberries.

CRANBERRY

Callisto may be applied to bearing or non-bearing cranberry beds for control or suppression of bog St. John's wort (*Hypericum boreala*), rushes (*Juncus canadensis*, *J. effuses*, *J. bufonlus*, *J. tenuis*), sedges spp. (*Carex* spp.), yellow loosestrife (*Lysimachia terrestris*) and silverleaf (*Potentilla pacifica*) in addition to the weeds listed in Tables 1 and 2. Callisto may be applied in cranberries at a rate up to 8 fl. oz./A. Apply no more than two applications per crop per year and not more than 16 fl. oz./A in total per year. If two applications are made, they must be made no closer than 14 days apart. The use of a crop oil concentrate (COC) type adjuvant at 1% v/v or non-ionic surfactant (NIS) at 0.25% v/v is recommended. Avoid using COC adjuvants that are

injurious to cranberry leaves. In non-bearing cranberries, make the Callisto application(s) after the bud break stage, but not less than 45 days before flooding in fall or winter. In bearing cranberries, make the Callisto application(s) after the bud break stage, but not less than 45 days prior to flooding or harvest.

Callisto may be applied through irrigation systems (chemigation) including center pivot or solid set.

Chemigation – Sprinkler Irrigation Application for Cranberry Only

Check the irrigation system to ensure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period. Apply by injecting the recommended rate of Callisto herbicide into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target areas in 0.1-0.2 acre-inch of water. In general, use the least amount of water in this range required for proper distribution and coverage.

Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system. In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the recommended rate of Callisto herbicide for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

Chemigation Use Precautions - Sprinkler Irrigation Application

- 1. Apply this product only through sprinkler irrigation systems including center pivot or solid set. Do not apply this product through any other type of irrigation system.
- 2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- 3. If you have any questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts.
- 4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- 5. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

- 6. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow.
- 7. The pesticide injection pipeline must contain a functional, automatic, quickclosing check valve to prevent the flow of fluid back toward the injection pump.
- 8. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when pressure decreases to the point where pesticide distribution is adversely affected.
- 11. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and are capable of being fitted with a system interlock.
- 12. Any alternatives to the above required safety devices must conform to the list of EPA approved alternative devices.
- 13. Do not apply when wind speed favors drift beyond the area intended for treatment or nonuniform distribution of treated water.

Additional Restrictions: 1) Do not apply directly to water or areas where surface water is present outside the bog system. 2) Do not contaminate water when disposing of equipment wash water or rinsate. 3) Do not apply within 10 feet of surface water outside the bog system. 4) Do not spray to runoff.

Callisto® and the Syngenta logo trademarks of a Syngenta Group Company

©201X Syngenta

CALLISTO 1131A S6-D - bb - 11-8-13

SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC

P. O. Box 18300

Greensboro, North Carolina 27419-8300

GROUP 27 HERBICIDE

Callisto®

Herbicide

This supplemental label expires on October 30, 2016 and must not be used or distributed after this date.

Active Ingredient:

Other Ingredients:

60.0%

Total:


100.0%

Contains 4 lbs. of active ingredient mesotrione per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1131



All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Callisto Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

-2-

SORGHUM (GRAIN AND SWEET)

Preemergence Application: Callisto can be applied preemergence or preplant non-incorporated up to 21 days before planting sorghum for control or partial control of the weeds listed in Table 2.

Apply Callisto preemergence at a rate of 6.0-6.4 fl. oz./A as a broadcast non-incorporated application prior to sorghum emergence. Applying Callisto less than 7 days before sorghum planting will increase the risk of crop injury, especially if irrigation or rainfall is received following the application. Injury symptoms include temporary bleaching of newly emerging sorghum leaves. Applying Callisto more than 7 days (but not more than 21) prior to planting will reduce the risk of crop injury.

If Callisto is applied prior to planting, minimize disturbance of the herbicide treated soil barrier during the planting process in order to lessen the potential for weed emergence.

If emerged weeds are present at the time of the preemergence application, it is recommended that a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v or a crop oil concentrate (COC) type adjuvant at a rate of 1% v/v be added to the spray solution. In addition to COC or NIS, a spray grade UAN at a rate of 2.5% v/v or ammonium sulfate (AMS) at a rate of 8.5 lb./100 gallons of spray solution can be added to the spray solution.

Preemergence Application Restrictions:

- 1. Do not apply more than 6.4 fl. oz./A of Callisto per year.
- 2. Do not apply Callisto to emerged sorghum or severe crop injury may occur.
- 3. Do not use Callisto in the production of forage sorghum, sudangrass, sorghum-sudangrass hybrids, or dual purpose sorghum.
- 4. Do not apply Callisto to sorghum that is grown on coarse textured soils (e.g. sandy loam, loamy sand, sand).
- 5. In the State of Texas, do not apply Callisto to sorghum grown south of Interstate 20 (I-20) or east of Highway 277.

Post-Directed: Callisto can be applied post-directed to grain sorghum for control or partial control of the weeds listed in Table 1. For best results, apply Callisto to actively growing weeds.

55 | 55

Apply Callisto at a rate of 3 fl. oz./A as a post-directed application when the grain sorghum is a minimum of 8 inches tall. Make the application by directing the spray between the crop rows and towards the base of the grain sorghum plant. Direct application of Callisto onto grain sorghum foliage can result in crop injury including temporary bleaching. If crop injury does occur, newly emerging leaves following application are typically unaffected.

It is recommended that a nonionic surfactant (NIS) type adjuvant at a rate of 0.25% v/v or a crop oil concentrate (COC) type adjuvant at a rate of 1% v/v be added to the spray solution. In addition to COC or NIS, a spray grade Urea Ammonium Nitrate (UAN) at a rate of 2.5% v/v or ammonium sulfate (AMS) at a rate of 8.5 lb./100 gallons of spray solution can be added to the spray solution.

Callisto may be tank mixed with other herbicides registered for grain sorghum for improved spectrum of weed control. Additionally, these tank mixtures can be used to include a herbicide with a different mode of action to help control or manage the development of resistant weed biotypes.

Post-Directed Restrictions:

- 1. Do not apply more than one post-directed application of Callisto.
- 2. Do not apply more than 3.0 fl. oz./A of Callisto post-directed and not more than 6.4 fl. oz./A of Callisto per grain sorghum crop year.
- 3. Do not apply Callisto broadcast over-the-top to emerged sorghum or severe crop injury may occur.
- 4. Do not harvest grain sorghum for forage for 30 days following application.
- 5. Do not harvest for grain or stover for 60 days following application.
- 6. Do not apply Callisto after the sorghum seedhead has begun to emerge.
- 7. Do not use Callisto in the production of forage sorghum, sudangrass, or sorghum-sudangrass hybrids.

Callisto® and the Syngenta logo trademarks of a Syngenta Group Company

©201X Syngenta

CALLISTO 1131A-S7-B - bb - 10-29-13