



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
100-1163

Date of Issuance:
FEB - 5 2004

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance:
Conditional

Name of Pesticide Product:
Suprend Herbicide

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) provided that you:

1. Submit the result of the ongoing storage and stability (830.6317) and corrosion characteristics (830.6320) studies once they are complete.
2. Submit/cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
3. Make the labeling changes listed below before you release the product for shipment.
 - a. Add the phrase "EPA Registration No. 100-1163".
 - b. Incorporate the sentence "Causes moderate eye irritation" into your Precautionary Statements, Hazards to Humans and Domestic Animals section of both your container label and label booklet.
 - c. At the beginning of the list of Personal Protective Equipment (PPE) within the Precautionary Statements, add the statements "Some of the materials that are chemical resistant to this product of listed below. If you want more options, follow the instructions for Category A on an EPA Chemical Resistance Chart.
 - d. Add the statement " This product is toxic to vascular plants and should be used strictly in accordance with the drift precautions on this label in order to minimize off-site exposures." to the Environmental Hazards section of both your container label and label booklet.

Signature of Approving Official:

Date:

2-5-04

e. Incorporate the following statements into your "Instructions to Avoid Spray Drift" section.

1. Do not apply under circumstances, where possible drift to unprotected persons, to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use, or consumption, can occur.

2. Revise the second bullet to read "Make applications when the wind velocity favors on-target product deposition. Do not spray if wind speed is 10 mph or greater or if winds are gusty. Wind speed must be measured adjacent to the application site on the upwind side immediately prior to application.

3. Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased drift. Avoid spraying during conditions of low humidity and/or high temperatures.

Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of labeling is enclosed for your records..

Suprend™

Herbicide

A herbicide for control of certain broadleaf, sedge, and grass weeds in cotton

Active Ingredients:

| | |
|---|----------------|
| Prometryn (CAS No. 7287-19-6) | 79.30% |
| 2-pyridinesulfonamide,N-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]-3-(2,2,2-trifluoroethoxy)-,monosodium salt, monohydrate; Trifloxysulfuron-sodium (ISO draft proposal) (CAS No. 199119-58-9) | 0.70% |
| Other Ingredients: | 20.00% |
| Total: | 100.00% |

KEEP OUT OF REACH OF CHILDREN.

CAUTION — PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100 - RRAG

EPA Est.

Product of France

SCP

15 pounds
Net Weight

ACCEPTED
with COMMENTS
In EPA Letter Dated:
FEB - 5 2004
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under EPA Reg. No.
100-1163

| FIRST AID | |
|--|---|
| If in eyes | <ul style="list-style-type: none">• Hold eye open and rinse slowly with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a Poison control center or doctor.• Do not give anything by mouth to an unconscious person. |
| If on skin or clothing | <ul style="list-style-type: none">• Take off contaminated clothing• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| If inhaled | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. | |
| HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call 1-800-888-8372 | |

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves, such as barrier laminate or viton

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing,
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water or rinsates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitations of Warranty and of Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Suprend should be used only in accordance with recommendations on this label or in separately published supplemental labeling recommendations for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and the restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR WEED CONTROL, AND/OR ILLEGAL RESIDUES.

STORAGE AND DISPOSAL

Pesticide Storage

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by the use according to label instructions, contact you State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

GENERAL INFORMATION

Suprend is a herbicide for postemergence control of certain broadleaf, sedge, and grass weeds in cotton. Use post-directed application methods to provide coverage of weed foliage while adjusting nozzles to minimize contact of cotton foliage with spray or drift. Over-the-top application of Suprend in cotton will result in cotton injury and may cause yield reduction.

For a list of weeds controlled, refer to Table 1 in the **Target Weeds** section of this label. The level of control resulting from application of Suprend is primarily dependent upon application rate, weed species, weed size at application, coverage of the weed (including terminal), environmental conditions and growing conditions. For best results, apply Suprend when target weeds are less than 6 inches in height. Weed control is enhanced when emerged weeds have ample soil moisture and weeds are actively growing. Growth of susceptible weeds is inhibited soon after application of Suprend. Complete plant death occurs within a few days after application, depending upon weed species and growing conditions.

HERBICIDE RESISTANCE

Suprend is a mixture of a photosynthetic inhibitor (prometryn) and an acetolactate synthase (ALS) inhibitor (trifloxysulfuron-sodium).

Suprend controls weeds by inhibiting (stopping) a biochemical process that produces certain essential amino acids necessary for plant growth and by inhibition of photosynthesis.

Certain weeds species have naturally occurring biotypes within the population that are resistant to ALS inhibiting herbicides or photosystem II inhibitors. Multiple applications of the same herbicide class in the same areas(s) applied continuously over a number of years, can lead to an increased presence of the resistant biotypes in weed populations. This may reduce the utility of herbicides for controlling specific target weeds in the population. To prevent or delay the build-up of resistant weed species biotypes, weed management programs should include the use of appropriately registered herbicides within the same or sequential years that: (1) have a different mode of action (MOA), and (2) control of the target weed. Mechanical control by tillage, cultivation, etc., or hand weeding before weeds set seed may also be helpful in reducing the build-up and spread of herbicide resistant weed biotypes.

GENERAL USE PRECAUTIONS OR RESTRICTIONS

- Do not spray over-the-top of cotton or crop injury may result.
- Apply Suprend post-directed using properly adjusted and calibrated precision applicators (e.g., hooded sprayer).
- Do not exceed a total of 2.7 lbs./A of Suprend per season from all applications. Suprend is made of two herbicidally active components, Envoke™ (trifloxysulfuron-sodium) and Caparol® (prometryn). Do not exceed a total of 0.0188 lb. active ingredient of trifloxysulfuron-sodium per acre per season resulting from all applications of Suprend or Envoke. Each pound of Suprend contains 0.007 lb. active ingredient of trifloxysulfuron-sodium and is equivalent to 0.15 oz. of Envoke. Each pound of Suprend contains 0.79 lb. active ingredient prometryn. Do not exceed 5.15 lbs. active ingredient of prometryn per acre per season resulting from all applications of Suprend or Caparol 4L, or other prometryn containing products.
- Due to reduced crop tolerance, do not apply Suprend post-directed on stripper-type cotton varieties. Post-directed applications on stripper cotton in OK, TX, NM is permitted.
- Degradation of Suprend in the soil is enhanced by soil with pH <7 and moist conditions. Application of Suprend to soils with pH >7.5 may increase the potential for rotational crop injury and may reduce rotational crop yield. Alkaline soils increase the potential for injury to rotational crops. If severe drought conditions

develop (less than 12 inches of rainfall/irrigation within the first five months following application of Suprend and/or less than 1 inch of rainfall/irrigation within the first month after application) rotational crop injury may occur. In areas where soil pH is >7.5 and/or drought occur, a field bioassay prior to planting of the rotational crop is recommended. See Table 2 for recommended crop rotation intervals following normal harvest.

- Do not tank mix Suprend with malathion, profenofos (Curacron®) or emamectin-benzoate containing insecticides (Denim™), acephate, Bidrin®, Capture®, Karate® or unacceptable cotton injury may occur.
- ALS resistant plants or photosystem II resistant plants may not be controlled by Suprend.
- Do not feed treated forage to livestock or graze treated areas.
- To avoid crop injury, do not apply to furrow-planted cotton until furrows are leveled (plowed-in).
- Do not treat cotton under stress caused by adverse environmental conditions, insect or disease damage, chemical or mechanical injury, or nutrient deficiency.
- Do not apply Suprend using liquid fertilizer as the carrier.
- Do not use in the Rio Grande Valley of TX, the Coachella Valley of AZ and CA, nor on sand and loamy sand in AZ and CA.
- Do not apply Suprend within 60 days of cotton harvest.
- Sequential applications of Suprend should be made at least 14 days apart.

MIXING PROCEDURES

MIXING INSTRUCTIONS

1. Clean the spray tank before using. If it is contaminated with other materials, mixing problems and/or clogging may occur which could cause injury to the crop or reduced performance. Prepare no more spray mixture than is required for the immediate application.
2. Fill the spray tank 1/4-1/2 full with clean water and begin agitation. Make certain that the agitation system is working properly and creates a rippling or rolling action on the water surface. Maintain agitation throughout the mixing and spraying process.
3. Add any products packaged in water-soluble film to the tank first. Allow the packets to completely dissolve and the contents of the packets to fully disperse into the mix water. **Important: Water-soluble packets must always be the first material put into the spray tank after water.** For products packaged in water-soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment has been thoroughly cleaned (see **Instructions for Cleaning Spray Equipment After Application**).
4. Add the required amount of Suprend to the spray tank while maintaining agitation. Allow the product to wet and thoroughly disperse into the mix water.
5. While maintaining agitation, continue filling the spray tank. When the tank is 3/4 full, add any tank mix partners. Add any water-dispersible granule or other dry formulation first, and allow that material to fully and uniformly disperse. Then add any emulsifiable liquid formulation.
6. Maintain agitation while adding a good nonionic surfactant with a minimum of 80% of the constituents effective as a spray adjuvant, at a rate not greater than 1qt./100 gals. spray mixture (0.25% v/v).
7. Maintain agitation while adding a nonphytotoxic COC containing 15-20% approved emulsifier at 0.5-1.0% v/v of finished spray solution.
8. Complete filling the tank maintaining sufficient agitation at all times to ensure surface action until the spray tank mixture is uniform.
9. An anti-foaming agent may be added to reduce excessive foaming, if it occurs.
10. **Do not leave spray in the spray tank without continuous agitation.** Always maintain agitation to maintain a uniform mixture in the spray tank.

11. Make only sufficient spray mixture that can be used the day in which it was mixed.

APPLICATION PROCEDURES

GROUND APPLICATION EQUIPMENT

Apply Suprend post-directed to prevent contact of the spray with cotton leaves, or crop injury may occur. Use precision application equipment so the spray is directed to the base of the cotton plant and still thoroughly covers the soil and weeds beneath the cotton plant. Apply during calm periods to avoid drift. Use leaf lifters or shields if leaf contact cannot be avoided merely by directing the spray.

For best results, apply to level, well prepared surfaces, such as relatively clod-free beds made with bed-shapers.

To help ensure accuracy, calibrate sprayer at the beginning of the season before use and recalibrate frequently. For ground application, use a minimum of 10 gallons water per acre. Higher volumes (i.e., at least 20 gals./A on a broadcast basis) should be used for severe weed infestations to ensure adequate spray coverage. Always include a nonionic surfactant spray adjuvant, approved for application to growing crops, in the spray mixture (see the **Mixing Procedures** section).

Use a pump with capacity to: (1) maintain at least 25-40 psi pressure at nozzles, and (2) provide sufficient agitation within the tank to keep product in suspension. Lower pressures may be used with extended range or drift reduction flat fan nozzles. A centrifugal pump that provides shear action for dispersing and mixing the product is recommended. The pump should provide a minimum of 20 gallons/minute/100 gallons tank size circulated through a correctly positioned sparger tube or jet agitators. If jet agitators are used, at least two agitators should be aligned on the bottom of the tank pointing toward each end. Agitation during both mixing and application is essential. Screens or strainers placed on the suction side of the pump should be 16-mesh or coarser. Use 50-mesh or coarser screens between the pump and boom, and when required, at the nozzles. Check nozzle manufacturer's recommendations.

Good weed coverage with the spray mixture is essential for optimum weed control. Observe sprayer nozzles frequently during the spraying operation to ensure that the spray pattern is uniform. Avoid spray overlaps that result in excessive rates in the overlap areas. Also, avoid application under conditions when uniform coverage cannot be obtained or when spray drift may occur (see section titled **Instructions to Avoid Spray Drift**).

Avoid all direct or indirect contact (such as spray drift) of this product with crops other than those recommended for treatment on this label, since injury may occur. Always

follow the **Instructions to Avoid Spray Drift** and the **Instructions for Cleaning Spray Equipment After Application** sections of this label.

Rainfastness: Suprend is rainfast within 3 hours after application.

Aerial Application Prohibition: Do not apply Suprend aerially.

Chemigation: Do not apply this product through any type of irrigation system.

INSTRUCTIONS TO AVOID SPRAY DRIFT

As with all crop protection products, it is important to avoid off-target movement. Do not allow spray to drift onto adjacent land or crops, as even small amounts may injure sensitive plants. When drift may be a problem, take steps to reduce spray drift, such as the following.

- Do not spray if wind speed is 10 mph or greater or if winds are gusting.
- Use extreme caution when conditions are favorable for drift (high temperatures and low relative humidity), especially when sensitive plants are located nearby. All plants not listed as target crops on this label should be considered as sensitive plants.
- If sensitive crops or other non-target plants are downwind, extreme caution must be used under all conditions. Drift from applications of this herbicide is likely to result in damage to sensitive non-target plants adjacent to the treatment site. This damage can occur at extremely low concentrations.
- Allow adequate distance between target area and non-target areas (e.g., other crops, neighboring plants, or surface water) to prevent drift onto non-target areas.
- Do not apply when a temperature inversion exists. If an inversion condition is suspected, consult with local weather services before making an application.
- Further reductions in drift can be obtained by:
 - Using nozzles that provide a uniform droplet size. Use nozzles that produce medium or coarse droplets (250-400 microns VMD) that are less prone to result in spray drift.
 - Use of precision application equipment with fenders or shields, can reduce herbicide contact with cotton foliage.
 - Recalibrate sprayer by reducing spray pressures and by increasing spray volumes to produce larger droplets when conditions favor drift.

- Applying as close to target plants as practical, i.e., a nozzle height that allows a good spray pattern for adequate coverage according to the manufacturer's recommendations, but minimizes drift potential.

SPRAY EQUIPMENT

INSTRUCTIONS FOR CLEANING SPRAY EQUIPMENT AFTER APPLICATION

Because some crops are extremely sensitive to low rates of Suprend, special attention must be given to cleaning equipment before spraying a crop other than those listed on this label. Mix only as much spray solution as needed. Immediately after spraying, clean equipment thoroughly using the following procedure.

1. Flush tank, sump, hoses, boom, and nozzles with clean water.
2. Prepare a tank cleaning solution of 1 gal. of household ammonia per 50 gals. of water. **Do not** use chlorine-based cleaners, such as Clorox®. Many commercial spray tank cleaners may be used. Refer to the publication "Clean It Up! A Guide to Cleaning Sprayers" from Syngenta, for additional information. The guide may be obtained from your local Syngenta Sales Representative.
3. When available, use a pressure washer to clean the inside of the spray tank with this solution. Take care to wash all internal parts of the tank, including the inside top surface. Completely fill the sprayer with the ammonia cleaning solution to ensure contact of the cleaning solution with all internal surfaces of the tank and plumbing. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.
4. Flush hoses, spray lines, and nozzles for at least 15 minutes with the cleaning solution.
5. Dispose of rinsate from steps 1-4 in an appropriate manner. Spray the cleaning solution on an untreated crop on which Suprend is registered, or return to a rinsate tank for later use as make-up water for spraying crops on which Suprend is registered, or use other approved disposal.
6. Repeat steps 2-5.
7. Remove nozzles, screens, and strainers and clean separately in the ammonia cleaning solution after completing the above procedures.
8. Rinse the complete spraying system with clean water.

Note: If the tank is equipped with the proper number of correctly mounted 360° tank washing nozzles that are attached to a dedicated rinsing system, less cleaning solution than a full tank may be used. Use sufficient cleaning solution to thoroughly rinse all surfaces. Start the sprayer agitation and recirculate the cleaning solution for at least 15 minutes. Flush the spray boom with the cleaning solution. Repeat the rinsing procedure 1-2 times.

CROP USE DIRECTIONS

COTTON (PICKER AND PIMA VARIETIES ONLY)

Apply Suprend as a directed spray to provide good coverage of weed canopy including weed terminals while minimizing contact of cotton leaves with spray or drift. Directed applications are generally best achieved when cotton is at least 6 inches tall. **DO NOT SPRAY OVER THE TOP OF COTTON OR CROP INJURY WILL RESULT.**

Suprend can be used post-directed on picker and pima cotton varieties in AL, AR, AZ, CA, FL, GA, KS, KY, LA, MS, NM (certain counties), NC, SC, TN, VA, the boot heel area of MO, and portions of OK and TX (east of State Highway 83 and east of I-35) including/excluding the high plains region of West Texas and Oklahoma.

Suprend can be used with all cotton tillage systems.

TARGET WEEDS

For best results, apply when target weeds are less than 6 inches in height. Apply to actively growing weeds listed below.

Table 1: Weeds Controlled or Partially Controlled With Suprend

| Weeds Controlled ^{1,2} | Latin Name | Control Level ³ | Suprend | |
|---------------------------------|--------------------------------|----------------------------|---|------------|
| | | | 1.0-1.25 lbs./A | 1.5 lbs./A |
| | | | Weed Size Ranges for Optimum Control (Inches) | |
| Barnyardgrass | <i>Echinochloa crus-galli</i> | PC | 0.25-1 | 0.25-1 |
| Bristly starbur | <i>Acanthospermum hispidum</i> | C | 1-4 | 1-6 |
| Broadleaf signalgrass | <i>Brachiaria platyphylla</i> | PC | 0.25-1 | 0.25-1 |
| Burgherkin | <i>Cucumis anguria</i> | PC/C | 1-3 | 1-4 |
| Carpetweed | <i>Mollugo verticillata</i> | C | 0.5-2 | 0.5-3 |
| Citronmelon | <i>Citrullus lanatus</i> | PC/C | 1-3 | 1-4 |
| Coffee senna | <i>Cassia occidentalis</i> | C | 1-5 | 1-6 |
| Common cocklebur | <i>Xanthium strumarium</i> | C | 1-5 | 1-6 |
| Corn, volunteer (non-IR or IT) | <i>Zea mays</i> | C | 1-5 | 1-6 |

| | | | | |
|------------------------------|--|------|-------|-------|
| Copperleaf hophornbeam | <i>Acalypha ostryifolia</i> | PC | 1-3 | 1-3 |
| Cowpea | <i>Vigna unguiculata</i> | PC/C | 1-3 | 1-4 |
| Crabgrass, large | <i>Digitaria sanguinalis</i> | PC/C | 1-5 | 1-5 |
| Crabgrass, Southern | <i>Digitaria ciliaris</i> | PC | 1-5 | 1-5 |
| Croton, tropic | <i>Croton glandulosus</i> | PC/C | 1-3 | 1-4 |
| Devil's-claw | <i>Proboscidea louisianica</i> | PC/C | 1-3 | 1-4 |
| Florida beggarweed | <i>Desmodium tortuosum</i> | C | 1-4 | 1-5 |
| Goosegrass | <i>Eleusine indica</i> | PC/C | 1-3 | 1-4 |
| Hemp sesbania | <i>Sesbania exaltata</i> | C | 1-4 | 1-5 |
| Horse purslane | <i>Trianthema portulacastrum</i> | PC | 0.5-1 | 0.5-2 |
| Horseweed | <i>Conyza canadensis</i> | PC | 1-3 | 1-4 |
| Johnsongrass (seedling) | <i>Sorghum halepense</i> | C | 1-4 | 1-6 |
| Lambsquarters, common | <i>Chenopodium album</i> | C | 0.5-2 | 0.5-3 |
| Morningglory: | | | | |
| entireleaf | <i>Ipomoea hederacea</i> var. <i>integriuscula</i> | C | 1-4 | 1-5 |
| ivyleaf | <i>Ipomoea hederacea</i> | C | 1-5 | 1-6 |
| pitted | <i>Ipomoea lacunosa</i> | C | 1-5 | 1-6 |
| red/scarlet | <i>Ipomoea coccinea</i> | C | 1-4 | 1-5 |
| smallflower | <i>Jacquemontia tamnifolia</i> | PC/C | 1-4 | 1-5 |
| tall morningglory | <i>Ipomoea purpurea</i> | C | 1-3 | 1-4 |
| Nutsedge | | | | |
| purple | <i>Cyperus rotundus</i> | PC | 2-4 | 2-4 |
| yellow | <i>Cyperus esculentus</i> | C | 2-6 | 2-6 |
| Panicum, Fall | <i>Panicum dichotomiflorum</i> | PC/C | 1-5 | 1-6 |
| Peanut, volunteer | <i>Arachis hypogea</i> | PC | 1-2 | 1-3 |
| Pennsylvania smartweed | <i>Polygonum pennsylvanicum</i> | PC/C | 1-4 | 1-6 |
| Pigweed: | | | | |
| Palmer amaranth | <i>Amaranthus palmeri</i> | PC | 1-2 | 1-2 |
| redroot pigweed | <i>Amaranthus retroflexus</i> | C | 1-4 | 1-6 |
| prostrate pigweed | <i>Amaranthus blitoides</i> | PC/C | 1-3 | 1-4 |
| smooth pigweed | <i>Amaranthus hybridus</i> | C | 1-4 | 1-5 |
| spiny amaranth | <i>Amaranthus spinosus</i> | PC | 1-3 | 1-4 |
| tumble pigweed | <i>Amaranthus albus</i> | PC/C | 1-3 | 1-4 |
| waterhemp, tall | <i>Amaranthus tuberculatus</i> | C | 1-2 | 1-2 |
| Poinsettia, wild | <i>Euphorbia heterophylla</i> | C | 0.5-2 | 0.5-3 |
| Prickly sida | <i>Sida spinosa</i> | PC | 1-4 | 1-4 |
| Ragweed: | | | | |
| giant | <i>Ambrosia trifida</i> | PC/C | 1-5 | 1-6 |
| common | <i>Ambrosia artemisiifolia</i> | C | 1-4 | 1-6 |
| Redweed | <i>Melochia corchorifolia</i> | C | 0.5-2 | 0.5-3 |
| Soybean, volunteer (Non-STS) | <i>Glycine max</i> | C | 1-3 | 1-4 |
| Sicklepod | <i>Cassia obtusifolia</i> | C | 1-4 | 1-6 |
| Spiderwort, tropical | <i>Commelina benghalensis</i> | PC/C | 1-3 | 1-4 |
| Spurred anoda | <i>Anoda cristata</i> | PC/C | 1-3 | 1-4 |
| Sunflower, common | <i>Helianthus annuus</i> | C | 1-4 | 1-5 |
| Velvetleaf | <i>Abutilon theophrasti</i> | C | 1-4 | 1-4 |

¹ ALS resistant plants or photosystem II resistant plants may be controlled by Suprend.

² For improved control of grass weeds tank mix with MSMA or glyphosate.

³ C = Control (85-100%) of weeds present at the time of application. C = may require use of higher rates or repeated applications to achieve control. PC = Partial control; partial control means significant activity but not always at a level generally acceptable for commercial weed control.

TANK MIXES WITH OTHER HERBICIDES

Suprend can be tank mixed with Dual MAGNUM®, MSMA, Aim™, diuron, linuron, Cobra®, and Caparol for post-directed applications where use of these products is also registered for use in cotton. Suprend can also be tank mixed with Buctril® for post-directed applications in BXN cotton varieties. Suprend can be tank mixed with Touchdown® (glyphosate) for post-directed use in Roundup Ready® Cotton varieties.

Suprend can be tank mixed with Touchdown (glyphosate) for use in conventional cotton varieties if applied with hooded sprayers that completely enclose the spray pattern for weed control between the rows. Adjust the hooded sprayer in raised seedbeds to ensure the rear flaps touch the ground to completely enclose the spray solution. Keep the spray and drift off cotton leaves and stems. Spray contacting cotton leaves and stems may cause cotton injury.

For all tank mixtures of Suprend refer to individual product labels for precautionary statements, restrictions, rates, and a list of weeds controlled. Spray contacting cotton leaves may cause cotton injury with many of these tank mixtures.

REPLANTING AFTER COTTON CROP FAILURE

If a cotton crop is lost (e.g., due to hail), and adequate growing season remains, and not more than 1.0 lb./A of Suprend has been applied, cotton may be replanted 30 or more days after the Suprend application, or 14 or more days after the first significant rainfall event (≥ 0.5 inches) following the Suprend application. If Suprend has been applied at > 1.0 lb./A, cotton may be replanted 30 days after the first significant rainfall event (≥ 0.5 inches) following the Suprend application.

ROTATIONAL CROP RESTRICTIONS FOLLOWING COTTON

The crops listed below may be planted at, or after, the time interval specified from the last application of Suprend. Table 3 shows the amount of trifloxysulfuron-sodium from various application rates of Suprend and Envoke herbicides. Do not exceed a cumulative total of 0.0188 lb. a.i./A of trifloxysulfuron-sodium per season resulting from all applications of Suprend and Envoke.

Suprend contains Caparol, with the active ingredient prometryn. Do not exceed a total of 5.15 lbs. a.i./A prometryn per season resulting from all applications of Suprend or prometryn containing products.

If another herbicide with a longer rotational interval was used, follow the longer rotational limitation.

Table 2. Rotational Crop Restrictions

| Rotational Crop | Months |
|----------------------------|--------|
| Bell pepper (transplanted) | 12* |
| Cotton | 7 |
| Corn, field | 7 |
| Corn, sweet | 7 |
| Grain sorghum | 7 |
| Parsley | 12* |
| Peanut | 7 |
| Potato, Irish | 12* |
| Radish | 12* |
| Rice | 7 |
| Soybean | 7 |
| Sugarcane | 1 |
| Tobacco (transplanted) | 7 |
| Tomato (transplanted) | 3 |
| Wheat, winter | 3 |
| All other crops | 18* |

* Field bioassay.

Table 3: Envoke/trifloxysulfuron and Caparol/prometryn Dose Conversion

| Suprend lbs. Product/A | Envoke oz. Product/A | Trifloxysulfuron lb. a.i./A | Caparol 4L pts./A | Prometryn lbs. a.i./A |
|---------------------------|-------------------------|--------------------------------|----------------------|--------------------------|
| 1.00 | 0.15 | 0.007 | 1.58 | 0.79 |
| 1.25 | 0.19 | 0.00875 | 1.97 | 0.98 |
| 1.50 | 0.23 | 0.0105 | 2.37 | 1.18 |
| 2.01 | 0.30 | 0.0141 | 3.18 | 1.59 |
| 2.69 | 0.40 | 0.0188 | 4.25 | 2.12 |

Caparol®, Curacron®, Denim™, Dual MAGNUM®, Envoke™, Karate®, Suprend™ Touchdown®, and the Syngenta logo are trademarks of a Syngenta Group Company U.S. Patent Nos. xxx, xxx, xxx, xxx

Aim™ and Capture® trademarks of FMC Corporation

Bidrin® trademark of Amvac Chemical Corporation

Buctril® trademark of Bayer CropScience

Clorox® trademark of Clorox Corporation

Cobra® is a trademark of Valent Agricultural Products

©2003 Syngenta

For non-emergency, (e.g. current product information), call
Syngenta Crop Protection at 1-800-334-9481.

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP

SUPREND – pd – 06/13/03
000100-xxxxx.20030616.SUPRENDLABEL .pdf

(CONTAINER LABEL)

Suprend™

Herbicide

A herbicide for control of certain broadleaf, sedge, and grass weeds in cotton

Active Ingredients:

| | |
|---|--------|
| Prometryn (CAS No. 7287-19-6) | 79.30% |
| 2-pyridinesulfonamide,N-[[[4,6-dimethoxy-2-pyrimidinyl)amino] carbonyl]-3-(2,2,2-trifluoroethoxy)-,monosodium salt, monohydrate; Trifloxysulfuron-sodium (ISO draft proposal) (CAS No. 199119-58-9) | 0.70% |

| | |
|--------------------|--------|
| Other Ingredients: | 20.00% |
|--------------------|--------|

| | |
|--------|---------|
| Total: | 100.00% |
|--------|---------|

KEEP OUT OF REACH OF CHILDREN.

CAUTION— PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-

EPA Est.

Product of France

15 pounds

Net Weight

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling.

| FIRST AID | |
|--|--|
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. | |
| <p align="center">HOT LINE NUMBER</p> <p align="center">For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call 1-800-888-8372</p> | |

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water or rinsates. . Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night

Chemigation

Do not apply this product through any type of irrigation system.

Suprend™ and the Syngenta logo are trademarks of a Syngenta Group Company
U.S. Patent Nos. xxx,xxx,xxx

©2003 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP

SUPREND – pd – 06/13/03
000100-xxxx.20030616.SUPRENDLABEL .pdf