




QUILT™ FUNGICIDE

EPA Reg. No. 100-1178

SUPPLEMENTAL LABELING



For Disease Control in Cereals

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This supplemental label must be in possession of the user at the time of application.

A C C E P T E D
JUN 23 2005
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 100-1178

DIRECTIONS FOR USE


Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Cereals Barley Wheat Triticale	Early season suppression of: Powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nordorum</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>)	7 - 14	Apply Quilt in the spring for suppression of early season diseases. Applications may be made no closer than a 14-day interval. Quilt may be applied up to Feekes growth stage 9. Quilt may be applied by ground, air or chemigation.
	Control of Leaf Diseases: Rust (<i>Puccinia</i> spp.) Powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nordorum</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>) Helminthosporium leaf blight (<i>Drechslera tritici-repentis</i>) Spot Blotch (<i>Bipolaris sorokiniana</i>) Barley Scald (<i>Rhynchosporium secalis</i>) Barley stripe (<i>Pyrenophora graminea</i>) Net blotch (<i>Pyrenophora teres</i>) Kernel blight (<i>Alternaria</i> spp.)	10.5 - 14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Quilt is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl. oz./acre may be applied. Applications may be made no closer than a 14-day interval. Quilt can be applied until the ligule of the flag leaf emerges (Feekes growth stage 9) on barley, rye, and triticale. For wheat only, Quilt can be applied through full head emergence (Feekes growth stage 10.5). Do not apply after this stage to avoid possible illegal residues. Quilt may be applied by ground, air or chemigation.
	Foot Rot/Eyespot (<i>Tapesia</i> spp.)	14	Apply full rate of Quilt plus half the rate recommended of other EPA registered fungicides such as Topsin® M. Apply at tillering but before elongation has occurred.
Specific Use Restrictions: Do not apply more than 2 applications/A/year. Do not apply within 45 days of harvest for grain and straw. Do not harvest wheat for forage. Do not graze or feed livestock treated forage or cut green crop for hay or silage. Do not apply more than 20.5 fl. oz./A/season of Quilt. Do not apply more than 0.167 lb ai propiconazole containing products/A/year. Do not apply more than 0.4 lb ai azoxystrobin containing products/A/ year. Under certain environmental conditions, tankmixes of Quilt plus herbicides and/or fertilizers may cause crop injury in barley, triticale and wheat.			

Trademark of Syngenta



© 2005 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com



SCP 1178A-SX XXXX

Qui 1178A-SX XXXX AMD to EPA nsp 06/20/05